concentrated solutions of ptomaines, which are highly poisonous. Most rational and safe is the treatment of this condition by irrigating the uterus with sterilized water at 104° F. The heat stimulates the uterus to contract and expel poisonous materials, while the water dilutes the ptomaine solutions until they are innocuous.

GYNECOLOGY.

UNDER THE CHARGE OF
HENRY C. COE, M.D., M.R.C.S.,
OF NEW YORK.

THE ACTION OF ANTISEPTICS UPON THE PERITONEUM.

DELBET, GRANDMAISON, and BRESSET (*Annales de Gynécologie et d'Obstétrique*, 1891) as a result of their united experiments and observations, arrive at the following conclusions:

1. The strong antiseptics, such as carbolic and salicylic acids, corrosive sublimate and bichloride of mercury, present more disadvantages than advantages, since they irritate a healthy peritoneum and thus favor the formation of adhesions.
2. The most useful solutions to be employed in laparotomy are one of sodium chloride (six or seven parts to one thousand), and of boric acid (three parts to one hundred).
3. Iodoform and salol exert but little action upon the peritoneum, and it is a question if they have any antiseptic effect.

URETERO-VAGINAL FISTULA.

ALTEN (Inaugural-dissertation, Munich, 1889; abstract in *Centralblatt für Gynäkologie*, 1890) analyzes thirty-five cases, the histories of three of which were recorded at Winckel's clinic, and finds that in the majority the condition could be traced to a previous severe instrumental delivery or operation, especially vaginal hysterectomy and the opening of pelvic abscesses. Ureteral fistulae are to be distinguished etiologically as congenital, traumatic, and spontaneous. In twenty cases an operation was performed, only ten of which were successful.

TORSION OF THE PEDICLE IN OVARIAN CYSTS.

KÜSTNER (*Centralblatt für Gynäkologie*, 1891) has made a number of observations bearing on this interesting subject, and differs from other authors in his explanation of the etiology. In his last thirty-six ovariectomies he found in fourteen cases (38.8 per cent.) torsion of the pedicle up to at least 180°. Rokitansky records 13 per cent., Thornton 9.5 per cent., Olshausen 6.3 per cent., and Hinrutz 23.2 per cent. He explains the greater frequency of this complication in the Dorpat clinic by the fact that in most

of the cases inflammatory adhesions existed. In the majority of the cysts of the right ovary the torsion was toward the *left*, in those of the left ovary toward the *right*. His explanation is as follows: As soon as the tumor rises out of the pelvis and comes in contact with the abdominal wall it tends to fall forward, displacing the uterus backward. The ovarian ligament is now put on the stretch and crosses the tube, so that when the tumor lies in contact with the abdominal wall there is really a torsion of its pedicle to the extent of 90° , which is usually overlooked at the time of operation. If the pedicle is small, and the abdominal wall relaxed, changes in the centre of gravity of the tumor are readily produced by changes in the position of the patient; when she lies upon her left side it tends to rotate from right to left, since the infundibulo-pelvic ligament (which is its principal attachment) extends from the posterior wall of the pelvis anteriorly from right to left. The reverse is the case when she lies on her right side. This torsion is probably only temporary. After the neoplasm occupies the abdominal cavity the pressure of the intestines upon its posterior surface during peristalsis tends to increase the existing torsion; this pressure on the right side is exerted most upon the left side of the tumor and *vice versa*. This theory was supported by a case of double ovariectomy of the writer's, in which the pedicle of the left tumor was twisted 180° to the right, and that of the right to the left.

THE TREATMENT OF UTERINE FIBROIDS ACCORDING TO APOSTOLI'S METHOD.

Two recent communications by well-known and successful abdominal surgeons are of considerable interest from the different conclusions at which the writers arrive.

KEITH (reprint from *British Medical Journal*) reports further progress in the electrical treatment of fibro-myomata. During the last three years and a half he has operated only upon fibro-cystic tumors, and believes that with care there ought never to be any doubt regarding the selection of proper cases. The best results are obtained in the case of small bleeding tumors of recent origin, which require shorter treatment than the old, large ones. He summarizes as follows: "This treatment *almost always* relieves pain. It *almost always* brings about diminution of the tumor—sometimes rapidly. It *almost always* stops hemorrhage—sometimes rapidly. The results are *almost always* permanent, and the growth of the tumor, if it be not lessened, is stopped. The general health is immensely improved. By '*almost always*,' I mean nineteen cases out of every twenty."

HOMANS (*Boston Medical and Surgical Journal*, 1891) records the results of his experience with the electrical treatment of uterine fibroids since December, 1887, including thirty-four cases, the subsequent histories of which were carefully traced. Of these the principal results noted were relief of pain and hemorrhage, and improvement in the general health. In ten cases the tumor continued to grow, in sixteen no change was noted, and in only two was there a marked diminution in its size; in the latter the menopause appeared soon after the cessation of the treatment. In general, he regards the treatment as merely symptomatic and rather unsatis-

factory. It is not entirely devoid of danger, and is not capable of successful application by the general practitioner. "It may be inferred," he adds, after citing fourteen successful laparo-hysterectomies out of fifteen cases, "that I have found nothing sufficiently curative in electrolysis to make me lay down my knife and ever take it up."

INTESTINAL OBSTRUCTION AFTER OVARIOTOMY.

ANDERSON and HANDFIELD JONES (*Lancet*, 1890) report a case in which obstruction occurred seventeen days after operation. The abdomen was reopened three days later and a strong adhesion was found, attaching a coil of small intestine to the anterior abdominal wall. It was separated, the collapsed coil dilated, and the obstruction was overcome. The patient continued for a short time to have fecal vomiting, but eventually recovered. The writers distinguish two forms of obstruction following abdominal section, the first being almost immediate, the second developing after convalescence. In the first the symptoms are masked by those of peritonitis, and a secondary operation is usually fatal; in the second the diagnosis is easier and an operation promises relief. A second case is reported in which the obstruction was relieved twenty-five days after the primary operation.

COLLAS (Thèse de Paris, 1891) has collected twenty-three cases, eighteen of which followed ovariectomy. In eight cases a secondary operation was performed, four being successful. In fifteen the actual condition was only revealed at the autopsy. The symptoms of obstruction usually developed within ten days after operation, but in one instance six years elapsed. Peritonitic adhesions were the usual factors. The symptoms appeared suddenly and were not relieved by palliative treatment. The writer's deduction is that laparotomy offers the only positive means of relief.

THE TREATMENT OF OSTEOMALACIA BY CASTRATION.

HOFMEIER (*Centralblatt für Gynäkologie*, No. 12, 1891) reports the case of a virgin, aged thirty years, who had osteomalacia of three years' standing, the disease being progressive at the time of operation, as shown by the presence of severe pains in the pelvic bones, inability to walk, and marked pelvic deformity. Menstruation regular and painless. Four weeks after removal of the ovaries the patient could walk without assistance, and the pains were much less severe. She received cod-liver oil and peptonate of iron; six weeks later she felt quite well, had no pain, and could walk a long distance. The pelvic organs were normal. The ovaries were atrophied, as in a woman after the menopause, and presented a similar appearance microscopically, only a few ovisacs being seen. The case was interesting, not only because the patient was a virgin (hence pregnancy could not be regarded as an etiological factor), and there was no disturbance of menstruation, or evidence of pelvic congestion.

As regards the effects of castration under such circumstances the writer admits, with Fehling, that it is impossible to give a satisfactory explanation. It may be due to some reflex action upon the vasomotor nerves supplying the nutrient vessels of the pelvic bones, the disease itself being regarded as a tropho-neurosis of the bones directly dependent upon ovarian activity. This

theory receives additional support from a similar case reported by Truzzi. Fehling has collected twenty other cases of castration for osteomalacia, in none of which was there a failure on the part of the operator to secure at least a temporary benefit.

PARALYSIS FOLLOWING SUBCUTANEOUS INJECTIONS OF ETHER.

EBERHART (*Centralblatt für Gynäkologie*, No. 12, 1891) reports the following interesting case, which is of considerable interest to laparotomists, since ether is frequently used hypodermatically as a cardiac stimulant in cases of collapse during abdominal section.

In the course of an operation upon the cervix and perineum Eberhart's patient suddenly collapsed, whereupon two injections of ether were given, one on the anterior, and the other on the posterior aspect of the right forearm. The following day the patient had extensor paralysis of the middle, ring, and little fingers of the right hand, which disappeared in six weeks under faradization. It was inferred that a branch of the radial had been affected. Similar cases have been reported by Heymann, Mendel, and Remák; the latter attributes the paralysis to the direct irritation of the peripheral nerve by the ether. [We have long held the opinion that ether, although one of the most rapidly-acting cardiac stimulants, is too irritating for hypodermic use, even in emergencies in which it is invariably so employed. Even if paralysis does not result, persistent localized pain, paresthesia, dermatitis, and abscesses are not infrequent results. We have found that deep injections of camphorated oil (one part to four), as recommended by the writer, were equally efficacious, and much less irritating.—ED.]

CYSTOPEXY.

TUFFIER (*Centralblatt für Gynäkologie*, No. 15, 1891) reported at the Société de Chirurgie several cases of cystocele treated by cystopexy, the technique of the operation being as follows: The bladder was first dilated with five ounces of boric acid solution, an incision two and one-half inches long was made in the hypogastrium, and a couple of catgut sutures were passed through the edges of the wound and through the superficial muscular layer of the anterior wall of the bladder. The abdominal incision was closed in the usual manner, and the patient was catheterized during the first week. In another case the bladder was suspended by four silk sutures, which included the serosa only, and were not passed through the skin.

Tuffier had operated thrice successfully by the extra-peritoneal method (as in epicystotomy), in one instance performing hysterorrhaphy, cystopexy, and anterior colporrhaphy upon the same patient. [This operation impresses us as being rather heroic treatment for a minor ailment. In the discussion of the paper Verneuil, Pozzi, and Richelot took the same ground. The same procedure was suggested and carried out by Dr. Henry Byford, of Chicago, at least two years ago.—ED.]

THE RELATION OF HEART LESIONS TO PELVIC DISEASE.

NEVUS (*Centralblatt für Gynäkologie*, No. 15, 1891) calls attention to the frequent coincidence of cardiac and pelvic disease, mitral stenosis being

most frequently observed. Since the pelvic symptoms usually absorb most attention, the cardiac complication is often overlooked. Among 419 women who were examined, 110 (26 per cent.) had mitral disease; of these, 55 per cent. had erosions of the cervix, 43 per cent. prolapse of the ovaries, 38 per cent. displacements of the uterus, and 43 per cent. menorrhagia or metrorrhagia. [We are forced to believe that the writer exaggerates the frequency of heart lesions in women with pelvic troubles, and would offer in support of this criticism the fact that in institutions like the New York State Woman's Hospital, in which the condition of the heart is carefully noted before operations, no such proportion has been observed. If this fact were well established, the number of gynecological operations, both in private and hospital practice, would be diminished at least one-half.—ED.].

VESICAL PAIN AS A SYMPTOM OF PYONEPHROSIS.

GUYON (*Centralblatt für Gynäkologie*, No. 15, 1891) reports two cases bearing upon this question. In one the extreme tenderness of the bladder in making pressure over the fundus, and in introducing a sound, the amount of vesical tenesmus, and the purulent character of the urine, seemed to justify the inference that the condition was primarily cystitis. Cystotomy was performed, and the pain was promptly relieved. The kidneys, which had previously been so large that they could be palpated, and were quite sensitive to pressure, rapidly diminished in size.

In the second case, in which the presence of enlarged kidneys, fever, and pyuria pointed to an existing pyonephrosis, cystotomy was followed by a disappearance of the symptoms, although the kidneys still remained large. The writer's deduction is that, as renal affections may be secondary to vesical inflammation, so the former may be relieved by treating the latter. [The writer's conclusions would be more valuable if his premises were more trustworthy. It is not clear that he possessed sufficient evidence upon which to base a reliable diagnosis of pyonephrosis. It would seem as if the disease in both instances was situated *below* the kidneys. Under similar conditions we recently made an explorative lumbar incision, and punctured the supposed pyonephrotic kidney, but found no pus. Subsequent observations showed that the symptoms were due to metritis. The recommendation to make an artificial vesico-vaginal fistula is none the less a good one in a doubtful case.—ED.].

EXTIRPATIO UTERI SACRALIS.

Under this term CZERNY (*Centralblatt für Gynäkologie*, No. 16, 1891) describes the operation of removing the uterus after resecting the sacrum, as practised by him in three cases. His technique is as follows: The patient being on the left side, an incision is made along either side of the sacrum. The coccyx, fifth sacral vertebra, and a portion of the fourth, are removed, osteoplastic resection not being recommended, since it retards the healing process. The peritoneum is divided, the coccyx is drawn downward, and the broad ligaments are ligated. The bladder is dissected off from above. The peritoneal wound is finally closed with catgut. The operation consumes at least two hours, and the healing process is slow; its advantages are not great, and it is

only to be preferred in complicated cases—stenosis of the vagina, disease of the parametric tissues, etc.

MÜLLER (*Correspondenzblatt für Schweiz. Aerzte*, Bd. xxi., 1891) has performed the operation with success in three instances. The disadvantages of the operation are the necessary external incision and the preliminary separation of the rectum, and slow convalescence due to the size of the wound. It is sufficient to make an incision from the lower end of the sacrum to a point half an inch from the anus; it is only necessary to remove the coccyx, which will leave a much smaller wound. The advantages of the method are the perfect view of the field of operation, even in complicated cases, the possibility of controlling hemorrhage, and of avoiding injury to the uterus; finally, the peritoneal cavity is more easily and exactly closed. It is particularly applicable to cases of advanced carcinoma.

UNIQUE CASE OF FOREIGN BODY IN THE VAGINA.

SZIGETHY (*Orvosi Hetilap*, No. 52, 1890) reports the case of a woman, aged seventy-five years, who, thirty years before, in order to support a prolapsed uterus, introduced into the vagina a ball of string previously dipped in wax. This entirely relieved her, and was worn without discomfort, so that she forgot its existence, when it was forced out of place by a violent effort. When extracted, with some difficulty, it measured seven inches in circumference, and was covered with mucus, but otherwise unchanged.

PÆDIATRICS.

UNDER THE CHARGE OF
JOHN M. KEATING, M.D.,
OF PHILADELPHIA;

A. F. CURRIER, M.D.,
OF NEW YORK,

AND

W. A. EDWARDS, M.D.,
OF SAN DIEGO, CAL.

ASTHMA OF CHILDREN.

R. BLACHE (*L'Union Médicale*, 1890, No. 133) defines asthma as a bulbous disease, in which attacks are produced by impressive irritations of the vagus or peripheral nerves, particularly the trigeminus; the reflex action manifests itself by successive or simultaneous spasms of all the intrinsic inspiratory muscles, intercostals, aculænx, trapezii, etc., and by a tetaniform contraction of the diaphragm.

Blache thus classifies the disease into three forms: Pneumo-bulbous asthma, essential or nervous asthma, emphysematous or alveolar asthma, catarrhal or bronchitic asthma.

Nervous asthma governs all asthmatical pathology of children. A class of asthma in infantile pathology which holds a place whose importance in-