cervix. The uterine sound was passed into each cervix and passed the same distance on each side—viz., four and a half inches. The right cervix (A) was now further dilated by means of Hegar's dilators till the finger could be passed through it into the uterus. The submucous fibroid (D), the size of a large walnut, was felt attached at the fundus and to the right side. It was seized with polypus forceps and twisted till it seemed loose, but it could not be brought away till the cervix was further dilated up to the size of No. 24 Hegar. The condition of the uterus was then examined more closely. The uterine sound was passed into the left cervix and the finger into the cervix on the right

A, Right cervix. B, Left cervix. C, Common cavity for the body of the uterus. D, Submucous fibroid. E, Remains of septum just above the internal os on each side. F, Falciform fold of vaginal mucous membrane. V, Vagina.

side through which the fibroid tumour had just been removed. The condition of things figured in the illustration was then found—that is, the finger felt the sound in the cavity (C) of the body of the uterus, showing that there was a single cavity for the body of the uterus into which led the two separate cervical canals. There was a sort of spur (E) felt a little above the level of the internal os. Bimanually the upper part of the uterus was felt to be slightly irregular, there being three distinct prominences, as figured in the illustration, most probably due to the presence of sub-peritoneal fibroids. The patient made the uneventful recovery usual nowadays after operations of this nature. The notes of this case were taken by Dr. G. H. Cowen, resident accoucheur and Mr. P. K. Nix, clinical clerk, and revised by myself.

Remarks.—Cases of double uterus are very rare. In nearly eleven years' experience at the London Hospital I have only seen two examples, one the case just recorded in which there was a submucous fibroid tumour, and the other a case of

double uterus with double hæmatometra and complete absence of the vagina, on which I operated successfully some weeks ago, and which I propose to record fully on another occasion. Probably a case of double uterus of this variety affected with fibroid tumours has been met with before, but I have not come across any record of such a case, and if cases of double uterus are admittedly rare a case of double uterus complicated with fibroid tumour must be a great deal rarer. It will be observed that there was only one cavity for the body of the uterus, the septum originally present there having disappeared; and that the double condition persisted from a little above the level of the internal os downwards on each side. More generally in cases of double uterus it is the upper part of the septum between the uteri that persists, while that corresponding to the cervical region disappears.

Harley-street, W.

SELF-POISONING IN HEART DISEASE: ITS RELATION TO THE SCHOTT METHODS OF TREATMENT.¹

BY W. BEZLY THORNE, M.D. ST. AND.

BEFORE entering in detail on the subject it may be well, as a preliminary step, to endeavour to define what it is proposed to include in the terms "self-poisoning" and "heart disease." By "self-poisoning" I refer, for the purpose of the present paper, to that particular dyscrasia which is due to depraved digestion, a polluted blood-stream, abnormal metabolism, and inadequate excretion. With regard to "heart disease," I may be permitted to say that it would be difficult to find, in the whole range of medical terminology, an expression of more indefinite meaning. In the lay mind it is associated with such grave valvular lesions as expose the subject to an insecurity of life which is without reprieve or remedy. Not infrequently it is employed by members of the medical profession, possibly in deference to the popular interpretation, in a sense scarcely less restricted, and thus it may happen that the subject of a valvular affection, in itself so insignificant as to involve no material increase of life risk, is pronounced to be afflicted with "heart disease," while another, in whom processes more insidious and more dangerous to life are making unimpeded advance, is said to have "nothing organically wrong." I make, therefore, no further excuse for employing the word "disease" to designate such conditions as involve deviations from those which may properly be regarded as pertaining to an organ in a state of average health. I shall perhaps make my meaning clearer if I state, without further preface, that of some five hundred cases presenting impairment of health, in which abnormal conditions of the heart and bloodvessels have played a leading part, I have recorded in 90 per cent. collateral symptoms which point to serious and prolonged contamination of the blood-stream. The subjects may be fat or lean, florid or pale, but underlying their facial aspect they one and all present, in greater or less degree, that feature which Eustace Smith long ago, in connexion with atrophic diseases of children, denominated the "café-au-lait complexion." On inquiry it is elicited that their stools are not infrequently, perhaps habitually, deficient in biliary colouring matter. The fæces are generally found to emit an acid odour and, when not too dry to respond to the test, to redden blue litmus paper. The urine on cooling deposits either amorphous urates or crystals of uric acid; the latter are occasionally so small and so colourless as to be not readily apparent. The distal articular extremity of the index metacarpal bone presents on the radial side an osteoid deposit of the size of a split pea or of greater size and prominence. Other bones, especially the external aspect of the carpal extremity of the ulna, are found to be similarly affected. The arteries are small, and in and after middle age are found to be in a state of undue tension and not infrequently tortuous and otherwise obviously atheromatous. Some give a history of podagra and of other forms of gout and others have suffered one or more accesses of acute rheumatism; the greater part acknowledge to an exceptional liability to so-called "rheumatic" pains. Where

¹ An address delivered before the Farnington Medical Society.

evidence on the subject is forthcoming it is often found that difficulties of feeding have added to the care of early infant life.

I have already said that one feature which is common to these cases is the occasional or habitual deficiency of biliary colouring matter in the evacuations; and here, I think, we have the clue to a primary and fundamental pathological condition. In treating them we have to take into consideration a chronic, though in some cases frequently latent, catarrh of at least one portion of the digestive tract, involving more or less obstruction of the orifice of the common bile duct, and, therefore, a not infrequent and perhaps habitual deficiency of the important pancreatic and biliary juices supplied to the passing products of a gastric digestion which itself has been vitiated by the teeming bacteria of fermentation which proliferate in the depraved mucus with which it is always, in greater or less degree, encumbered. Speaking as a clinical observer, it would be difficult for me to say how far in most cases the catarrhal condition extends along the course of the small intestine; but certain it is that in some, and those by far the more grave, it involves the colon. In such we find that, instead of normally formed or constipated faecal evacuations, one or more, sometimes several, fragmentary or liquid stools are daily evacuated. What, then, is the chemico-physiological condition with which we are confronted as regards the digestive functions in a large proportion of the subjects of cardio-vascular disease? Reasonable inference and clinical observation alike point to an excess of gastro-intestinal fermentation, and the conversion of more or less of the ingested food into products which, transmitted into the blood-stream, exercise a definitely toxic influence on the tissues of the organs of circulation and on the structures which it should supply throughout the body with material for nutrition and repair. If we proceed to inquire what are the effects produced on the cardio-vascular system we may expect to meet with the following train of results: long-sustained contraction of the arterial channels, more especially of the remoter ramifications, inducing sooner or later structural change; increased peripheral resistance and later an excess of labour thrown on the left ventricle; a slowly diminishing coronary supply of nutritive blood; the effects of impeded circulation creeping backwards through the pulmonary circuit to the chambers of the right side; veins gorged with a languid stream which is charged with an undue proportion of depraved and toxic constituents; a steadily increasing transfer of the vital stream from the area of life on the arterial side to the venous store of used and unregenerated blood; and venous congestion of the great viscera, including special organs and secreting glands, with consecutive impairment of function, and more especially of the pelvic organs, involving in the male prostatic congestion, in the female a similar condition of the uterus and its appendages, and in both the hæmorrhoidal system of vessels with the well-known attendant consequences. Indeed, it would be impossible to refer in detail to the multifarious consequences of the condition to which I have alluded without engaging in an unnecessary enumeration of an almost endless series of morbid conditions with which we are all only too familiar as following on deviations of the circulatory system from a condition of health and efficiency. Nevertheless, there are some which claim special attention. I will make no more than a passing allusion to the malnutrition of the nerve centres and the languor of mind and body and depression of spirits which of necessity ensue, adding heavily to the general burden of life, and pass on to refer to changes specially affecting cardio-vascular structures themselves.

I have drawn attention to the fact that the general diminution of the arterial lumen must affect the coronary circulation no less than that of other parts. It is no less obvious that the prolonged strain thrown on the heart walls must tend to put them on the stretch to such a degree as further to impede the flow of blood through their nutrient vessels. But the evil series does not stop at this point. Not only do the vasa vasorum and the interstitial nutrient spaces become involved in their turn, but the dyscrasic condition of the blood has a direct influence on the now vulnerable elements of the vascular coats. Deterioration and degeneration of structure must inevitably follow. In view of such a combination of morbid influences it is not difficult to account for the changes which are included under the denomination of atheroma, and for the several forms of chronic and acute inflammation to which the heart, the arteries, and the veins are subject. More especially would

I draw attention to the processes which lead to aneurysm of the heart and great vessels, and to deforming and ulcerative lesions of the valves, as well as to endocardial inflammation, with their often formidable consequences; and I cannot refrain from pausing to ask whether the affection known as rheumatic fever, the cause and incidence of which have been the subject of so much learned investigation, may not be an acute and culminating incident, determined by relatively adventitious circumstances, in a long protracted course of self-poisoning in which uric acid has played an important part. If such be the case the frequent and occasionally exclusive incidence of that disease on the structures of the heart would be accounted for.

Before passing from the enumeration of the changes which the dyscrasia under consideration is capable of inducing in the organs of circulation I must not omit to allude to a condition which has been described by Dr. Savill under the name of "Arterial Hypermyotrophy."² He has pointed out that in a number of aged people death ensues on a slowly progressive failure of the functions of organic life, which can only be accounted for by the general occlusion of the arterioles, leading in its turn to slow degeneration of the tissues which they supply, and, further, that microscopic examination of those small vessels actually reveals changes in the muscular tunic which are due to a hypertrophy of its constituent fibres, which tends at one and the same time to obliterate the vascular lumen and to proceed to a degeneration of the redundant muscular tissue essentially different from that which is characteristic of atheroma. I cannot help suspecting that the long-sustained effort of contraction which is imposed on those tissues by the transmission of toxic blood may be mainly responsible for the morbid process to which Dr. Savill has called attention. I do not pretend to define exactly the identity and the chemical constitution of the noxious substances which are capable of working such widespread and multifarious mischief in the human economy. Urea and its derivatives are doubtless among the chief offenders, but there is reason to believe that they do not stand alone. I am partly led to that conclusion by the fact that the urine of the subjects in question, even when freed by deposition from a great part of its lithic constituents, possesses a notable power of decolourising Fehling's test solution; and when it is borne in mind that that special quality often precedes in point of time the actual occurrence of glycosuria and follows its subsidence, that pancreatic inhibition and duodenal indigestion are believed to influence the incidence of the diabetic condition, and that the greater part of glycosuric subjects present weak and dilated hearts, I cannot resist the conviction that in the direction which I have indicated there lies before the investigator a field of research of almost fascinating interest.

Returning now to the subject of our special consideration, we have before us a condition presenting the following principal factors: a catarrhal condition of one or more regions of the digestive tract, an excessive and depraved process of bacterial fermentation inducing putrefactive changes and hyper-acidity, the passage of toxins from the gastro-intestinal system into the blood-stream, and consecutive functional and organic changes in the tissues of the heart and bloodvessels. If, therefore, those several conditions form in many cases of disease consistent parts of a whole, and stand to each other in the relation of cause and effect, it remains to inquire how they can best be met by the various therapeutic resources at present available. It would seem to be obvious that the objects to be kept in view are the relief and prevention of catarrhal conditions, the correction of bacterial fermentation, the avoidance of fermentable articles of diet, intestinal antiseptics, the promotion of adequate elimination, the opening up of the arterial channels, and the repair of cardiac and vascular structures. If those ends be attained the purification of the blood-stream and the establishment of a better habit of metabolism must follow as a natural consequence. That pharmaceutical agents may be employed with great advantage will at once be admitted by the most ardent advocates of physical treatment, and among those which are calculated to exercise a beneficial influence on the digestive mucous surfaces preparations of bismuth stand first in order. My experience leads me to place most reliance on the salicylate as possessing special bactericidal and antiseptic properties and being

² Vide paper read in the Section of Pathology of the British Medical Association meeting at Bournemouth, 1891.

peculiarly well adapted to meet the exigencies of the numerous cases in which uric acid plays a leading part. Its efficacy is often enhanced by combination with small doses of one of the alkaline iodides. Next in order come mercury and its salts. Grey powder, in doses of a fourth of a grain, administered in due relation to meals, is well adapted to those cases in which there is a tendency to a too free action of the bowel, while still smaller doses of calomel, and the solution of the bichloride, or fractional doses of the biniodide, are indicated where, the catarrh being limited to the gastro-duodenal region, the evacuations are either normally formed or hard and scybalous. Guaiacol, resorcin, and thiocamf, enclosed in gelatine capsules and taken with a drink of warm water from two and a half to three hours after meals, do much to correct intestinal putrefaction, especially if there be loose and offensive stools. I have already stated that such cases are among the more grave. The colon is endowed with great powers of absorption, and if it be allowed to transmit liquid or semi-liquid matter highly charged with toxins serious contamination of the blood-stream must ensue. In long-standing cases chronic desquamative catarrh, and even ulceration, present formidable obstacles to successful treatment. I have not met with one such in which there has not been a weak, dilated, and presumably degenerated heart, with or without valvular lesion and more or less tachycardia. The saline laxatives are remedies of unquestionable value, but they are double-edged weapons; and if their exhibition be unduly prolonged they tend to promote the catarrh which it should be one of our main objects to relieve and forestall. Generally speaking, they are of greatest use where the catarrh is mainly limited to the gastro-duodenal tract; but, on the other hand, they are not only tolerated but often prove to be actually curative in colic cases, if taken at the spring from which they rise as natural mineral waters. Thus ingested they exercise a powerful influence on the kidneys, the great eliminators of waste and depraved products, and possess advantages over pharmaceutically prepared remedies which it would be difficult either to explain or exaggerate. A measure of primary importance, in the prevention of fermentive decomposition, is the elimination from the dietary of those substances which most readily undergo fermentation—namely, the carbohydrates and hydrocarbons. That precaution observed, no fear need be entertained of aggravating the uric acid diathesis by means of nitrogenous food. Uric acid in many instances readily disappears under the influence of a diet of diabetic severity. Of all agents, however, water is perhaps the most efficacious and at the same time the most harmless. It should be taken at a moderate heat, in quantities of from one-third to half a pint, when gastric digestion has been mainly completed and the chyme is actively passing through the pylorus.

Such measures and means as I have briefly enumerated may generally be relied on to effect a notable purification of the blood-stream and to promote the elimination of noxious substances from the tissues. Even if those objects be only partially attained diminution of arterial tension is rapidly ensured, and, peripheral resistance being relieved, not only is the work of the heart reduced, but the processes of tissue poisoning and of consequent degeneration are checked in proportion. But of all the therapeutic agencies which can be called into requisition for the relief of the class of cases under consideration, the methods of treatment which owe their origin and elaboration to August and Theodor Schott stand for every reason first and foremost. So much has recently been said and written on the subject that it is now scarcely necessary to enter on a detailed description of that mode of treatment. It consists, as is well known, of the administration of mineral baths and of physical exercises. Nevertheless, as some misapprehension appears to exist I may be allowed to say, with reference to the exercises, that the Schott method does not merely consist in causing the patient to execute certain movements which call his muscles into play. The essence of the system is that the efforts of the individual under treatment should be so regulated and timed as to produce no fatigue or sense of distress depending on incoördination of the functions of breathing and circulation; and that, whether baths or exercises be employed, they should be so regulated as never to overtax what, for want of a better expression, I may designate the patient's power of reaction. That condition will not have been observed if the rate of the pulse be increased or the volume of its wave diminished as the result of either proceeding. The effects produced by both baths and exercises are in most respects identical. The force of the myocardial contractions is

increased, and simultaneously, if not antecedently, the arterial capacity expands, tension is reduced, the volume and force of the pulse rise, and as a general rule its frequency diminishes, provided that the precautionary conditions just referred to are observed. Such important results are obtained in a space of time which is not to be measured by weeks or days, but by minutes. The chambers of the heart close on themselves at each contraction, inspiration becomes deeper, and loaded veins, released from the burden of back-pressure, discharge their contents through the right heart and the pulmonary circuit into the expanding arteries. The quickened capillary circulation imparts a glow of warmth and colour to the surface and the extremities, a healthy stimulus is imparted to the general metabolism, and relief is afforded to congested viscera throughout the body. Not many days or hours pass before the kidneys resume full activity of function, and free diuresis ensues with all its tissue-purging effects. The relief brought to the abdominal and portal veins and the increasing supply of arterial blood impart a healthy impetus to the entire process of digestion. Another effect of these revivifying changes which as far as I am aware has not hitherto been recorded is that the skin in many cases resumes its physiological office after, it may be, years of abeyance, and patients who have been long-unconscious of perspiration begin to experience a welcome sense of relief from a normal moisture of the surface. I think that it may safely be asserted that no drug or combination of drugs known to medical science is capable of effecting so immediate and far-reaching a health-restoring revolution in the human economy or one so perfectly adapted to correct the depraving and destructive form of self-poisoning which I have endeavoured briefly to portray, and which, if my suggestion be founded on fact, is so intimately associated with disabling functional and organic changes in the heart and bloodvessels. The best evidence of the truth of my contention is that recourse to the physical treatment is so often followed by the repair of cardio-vascular tissue, the restoration of heart competence even in spite of irreparable valvular deformity, the rehabilitation of the functions of digestion and elimination, the return of health as evidenced among other signs by renewed mental and physical power, and by the rapid resolution of adventitious products of many acute and chronic diseases to which special organs and the tissues generally are liable. On the other hand, I am far from wishing, as I have already sufficiently indicated, to cast doubt on the value of pharmaceutical treatment. It has its office—one with which no physician can afford to dispense—more especially in guarding the subjects of the diseases to which reference has been made from the danger of relapse. A judicious combination of drug treatment with the physical methods of the Schott system and with well-devised instruction as to how to eat, drink, and live is, happily, capable of endowing many, who have lost hope and are hastening on the road to dissolution, with years of unlooked-for healthy and happy life.

Upper Brook-street, W.

AN INQUIRY INTO THE PHYSIOLOGY OF THE ACTION OF THERMAL SALINE BATHS AND RESIST- ANCE EXERCISES IN THE TREATMENT OF CHRONIC HEART DISEASE (THE NAUHEIM AND SCHOTT SYSTEMS).¹

By R. F. C. LEITH, M.B., B.Sc., M.A., F.R.C.P. EDIN., &C.,
LECTURER ON PATHOLOGY, SCHOOL OF MEDICINE OF THE ROYAL
COLLEGES, EDINBURGH, AND SCHOOL OF MEDICINE FOR
WOMEN, EDINBURGH; PATHOLOGIST TO THE ROYAL
INFIRMARY, EDINBURGH; EXAMINER IN PATHO-
LOGY, UNIVERSITY OF GLASGOW, AND
ROYAL COLLEGE OF PHYSICIANS,
EDINBURGH, ETC.

THIS method of treating chronic heart disease, commonly called in this country "the Schott System," in recognition of its practical discoverers and exponents, was lately brought into prominence in England by Dr. Bezly Thorne and is now attracting much attention. It is claimed for it that it is applicable to an immense number of cases which hardly improve if at all under rest and ordinary cardiac tonics and

¹ A paper read before the Edinburgh Medico-Chirurgical Society, Jan. 15th, 1896.