the initial hard chancre, but it may develop at a later period. If the disease is seen early and anti-syphilitic remedies are vigorously pushed resolution will usually occur; but in the cases where the condition has persisted for a long period of time the anti-syphilitic remedies have little if any effect.

But little difficulty was encountered in the operation, although the hemorrhage was rather profuse. The flaps were brought together with some tension, but strange to say, healed

kindly. The woman today, four weeks after the operation, is perfectly well.

The lesson to be drawn from this case is this—that if the condition had been recognized at an early date and appropriate treatment given, this woman would have been saved seven years of suffering and in addition a severe surgical operation. The illustration gives but a poor idea of the enormous size of the growth.

TERATOMA OF OVARY.

F. G. DU BOSE, M.D., SELMA, ALA.

(Report of a case brought to operation on account of strangulation resulting from a twist of its pedicle.)

In the course of operations for the removal of tumors it is not rare to meet with dermoids containing heterogeneous tissue elements in varying stages of development from a mere resemblance up to a complete duplication of one or more mature body structures.

These result from embryonal cell misplacement. The case here presented is of interest not only on account of the tumor contents, but because the clinical history and physical examination was sufficiently clear to permit a diagnosis of a tumor strangulated from torsion at the first visit I made this patient.

CASE REPORT.

Mrs. M., age 48; multipara; healthy; robust; past history, negative; menstruation, regular and normal; suffered no pain in side till the morning of Nov. 28, 1907, when, on getting out of bed she felt pain in her left side. She had stooped and knelt down on the floor, wiping off and under a new couch, which had been placed in the room the previous evening, and first thought the pain was a stitch in her side. In an hour's time this pain

had so increased in severity that it was unbearable and a doctor was hurriedly sent for; he spent some time with the patient and gave several hypodermics of morphine before the pain was relieved. Though under the influence of opiates, she suffered pain at intervals. The following morning I was called and found, on palpation over left inguinal region, a movable tumor, painful on pressure, and some rigidity of the overlying muscles. Further examination per vaginam disclosed a pedunculated tumor of the left ovary, larger than a man's fist. The neoplasm was exquisitely sensitive and painful on bimanual palpation. Temperature and pulse normal. Operation advised and patient admitted to my sanatorium at 4 p. m. A saline enema was given with effectual result. Patient still suffered with pain in side, become severe at intervals. Hypodermic of codein, $\frac{1}{2}$ grain, was given. She spent a restless and painful night, and morphia and atropia were given at 7 a. m.; at 8:30 patient was carried to operating room. Gas-ether sequence anaesthesia. Left rectus incision, close to middle line. A reddish-black and ecchymotic tumor was delivered through incision. The pedicle, about two inches long, was twisted on itself, and the vessels supplying the tumor were thrombosed;

the pedicle was ligated flush with the broad ligament and the tumor removed. The right ovary and the appendix were also removed. Time of operation, forty minutes. The patient made a rapid recovery, and is in robust health at present.

attached to the cyst wall. The strands of hair are from 6 to 14 inches in length and of a red color. A mass of yellow fat filled the cyst cavity, and the hair was intermingled with this fatty substance. The shape of the tumor was ovoid and measured ten and eleven inches

The accompanying photograph shows the tumor turned inside out. The shaft of bone is $1\frac{3}{8}$ inches high by $2\frac{1}{4}$ inches in circumference. There are six well-formed teeth rather symmetrically arranged around the free end of the bone, the base of the bone being firmly

over its smaller and greater circumferences, respectively. The six teeth comprise 1 well-developed superior molar, 2 well-developed superior bicuspid, 1 poorly-developed bicuspid, 2 poorly-developed lateral incisors.