weeks after inoculation, produced generalised tuberculosis in the guinea-pig when its splenic emulsion was injected into that animal. The fourth pigeon was killed six weeks after inoculation; the result was similar to the last. The fifth pigeon was killed eight weeks after inoculation; its spleen on being inoculated into the guinea-pig produced general tuberculosis at the end of 44 days from the date of inoculation.

The relationship of the human to the avian bacillus as tested by means of the "opsonic" index.—A large number of experiments were made with the serum obtained from cases of human tuberculosis and avian tuberculosis on human and avian bacilli. In most cases normal human leucocytes were used in these experiments but in some cases birds' serum was treated with washed birds' leucocytes. The results of our observations show uniformly and in each case clearly that the opsonin in the blood serum of a tuberculous patient is removed or, more correctly, reduced when the serum is saturated with the avian bacillus as markedly as when it is saturated with the human bacillus. No specific difference in the behaviour of these two bacilli is brought out by these observations.

CONCLUSIONS.

1. The human tubercle bacillus is pathogenic to the pigeon in a very limited degree only. It sets up no disease of the intestine or of the abdominal viscera when introduced (in sputum) with food into the alimentary tract, even for prolonged periods (six months), and in either moderate or excessive quantities. It sets up a local, or a local and glandular, disease only in the pigeon when injected (in sputum) into the muscles or subcutaneous tissues. It sets up a local, or a local and glandular, disease only in the pigeon when injected into the muscles or subcutaneous tissues from the organs of guinea-pigs which have been infected by inoculation with human sputum. The impossibility of producing infection in the pigeon by means of tuberculous sputum, as shown in our work, does not allow us to accept the statements which have been made in the other direction, for these statements do not rest upon experiment but on inferences based only upon possibilities. Although the disease may be spread from bird to bird it has still to be determined whether there is any further source of infection. Our experiments only show that if there is such, it is not human. Neither can we accept the statement made by some that the subcutaneous inoculation of birds with human sputum will produce avian tuberculosis—i.e., the generalised and fatal disease as it is seen naturally in birds. In none of our pigeons has anything other than a local lesion or a local and a glandular lesion followed; moreover, the disease so induced instead of progressing has (as told by the examination of birds at different dates after infection) retrogressed, and in some disappeared.

2. The avian tubercle bacillus, as tested from various kinds of birds, is pathogenic to the guinea-pig in a very limited degree.

3. The human bacillus is not convertible into the avian by inoculation into the bird—i.e., the bacilli are not identical. Theoretically it would, of course, be quite possible that the human bacillus after its insertion into the pigeon might lose certain of its characters and become incapable of exciting more than a trivial disease in the guinea-pig or in the human subject from which it was originally derived. Our experiments do not, however, support this theoretical possibility. Using pigeons into the muscular and subcutaneous tissues of which a salt emulsion of human tuberculous sputum had been injected, we adopted the method of making a salt suspension of the spleen from these birds, killed at different dates after their inoculation (one week, two, four, six, and eight weeks), and injecting this into the subcutaneous or muscular tissues of guinea-pigs. In addition we similarly used the local lesion produced in the same pigeons. A fact of great importance to observe here is that the spleens of these pigeons in no case presented any naked-eye disease. In each bird there was a lesion at the site of inoculation; in the birds allowed to live the longest this was of the smallest extent. Although there was no disease of the spleen, the salt suspensions of this organ in every instance produced not merely a local lesion in the guinea-pig but a general tuberculosis. The human bacillus after a sojourn of as long as eight weeks in the spleen produced the glandular and visceral disease in the guinea-pig as though it had been transferred directly to this animal from human sputum. It did *not* produce merely a local or a local and glandular disease, such as it

should have done had it acquired the characters of the true avian micro-organism.

As evidence of the non-identity of the human and avian tubercle bacillus we attach more importance to the positive results afforded by feeding and inoculation experiments than to the negative evidence in the other direction provided by the "opsonic" test. The latter shows, it is true, that either bacillus will extract the "opsonin" from tuberculous serum. But we do not regard this as a proof of their identity; it is by no means a proof, i.e., of the identity of their physiological or chemical properties.

Certain investigations in course of publication (S. G. Shattock and L. S. Dudgeon) show, amongst other things, that the "opsonin" may be extracted from tuberculous serum by saturating it with a thick suspension of melanin—an observation proving that the removal of "opsonin" by bacilli is not wholly indicative of specificity but that it is largely mechanical.

NOTES OF

FOUR CASES IN WHICH FILIGREE-IMPLANTATION WAS SUCCESSFULLY APPLIED TO ABDOMINAL HERNIÆ PREVIOUSLY CONSIDERED INCURABLE.

BY LAWRIE H. MCGAVIN, F.R.C.S. ENG.,

SURGEON TO THE "DREADNOUGHT" HOSPITAL; LECTURER ON CLINICAL SURGERY AND TEACHER OF OPERATIVE SURGERY IN THE LONDON SCHOOL OF CLINICAL MEDICINE.

SOME time ago I published in the *Practitioner* the notes of seven cases of hernia in which I had applied the method of filigree-implantation as devised by Willard Bartlett. Since that time I have had occasion to employ it in a considerable number of cases, all of which have justified the claims put forward for its more extensive adoption. There are, however, among these cases four which merit publication for several reasons. First, three of them were of the same variety—viz., herniæ in the region of the appendix, two of them, indeed, resulting from the operation of appendicectomy. Secondly, all of them were genuinely incurable by any of the usual operations for ventral hernia, two of them being cases of recurrence and having been considered by surgeons other than myself as inoperable. Thirdly, all of them were complicated by pain (in one case by paroxysmal colic), retching, or inability to perform the ordinary duties of life owing to the constant prolapse of abdominal contents on the most trivial exertion. Lastly, two of them required filigrees of such size as to constitute (so far as I am aware) a record in the introduction of foreign bodies into the living tissues for curative purposes.

CASE 1.—In August, 1906, a woman was sent to me by Dr. A. H. Croucher of Eastbourne, with the following history. 18 months previously she had undergone an operation for uncomplicated appendicitis at the hands of a well-known operator in London. About nine months later, having removed the belt she was wearing, the cicatrix began to stretch and soon a well-marked hernia resulted. The belt was again resorted to but failed to control it, the hernia growing larger and being accompanied by pain in the iliac fossa and at times by nausea. She accordingly returned to her surgeon who endeavoured to repair the gap by approximating the muscles. Within three months the hernia again appeared, and this time the pain and nausea were such as seriously to incapacitate her. Returning to her surgeon, however, she was told that owing to the friability of her muscles her condition did not admit of any prospect of cure, and she was therefore recommended to wear her belt permanently. This she did for some weeks without relief, the hernia gradually increasing in size. On examining her I found her to be anæmic but otherwise healthy. The scar of her previous operations was very thin, pigmented, and widely stretched. It measured six inches in length, and on coughing or straining it bulged in an alarming manner. The breadth of the hand could easily be passed into the abdomen and coils of bowel could be felt to be adherent to the sac, manipulation of which produced feelings of nausea.

On Oct. 11th, 1906 Dr. F. W. Hewitt administering the anæsthetic, I excised the cutaneous scar and opened the abdomen at the extreme upper end of the incision which measured eight inches in length. Introducing two fingers

into the abdomen, I found a coil of bowel adherent to the right side of the gap and then completed the peritoneal incision in the whole length of the wound. The adhesions, which were of the ileum chiefly, were separated with some difficulty and the excess of sac was removed, the edges being sutured with a continuous catgut suture. The outer margin of the rectus muscle was now sought for and found to be much pushed inwards and rather fibrotic; it was freed as far as possible and the peritoneum was stripped back from the deep surface of the transversalis fascia for a distance of about two inches. The margin of the oblique muscles was next found and freed from scar tissue, and the peritoneum on this side was treated in a manner similar to that already described. All oozing was now controlled by the use of hot saline lotion, and the edges of the wound being widely retracted, the filigree, measuring $6\frac{1}{2}$ inches in length and 3 inches in width, was carefully introduced, being laid upon the peritoneum. The rectus muscle and the edge of the oblique muscles were drawn over the margins of the filigree and brought as closely into apposition as possible. This was only possible by the exercise of considerable tension, and even then the muscles were only approximated at the actual points of suture. Finally, the aponeuroses were closed over the muscles and the skin incision was closed by the use of Michel's clips. Convalescence was uninterrupted and since then the patient has never worn anything in the shape of a belt or binder. She tells me that she is in perfect health and that, in her own words, she is "beltless and bandageless, and it is a joy to live in comfort again." When I last saw her the abdominal cicatrix was firm and hard; no amount of straining affected it; there was no sensation of discomfort from the presence of the filigree.

CASE 2.—The patient was a woman, aged 21 years. She had been admitted to Guy's Hospital seven years previously, having been crushed between a wall and the tailboard of a cart. Her symptoms suggested the occurrence of internal hæmorrhage, and a few hours after admission her abdomen was opened by a six-inch incision above the umbilicus. No hæmorrhage was found, but there was an extensive condition of tuberculous peritonitis with ascites. For this reason a drainage tube was placed in the wound. Five days later, during a fit of coughing she ruptured the stitches above and below the tube and some coils of bowel appeared; these were washed and returned and the wound was closed by complete suture. Primary union resulted and the patient was discharged well in the course of a month. Soon after that time she began to notice that the wound was bulging and this it continued to do, becoming gradually larger. During the six months previous to her visit to me the hernia had progressed rapidly owing to her commencement of domestic service. I first saw her in the autumn of last year; she then had a well-developed epigastric hernia, the gap measuring 4 inches wide by 5 inches in length. Many adhesions could be felt in the sac and the patient suffered much from retching and actual sickness during her work and complained of gripping pain in the region of the stomach on attempting to lift any weight. She was of spare build but looked healthy and of fair muscular development.

On Nov. 23rd, 1906, I opened the abdomen by a seven-inch incision below the ensiform cartilage, removing all the cutaneous scar. No sign of tubercle was to be seen but the resulting adhesions were numerous and very dense, involving the stomach and transverse colon. Some omentum was found adherent to the sac and was removed. These adhesions were divided and the abdominal contents were returned. I did not attempt to place the filigree in this case on the peritoneum owing to its densely adherent condition, which would have made any attempt to strip it from the fascia a source of oozing. In such cases I prefer to place it between the rectus muscle and its posterior sheath. I therefore split the adjacent margins of the rectus sheath to the full length of the incision, and dividing the linea transversa in front of and behind the muscles, drew the latter from their sheaths. The peritoneum and posterior rectus sheath being inseparable were closed in one layer, a portion of the sac being utilised on either side to make up the deficiency and to render closure possible. Upon this the filigree, measuring $6\frac{1}{2}$ inches by 3 inches, was placed, and the muscles were then with some difficulty closed over it. The anterior layer of the sheath was united to its fellow separately and the skin incision was closed as in the last case. Recovery was uneventful and the patient remains at the present time in perfect health and comfort, retching, vomiting, and pain having ceased from the time of the operation. The epigastric region is firm and flat.

CASE 3.—The third case is that of a woman, aged 36 years, who was transferred to my care in the Hospital for Women in January, 1906. Three weeks previously she had undergone operation for a large and densely adherent cyst of the right broad ligament. At the time of her transference she had developed a firm, tender swelling in the right inguinal and iliac regions with some œdema over it, and a temperature of 103° F. This swelling was clearly due to pus and was thought to be the result of pelvic infection. I therefore opened it close to the anterior superior spine of the ilium and evacuated a large quantity of thin pus. On the next day nothing but pure urine came from the wound, and it was then clear that the abscess was in the ureter, which had been unfortunately ligated. A few days later, therefore, I opened the abdomen, after sealing off the incision, internal to the linea semilunaris, and endeavoured to find the end of the ureter by tracing it from above the pelvic brim. This was found to be impossible on account of the density of the adhesions, and in separating some of these faecal matter appeared in the wound in the vicinity of the cæcum. I accordingly abandoned the attempt, and having passed a large tube down to the site of extravasation and plugged it about with gauze, turned the patient over and removed the kidney through the loin. In a few days a gangrenous patch appeared about the drainage-tube in the abdomen, and gradually an area of about the size of a five-shilling piece sloughed out, including all structures down to the peritoneum. Although at this time the patient was extremely ill and was thought to be dying she showed wonderful vitality and gradually recovered, all her wounds being healed at the end of ten weeks.

In January of this year she returned to me at the Hospital for Women with one of the largest ventral herniæ I have ever seen, at the site of the gangrenous patch already mentioned. She had grown stout since her discharge, and the appearance of the mass, covered as it was by feeble-looking cicatricial skin, was sufficient to put any ordinary form of operative treatment quite out of court. She was in constant pain with attacks of severe dragging colic attended by violent retching and sometimes by actual vomiting. She begged that something might be done for her and readily consented to the proposal that the attempt to implant a filigree should be made. As far as could be judged through her stout abdominal wall the fibro-muscular edges of the gap inclosed an area of about 7 inches by 4 inches. A filigree was therefore constructed for me by Messrs. Down Bros. measuring 8 inches by 4 inches, and on Jan. 19th, Dr. Maynard Horne acting as my anæsthetist, I proceeded as in the first case to excise the cicatrix by an elliptical incision 11 inches in length. On opening the abdomen at the upper end of the incision and introducing the finger for exploration the cæcum was found to be adherent to the sac and to a large mass of omentum weighing about $1\frac{1}{2}$ pounds which was subsequently excised. I found it quite impossible to free the peritoneum from the aponeurosis of the external oblique muscle or from the posterior sheath of the rectus, and was therefore met by the difficulty of forming a bed for the filigree and finding material with which to cover it over. I accordingly employed a method which I had not attempted before. This consisted in incising the margin of attachment of the external oblique muscles to its aponeurosis for a distance of eight inches so as to expose the muscular fibres. This permitted of the detached aponeurosis being drawn inwards sufficiently to meet the posterior sheath of the rectus (Fig. 1), to which it was then sutured. The incision in the aponeurosis was thus converted into an ellipse which was floored by the peritoneum which was stripped from the posterior surface of the oblique muscles. A bed being thus provided for the filigree the latter was introduced so as to lie with its inner loops on the posterior sheath of the rectus muscle and the divorced aponeurosis of the oblique muscle, and its midrib and outer loops on the peritoneum forming the floor of the ellipse above mentioned. The rectus muscle was then brought out of its sheath from the outer side and in its whole width was transplanted outwards so as to meet the cut margin of the oblique muscles from which the aponeurosis had been divorced. This provided covering for the inner two-thirds of the filigree, while the outer third was covered by the inner margin of the oblique muscles. The rectus muscle was with difficulty sutured to this margin with stout catgut sutures passed at some distance from the muscular edges. The transplantation of the rectus muscle I then found had left an elliptical weak spot between itself and the left rectus, since the former was shrunken and partly atrophic;

to close this I opened the left rectus sheath and drew over the muscle of the left side across the middle line and sutured it there. The operation was finally completed by closing the skin incision by means of 35 Michel's clips, no aponeurotic tissue being available for covering the muscles. The operation lasted over two hours and was, I may say, of unusual

FIG. 1.

D, Divorced strip of external oblique aponeurosis sutured to (s) posterior sheath of rectus muscle. E, External oblique muscle. R, Rectus muscle. P, Peritoneum covered actually in whole area, shown by F, Filigree, a portion only of which is here shown diagrammatically *in situ*.

difficulty. The result has been extremely gratifying; the wound healed by first intention, and at the present time the patient is in excellent health, her abdominal wall being firm and resistant.

CASE 4.—The last case is even more striking than the above. In 1904 a young woman, aged 22 years, was admitted to Guy's Hospital, under the care of Mr. W. Arbuthnot Lane, for gangrenous appendicitis and general suppurative peritonitis. The abdomen was opened over the appendix and in the middle line above the pubes. These wounds were left open for drainage, and as the patient gradually recovered they were sutured four days later. Shortly after this there was a rise of temperature, and faecal matter appeared on the dressings over the appendical wound, and from this time there was a complete faecal fistula of the small intestine. From the character of the excreta it appeared that the fistula was high up in the bowel, and the girl began to waste rapidly. At this time she came under the care of Sir Alfred Fripp. Four attempts were made by suture to close the fistula without success, and finally, when her condition was of the gravest, a complete enterectomy, involving removal of six inches of the intestine, was performed and an end-to-end anastomosis successfully carried out. The patient now gradually recovered and was eventually discharged. Six months later she was again admitted to hospital, under the care of Mr. Lane, for a very large ventral hernia at the site of the appendical cicatrix. The sac was excised and the abdominal muscles approximated, but the gap appears to have been too great, for the hernia quickly reappeared and this time it assumed much larger proportions than before. She was again seen by Sir Alfred Fripp, who kindly sent her to me for submission to operation by filigree implantation, as he considered that any other method of treatment was out of the question. On examining her I found a hernia of really gigantic proportions, occupying a considerable part of the right lumbar, right inguinal, umbilical, and hypogastric areas of the abdomen (Fig. 2), and passing from two inches below the tip of the tenth rib, obliquely downwards and inwards to an inch above the pubic spine. The length of the actual gap was $7\frac{1}{2}$ inches and its width $4\frac{1}{2}$ inches. Coils of adherent bowel could be seen in the sac, and peristalsis was clearly visible through the thinned out and pigmented cicatrix. The entire hand placed flat upon the scar could be passed straight into the abdomen. The patient told me that from time to time she was seized with attacks of acute agonising pain in the abdomen, accompanied by sickness, and that her life was in that condition insupportable. In considering whether such a case came within the limits of justifiable surgery I was influenced in my decision by several points. The girl was

apparently in the best of health otherwise; she was dependent entirely on her work for her living; her life was, as above stated, wretched; the control of the hernia was beyond the scope of any form of belt or truss; the future state of the hernia was likely to be one of rapid progress; the cases of filigree-implantation I had already attempted had been uniformly successful; and finally, there was the extraordinary vitality of the girl as shown by her previous surgical history. In these circumstances I agreed to attempt the cure of her condition.

FIG. 2.

The shaded portion in this diagram represents the area of the hernial protrusion in Case 4.

The patient was admitted to the Dreadnought Hospital, Greenwich, in January of this year and on the 22nd, Dr. Cecil H. M. Hughes being the anaesthetist, I removed the entire cutaneous cicatrix by an ellipse measuring 13 inches by 3 inches. In doing this, as I had anticipated, the peritoneal cavity was opened about the centre of the wound. The abdomen was opened again, as in the other cases, at the extreme upper limit of the wound and with the finger as a

FIG. 3.

R, Rectus muscle partly covered by A, representing oblique aponeurosis and anterior sheath of rectus muscle. E, External oblique muscle. C, Caecum sutured on inner side to small intestine (I), and adherent on outer side to P, peritoneum. P', Conjoined peritoneum and posterior sheath of rectus muscle sutured to small intestine (I). F, Portion only of filigree shown *in situ*.

guide the peritoneum was laid open in the whole length, keeping to the umbilical side of the mass of adhesions which was glued to the right side of the incision. At the end of an hour I had succeeded in separating by dissection the greater portion of the adhesions between the ileum, omentum, colon, and abdominal wall, although much had to be left undone as unjustifiably dangerous. An attempt to strip

the peritoneum from the posterior surface of the abdominal wall resulted in so much tearing that a large portion of it was rendered quite useless for the purpose of carrying a filigree, a gap measuring 3 inches by $1\frac{1}{2}$ inches remaining patent. As the omentum was not long enough to reach beyond the upper portion of the wound, into which it might otherwise have been sutured, I chose the strongest loop of small intestine in the vicinity of the gap and drew it into the latter, suturing it on the outer side to the inner margin of the cæcum which was adherent to the peritoneum (Fig. 3) and on the inner side to the outer margin of the conjoined peritoneum and posterior sheath of the rectus muscle. Upon this patchwork of peritoneum, intestine, and fascia the filigree was placed; it measured 9 inches in length, 4 inches across the centre, and 3 inches at either end (Fig. 4). The operation was then completed in a manner similar to that described in the last case, the filigree being covered by the atrophic rectus to the inner side and by the oblique muscles on the outer side. The frayed remains of the aponeurosis of the external oblique were united to the anterior sheath of the rectus muscle and the skin incision was closed by 40 Michel's clips and four stout salmon-gut tension sutures. The patient stood the operation, which

FIG. 4.

The diagram shows the pattern of filigree used in these cases and is drawn to one-third the size of that employed in Case 4. A A, Midrib of filigree.

lasted nearly three hours, very well, her convalescence was uninterrupted, and the clips were removed at the end of the seventh day. She was discharged from hospital at the end of six weeks and from the time of her operation has been free from pain or discomfort of any kind. She declares herself fit for work, to which she has not yet, however, been allowed to return.

The above four cases are, I think, good examples of the class of case which, having hitherto been regarded as beyond the power of surgery, has been brought within our grasp by the application of Bartlett's method. Since this surgeon first introduced his filigree to the notice of the profession in America I have had occasion to make use of it in over a dozen cases. Seven of these cases I published some time ago in the *Practitioner*; the four here related, among others, have been done since that time. I have had little hesitation in publishing them so soon after operation, the results of my earliest cases having convinced me of the soundness of the method. In adopting Bartlett's method I have been struck with one or two points of importance which it may not be out of place to mention here. The situation of the filigree will usually be found to be determined not by any stereotyped rule but by the conditions presented by each individual case, and although it is advisable to place the filigree as close as possible to the abdominal cavity it will be found to be at times a matter of impossibility to place it actually on the peritoneum. I have elsewhere drawn attention to the firm

adhesions which often exist between this structure and the transversalis fascia in cases of old-standing umbilical hernia and in some cases of ventral hernia following operation or accidental trauma. The attempt to separate these structures in such cases in order to lay the filigree upon the peritoneum results in the tearing of holes in the latter and in most troublesome and persistent oozing. Time is thus lost, shock is increased, and suppuration is favoured. There exist other situations in which a filigree may be implanted. In the case of median hernia above the umbilicus, or in umbilical hernia and for two inches lower down than the umbilicus, the space between the rectus muscles and their posterior sheaths is admirably suited to the purpose. The separation of these structures is a very simple matter; there is little oozing to endanger the union of the wound and the bed so formed for the filigree is firm and resistant.

In the case of hernia following appendicectomy, or other operation through the lateral wall of the abdomen, the difficulty of forming a bed for the filigree may be very considerable. It will arise where there is a deficiency of peritoneum with which to close the abdomen, either from tearing of holes in the attempt to strip or from cutting out of sutures where the tension is great. This difficulty may be met in one of three different ways. First, the omentum, if it is long enough and not widely adherent, may be sutured into the gap and the filigree laid upon it. Secondly, where this is impossible the intestines themselves, if in good condition, and especially if more or less adherent to each other already, may be utilised, as in my fourth case, to close the gap. This method is, of course, only admissible as a last resource when no other method is available, and must be carried out with the greatest care lest the whole operation be rendered impossible by puncturing the bowel. Thirdly, where there is no lack of material but the peritoneum is so densely adherent to the fascia of the abdomen as to be inseparable, an incision may be made along the margin of attachment of the external oblique muscle to its aponeurosis down to the peritoneum, as shown in Fig. 1. The portion internal to this incision is isolated from the remainder of the aponeurosis by carrying the upper and lower ends of the incision inwards towards the middle line. It will then be found that the peritoneum can be stripped away from the oblique muscles, so that this peritoneo-fascial strip can then be made to meet the outer edge of the posterior sheath of the rectus muscle which is probably also inseparable from its peritoneum. The filigree can now be laid upon this bed and with a little care the rectus muscle can be drawn from its sheath and transplanted outwards so as to meet the edge of the external oblique muscle which has been laid bare by the divorce from its aponeurosis and which will serve to cover the outer loops of the filigree.

The first of these methods I have not had occasion to use; the latter two I devised to meet the requirements of the last two cases of the series. Recently I have found occasion to modify slightly the shape of the filigree originally described by Bartlett. As in all hernial gaps the widest part is usually the most central, it is clear that defence is more essential here than at either end; there is, therefore, little to be gained by making the filigree the same width throughout its whole length; and as there is often considerable difficulty in getting the corners of a large oblong filigree fitted underneath the muscles at the extreme ends of the incision, I now have all filigrees made for me for abdominal herniæ constructed as in the last case described—i.e., with ends slightly narrower than the measurement across the centre. For the sake of uniformity all are constructed, whether for inguinal, umbilical, or ventral hernia, of the same gauge of wire—viz., No. 28 standard wire gauge, and have the same number of loops to the inch—viz., eight. In conclusion, I desire, as the introducer of Bartlett's method into this country, to state my belief that in it we have the simplest, quickest, and surest means yet devised of commanding cases of hernia of all varieties (with the possible exception of femoral hernia) hitherto considered incurable.

For the accompanying diagrams I am indebted to the pen of Dr. Ernest T. Jensen.

Mansfield-street, W.

ROYAL UNITED HOSPITAL, BATH.—The Mayor of Bath (Mr. S. Bush) has for some time been making strenuous efforts to free the Royal United Hospital in that city from a debt of £6124, and he received a munificent offer of £1000 from Mr. Overton Wills provided that the further sum of £5000 was subscribed. On Nov. 8th the amount in question was raised and the hospital is now free from debt.