gated to the past. Of special interest are his suggestions for the management of disturbed and unclean patients. He estimates the proportion of patients in a mixed asylum who can with advantage be subjected to bed treatment as high as 60 per cent.

5. *Dementia Precox or Brain Tumor?*—Description of the interesting case of a young woman in whom for ten years typical symptoms of dementia precox, with eventually an extreme grade of dementia, were present. She then began to have Jacksonian convulsions specially affecting the right side of the face and right arm, which continued at intervals until her death about eleven months later. The eye-grounds were normal. Post-mortem examination showed a diffuse glioma with hemorrhage and softening affecting nearly the whole of the right hemisphere. The occurrence of the Jacksonian attacks on the right side, and not on the left, as would be expected from the location of the tumor, the author thinks due either to pressure or to an existing meningitis, as otherwise no macroscopic changes were evident on the left side. A microscopical examination does not seem to have been made. He states, however, that neither pyramidal tract appeared quite normal, and that he was able to assure himself that decussation was present. Discussing the question as to whether the tumor was actual cause of the initial mental symptoms or only a complication of preëxisting dementia precox, he inclines to the opinion that since the neoplasm was of probably very slow growth, the former view of the case is the correct one.

6. *Mental Condition of Deaf Mutes.*—Continued article.

7. *Persons Ordered Committed to German Asylums.*—Statistical table.

8. *Simulation of Mental Diseases.*—Continued article.

9. *Simulation of Attacks of Pain in a Morphine Habitue.*—An account of the case of a man of considerable education who had led the life of a swindler, had been frequently in prison and had contracted the morphine habit, who, interned at Waldheim both for cure of the drug addiction and for an opinion as to his mental condition, came under the care of the author. After careful study of the case the conclusion was reached that he was not insane, but an ordinary liar and swindler, and that his pains were simulated to secure morphine, to arouse sympathy and to escape or lessen punishment. The author does not, however, deny that the patient may belong to the class of psychically inferior, and thinks that institutions for the care of such people would afford great relief to doctors and to the public generally.

10. *Endogenous Symptom-Complexes in Exogenous Disease Forms.*—Description of a case of general paresis which began suddenly and for a year presented symptoms of manic depressive insanity without any of the somatic signs of general paresis, and with no impairment of memory or intelligence. The patient was sent home apparently convalescent. Shortly afterward he had a convulsive attack followed by typical symptoms, both somatic and psychical, of dementia paralytica. The author takes occasion to discuss the psychological mechanism of the production of various morbid mental manifestations in the so-called endogenous forms or degenerative insanities. He finds theoretically nothing opposed to the practically well-known fact that symptoms supposedly characteristic of these endogenous forms often occur both in general paresis and in other insanities due to exogenous causes. ALLEN (Trenton).

Miscellany

MENTAL DISTURBANCES IN REVIVAL AFTER HANGING. B. Bayerl (Wiener klinische Rundschau. Volume XIX, No. 4).

A young man, upon the restoration of the vital functions after an attempt to commit suicide by hanging, displayed a maniacal condition, with dilated, non-reacting pupils. After a number of days the mania subsided,

and he had never showed any mental disturbance before or afterward. His arrest on a charge of murder was the occasion of his deed.

JELLIFFE.

PATHOLOGICAL SLEEP IN CASES OF CEREBRAL TUMOR. Fr. Franceschi (Riv. di pat. nerv. e ment., October, 1904).

Franceschi describes a case of cerebral tumor attended from the first by a tendency to sleepiness, and raises numerous questions with regard to the nature both of physiological and of pathological sleep. He notes that while all authors mention pathological sleep as a symptom of cerebral tumor, it is usual to note its existence without any investigation of its cause or any explanation of the different stages of the disease at which the symptom shows itself. Some authors do not distinguish clearly between somnolence, sleep and coma, and it is, therefore, impossible to draw exact conclusions from cases reported by them. The principal questions raised are those of the mechanism of sleep and the effect of the localization and nature of cerebral neoplasms in producing sleep. Righetti has shown that pathological sleep is a more common symptom of tumors in the third ventricle and the hypophysis than of those situated elsewhere. In this case the patient suffered from somnolence, and later from mental torpor, for five months before an affection of vision and the increasing severity of the other symptoms led to her admission to hospital. At that time she presented the appearance of a continuous and tranquil sleep. She was capable of being aroused, but quickly slept again. Later she suffered from headache and vomiting, and she died of bronchopneumonia a month after admission. She was conscious when awake up to the end of life and frequently executed voluntary movements. Her sleep was much disturbed by the bronchopneumonia. It was, therefore, clearly sleep and not coma. At the autopsy a sarcoma was found originating probably in the dura mater, and afterwards involving the base of the third ventricle and the neighboring parts. The tumor was of such a size and in such a position as to be closely surrounded by the circle of Willis. The internal carotids were half their normal size, and the posterior communicating arteries were a quarter or a fifth of their normal size. The basilar arteries were compressed by the tumor. The most commonly-received explanation of sleep caused by a cerebral tumor is that it is due to increased intracranial pressure, but in this case there was no appearance at the autopsy of anything beyond slight increase of tension, and no likelihood that there was any appreciable increase of tension when somnolence was first observed. One of the best-established facts with regard to sleep is that it is attended, either as cause or effect, by anemia of the cerebral cortex. That theory of sleep which regards it as caused by anemia of the cortex fits best with the conditions observed in this case, and the frequency with which pathological sleep is associated with tumors of the optic thalami, the hypophysis and the floor of the third ventricle may be reasonably attributed to pressure on the circle of Willis, interfering with the circulation through the brain. Other theories—chemical, histological and evolutionary—of sleep causation the author accepts as complementary to and not incompatible with this. A much larger number of careful and accurately recorded observations is needed before each hypothesis can be credited with its true worth. Franceschi briefly reviews the opinions of those who maintain the existence of a special center for sleep located in the central grey substance of the brain, and points out that observations made in this case might well be cited in support of such a theory.

JELLIFFE.

THE CURABILITY OF EARLY PARESIS. C. L. Dana (Journal A. M. A., May 6).

The author suggests that paresis, like tabes—with which it is closely related as a parasyphilitic disorder—may be arrested in its earlier stages.