appeared to be some improvement. Oliver and Schäfer⁷ find that pituitary and thyroid extracts have opposite physiological effects. Pituitary extract resembles in its action that of the suprarenal capsules, though there are points of difference. Thyroid extract has been found to exercise a curative effect in a case of acromegaly recorded by A. H. Benson.⁸ Murray,⁹ however, thinks it doubtful whether thyroid extract has any specific effect on acromegaly; this opinion is endorsed by Osler.¹⁰ It is not improbable that any good it does is by influence on the general health. If acromegaly were due to a hypersecretion of the normal internal secretion of the pituitary body, the administration of pituitary extract should intensify and giving thyroid extract should remove the symptoms. If this is not the case—and it does not appear to be so—hypersecretion on the part of the pituitary gland cannot be regarded as a satisfactory explanation of acromegaly.

III. Perverted function due to a heterogeneous transformation of the structure of the pituitary body (Marinesco). But in what direction the secretion is perverted is a matter of speculation. Three alternatives may be suggested. 1. The secretion might be deficient in an active principle the function of which is to restrain exuberant growth in the extremities. This, perhaps, is rendered improbable by the general failure of pituitary extract when given medicinally, and there is of course no evidence that irregular growth is spontaneously likely to occur if not kept in check. 2. The abnormal secretion might contain some toxic principle which stimulated the tissues to overgrowth. That the disease is a toxæmia receives some support from its points of resemblance to hypertrophic pulmonary osteo-arthritis. Marie suggested that the changes in the limbs characterising the latter disease were due to the local action of poisons absorbed from the pulmonary lesions. 3. It might be thought that owing to alteration in the secretion of the pituitary the equilibrium and interaction between it and other internal secretions are disturbed. Possibly some toxic body such as was suggested in the preceding alternative might be thus produced. The thyroid and pituitary glands have been thought to be compensatory; recently they have been shown by Schäfer and Oliver to be physiologically antagonistic. Superficially there are some resemblances between acromegaly and myxœdema, and it might be thought that acromegaly is the result of a disturbance of a chemico-physiological equilibrium maintained in health by the normal activity of these two glands. Such a consideration suggests that acromegaly might be benefited by giving pituitary and thyroid extracts at the same time. The dose of the two extracts should be varied so as to get the desirable mean, but at first, perhaps, a larger quantity of the pituitary extract should be given. The failure of pituitary extract to ameliorate the symptoms of acromegaly is then somewhat in favour of the view that the cause of the disease is perverted and not suppressed or exaggerated functional activity of the gland.

BOLTON INFIRMARY.

A CASE OF TRAUMATIC VENTRAL HERNIA AND ONE OF HÆMOPHILIA.

(Under the care of Mr. GARSTANG.)

TRAUMATIC ventral hernia of the kind described in the first case is of very unusual occurrence—in fact, it has received little notice in our general text-books. Such a hernia, as the result of direct subcutaneous injury, may develop immediately in consequence of rupture of the muscles in the abdominal wall, the protrusion taking place through the opening made, or atrophy with resulting weakness may ensue at the point injured and a hernia develop secondarily. Examples of the former are given by Gay, Meckel, and Bryant. The patient under the care of Gay had a blow from a buffer in the abdomen, and on dissection some of the viscera were under the skin and had come through a rent which divided the left rectus and left epigastric artery, the linea alba, and part of the right

rectus. Included in the term traumatic, and much more common than those already referred to, are the herniæ which protrude through the scar resulting from the wound made by a surgeon for the removal of an abdominal tumour or for the evacuation of an abscess, also wounds of the parietes received in other ways by stab or gunshot. We are still in a condition of uncertainty as to the exact deviation from the normal which produces hæmophilia, and the facts that are known about it, such as its hereditary character, its more frequent appearance in the male, and also that it is congenital, do not help us in respect to the treatment of the hæmorrhages which are so apt to prove fatal. It is probable that antiseptic treatment with pressure will do more for the arrest of hæmorrhage in these cases than was formerly possible. When the wound suppurated and hæmorrhage frequently recurred from it the attempt to apply pressure or the application of styptics only seemed to make the wound a larger one and the fever more intense. Of the numerous drugs which have been recommended for the treatment of the hæmorrhages the only one which has seemed to us to exercise any control on the loss has been belladonna, applied externally in the event of large subcutaneous hæmorrhages, or given internally in cases of bleedings from the mucous surfaces. For the notes of the cases we are indebted to Dr. G. Wilkinson, late senior house surgeon.

CASE 1.—A boy aged fifteen years was admitted to the Bolton Infirmary on Feb. 7th, 1895, with a swelling in the left hypochondriac region the result of an injury. He gave the history that he was carrying a bundle of dye-sticks when he stumbled and fell with his abdomen on the blunt end of one of them. On admission there was found to be a swelling about the size and shape of half an orange, midway between the umbilicus and the cartilage of the tenth rib on the left side. When the patient coughed the swelling enlarged. It could be put back entirely into the abdominal cavity, and when this was done a well-defined rounded opening about one and a half inches in diameter could be felt in the abdominal wall beneath the skin. The substance protruded beneath the skin when the patient coughed and had the knotted feel of omentum. The skin over a part of the swelling was superficially abraded. The patient was put to bed, carefully dieted, and a pad and bandage were applied to the seat of injury. On March 6th the subcutaneous rent in the abdominal wall had nearly closed, so that only the tip of a finger could be introduced, and the abrasion had healed. The patient was supplied with an abdominal belt with pad and was allowed to get up. On Dec. 9th the patient came up for inspection. He had resumed work and discontinued the use of the belt. There was a prominent scar marking the seat of abrasion, which showed signs of cheloid growth. The rent in the muscular layers had quite closed and firm scar tissue was felt.

CASE 2.—A boy aged eleven years was admitted on June 18th, 1894, with hæmorrhage from the cheek and gums. He gave the history that he was struck on the left cheek with a cricket bat four days before admission and that there had been continual bleeding from his gums and cheek since. His previous history was as follows. When two years of age his tongue ulcerated and bled for three weeks; at three years of age he received a wound on the back of his head, which bled during five or six weeks; when seven or eight years of age he had a wound of the temple, which bled three or four weeks in spite of plugging, caustic, &c. There was no history of any male relative having been similarly affected. A sister aged twenty-five years had had four miscarriages and a boy who died at the age of eight months. The patient had a sallow complexion with transparent condition of skin. The conjunctiva was not unusually pale and there was no heart murmur. Tincture of perchloride of iron was applied to the wound of the cheek without much beneficial effect. Finally a silk suture was introduced and proved effectual. Arsenic and tincture of perchloride of iron were given internally and the mouth was frequently washed out with iced Condy's lotion. On July 9th the wound had quite healed. On Nov. 28th, 1895, the patient was readmitted on account of hæmorrhage from a wound of the upper lip inflicted by falling against a frame eight days previously. The hæmorrhage having ceased after admission it was not thought advisable to disturb him. Free hæmorrhage, however, came on in the night, which was controlled by the introduction of a silk suture. On Dec. 23rd the wound had not healed, but was slowly granulating.

⁷ Journal of Physiology, 1895, p. 277.

⁸ Brit. Med. Jour., vol. ii., 1895, p. 949.

⁹ Ibid., vol. i., 1895, p. 294.

¹⁰ Principles and Practice of Medicine, 1895, p. 1047.