might lead to persistent congestion of the uterus, such as traumatism during local treatment, sexual excess, etc. As soon as a small fibromyoma is detected especial care should be taken to place the uterus under the most favorable condition for limiting its growth. Ligation of both uterine arteries is a rational procedure at this early stage. In three cases in which the writer performed this operation there was a notable diminution in the size of the growth in two instances, while in the third it disappeared entirely (?), there being a notable relief of the menorrhagia.

The relief afforded by castration is to be referred rather to the diminution of the blood supply to the tumor consequent upon ligation of the ovarian arteries, in addition to the elimination of the periodical monthly congestion.

SARCOMA UTERI DECIDUO-CELLULARE.

SÄNGER (*Archiv für Gynäkologie*, Band xlv., Heft 1) presents another elaborate paper on the subject of this form of neoplasm, which he was the first to describe. He suggests, in common with Pestalozza, a possible parasitic origin, in view of the fact that several of the reported cases presented phenomena suggesting an infectious disease of the decidua. It seems to have no relation to septic infection. The principal clinical symptoms are persistent hemorrhages, occurring sometimes after an abortion or normal delivery, which may soon cease after curettage. Following the hemorrhages there is a constant foul, sanguinolent discharge, with later septic symptoms. The uterus gradually enlarges and assumes a nodular form, while progressive anemia, and finally cachexia are noted. Metastases develop, especially in the vagina. Secondary growths in the lungs, and hemorrhagic pleural effusion being indicated by the appearances of dyspnea, cough, and bloody expectoration. The disease runs a rapid course, terminating fatally within six or seven months. The condition should be suspected when profuse hemorrhages and foul discharges follow normal labor or abortion, especially if these discharges continue after the removal of retained products of conception. Puerperal septic endometritis and tuberculosis are most likely to be diagnosed in these cases. The growth is distinguished from carcinoma and ordinary round-celled sarcoma by the microscope, the history of a recent pregnancy being an important clue to the true condition. In order to confirm the diagnosis it is necessary to dilate the cervix with tents and to palpate the uterine cavity, instead of relying on the curette alone. The prognosis is bad on account of the tendency of the neoplasm to early metastasis, hence the necessity of prompt extirpation of the uterus. In every case of retention of the products of conception, and especially after the removal of placental moles, we should bear in mind the possibility of beginning sarcomatous degeneration.

URETERO-URETEROSTOMY.

KELLY (*Bulletin of Johns Hopkins Hospital*, Oct., 1893) reports an interesting case of hysteromyomectomy, in which the dilated ureter was ligated and divided under the impression that it was an enlarged vein. After removing the tumor the question of disposing of the incised ureter was considered, three modes of treating it being presented: to establish a fistula by

suturing the proximal end in the flank or in the abdominal wound, to remove the kidney, or to anastomose the ureter. The latter course was adopted, a slit being made in the lower portion of the ureter below the point of ligation, into which the proximal end was invaginated and secured by fine silk sutures, drainage with gauze being used in case of leakage. The patient made an uneventful recovery, being catheterized every four hours during the first day, after which she passed her urine voluntarily. There was no marked diminution of the daily amount of urine, and though albumin was present in considerable amount before the operation, none appeared subsequently.

REMOVAL OF THE UTERUS FOR SUSPECTED MALIGNANT DISEASE.

CORDIER (*Kansas City Medical Index*) properly holds that it is better to remove the uterus even when the microscopical evidence of malignant disease is not positive than to wait until the diagnosis has become certain, when the case may be inoperable. His conclusions are as follows:

Cancer of the cervix uteri is nearly always a local disease, which is sure to terminate fatally if allowed to run its usual course. Early extirpation is attended with a low rate of mortality and is curative in a considerable number of cases. Microscopical examination of suspected tissue does not always present the typical appearances of cancer, even when it is present; hence, one should not base his decision against operative interference on this criterion alone when there is other strong evidence of existing malignant disease.

THE SURGICAL SIGNIFICANCE OF DUST.

HAEGLER (*Beitr. zur klin. Chirurgie*, Band ix., p. 496) has found by experiment that floating germs in the air of an operating-room may be almost entirely removed by steam, and that their reaccumulation is best prevented by keeping the floor, walls, and furniture damp. Before removing the dressing from the field of operation it should be moistened with sterilized water. No dressings intended to be placed over the wound should be exposed, but they should be kept in closed dishes the interior of which is kept moistened with an antiseptic solution. The dust on the floor and walls of the operating-room is to be regarded as infectious material, the evil effect of which may be neutralized by saturating the room with steam a half or a quarter of an hour before the operation. The operator and his assistants should pay particular attention to the hair, which should be moistened or oiled. Sterilized gowns are safer if they are worn when they are still damp from the sterilizer.

PETROLEUM IN THE TREATMENT OF CANCER.

DESPRÈS (*Gaz. des Hôpitaux*, 1893, No. 68) employs refined petroleum in the treatment of inoperable carcinoma of the cervix uteri, making deep injections into the growth, which are painful but cause a speedy separation of sloughs, desiccation of ulcerating surfaces, and disappearance of odor. Injected into abscesses, petroleum causes speedy healing. In cases of acute vaginitis he uses thrice daily a douche of from three to five ounces of petroleum, which usually cures the trouble in six days. He calls attention to its

power as a disinfectant, while it does not irritate mucous membranes, though it may cause vesication (but not sloughing) of the already inflamed skin.

UNUSUALLY LARGE OVARIAN TUMOR.

MARITAN (*Gaz. Méd. de Paris*, 1833, No. 2) removed an ovarian cyst weighing two hundred pounds from a patient whose weight before the operation was two hundred and ninety pounds, her circumference at the umbilicus being ninety inches. The length of the abdominal incision was about twenty-eight inches. The patient suffered from collapse, but made a good recovery.

FOREIGN BODIES IN THE ABDOMINAL CAVITY.

ANONYMUS (*Revue des Mal. des Femmes*, 1892, No. 4) reports several cases in which foreign bodies were left in the abdominal cavity after celiotomy. In one instance a gauze compress a foot long was discharged per rectum eight months after myomectomy had been performed, the patient having no trouble until four months after the operation. In another a strip of iodoform gauze, fifteen inches long, was withdrawn from the intestines of a young girl who had undergone salpingotomy. A month later vaginal hysterectomy was performed, as she had not been relieved by the first operation. Several months after, a secondary celiotomy was performed on account of persistent abdominal pain. In endeavoring to separate an adherent loop of intestine, which was supposed to be the seat of the trouble, it was accidentally ruptured, when the gauze protruded. Five inches of the gut were excised, and the patient eventually recovered after having a fecal fistula. In two other cases a sponge and a forceps were missed soon after the patient had been placed in bed. In both cases the patient was returned to the operating-room, the wound reopened and the missing objects were found in the abdominal cavity.

TUBAL PREGNANCY; DIFFICULTY OF DIFFERENTIAL DIAGNOSIS.

PICHEVIN (*Nouv. Arch. d'Obstér. et de Gynéc.*, 1892, No. 9) reports the case of a woman, aged forty years, who had always menstruated regularly. Her period was delayed twelve days; then she had an irregular flow for several weeks; after this had ceased for two weeks, she had a sudden attack of severe abdominal pain, without collapse. The hemorrhages ceased and then returned, persisting for three or four weeks. The patient then presented the ordinary mammary signs of pregnancy; the cervix uteri was soft and patulous, and the fundus extended three inches above the symphysis. To the left of the uterus could be felt a tumor, which dipped down into Douglas' pouch and extended upward into the abdomen; no fluctuation could be detected. On opening the abdomen two ovarian cysts were found (one tubo-ovarian?), but no sign of ectopic gestation.

NEW OPERATION FOR PROLAPSUS.

FREUND (*Centralblatt für Gynäkologie*, 1893, No. 47) describes a new operation for prolapsus which he has performed in five cases with satisfactory

results. Its advantages are rapidity of execution, as there is no denudation, painlessness (so that it can be performed without an anæsthetic), and the fact that there are no sutures to be removed. It is especially applicable to old women, who have neither hypertrophy of the cervix, destruction of the pelvic floor, nor tumor, etc., requiring the usual operations. It should not be undertaken if the cervix or vaginal wall is ulcerated.

The operation consists essentially in inserting three or four purse-string sutures (as in Stoltz's method) of silver wire. A curved needle is first entered beneath the vaginal mucosa at a point just above the portio, and is run in and out, emerging at the point of entrance. As this suture is tightened the cervix is pushed upward (as in Lefort's operation), and the wire is twisted. A second circular suture is inserted an inch above the first, and is twisted until the calibre of the vagina at that point is narrowed to the size of the finger, then a third suture in the same manner, the last being placed close to the remains of the hymen. The latter must not be twisted so tight as to effect *kolpokleisis*. The operation is bloodless, takes only a few minutes, and is well borne by old patients, who need be kept in bed only one day. In one case the patient, who had a large cystocele and rectocele, still menstruated and performed her matrimonial duties. There was no recurrence of the prolapse with severe muscular efforts. The sutures are not removed at all.

USE OF CLAMPS IN VAGINAL EXTIRPATION OF THE UTERUS.

LANDAU (*Berliner klin. Wochenschrift*, 1893, Nos. 24-26) reports seventy-one cases, with five deaths, in all of which he secured the broad ligaments with clamps instead of using ligatures. The technique differs according as the uterus is perfectly movable or the reverse. In the former case the uterus is anteфлекed, after incising the vesico uterine fold of peritoneum, and is drawn down forcibly, so that the finger can be hooked over the left broad ligament, which is then secured with one or two clamps and is divided, when it is an easy matter to clamp the ligament on the right side. The ovaries and tubes are afterward drawn down and clamped off if desired. Bleeding vessels on the edge of the vaginal wound are caught with forceps. A gauze tent is pushed up into the opening and the operation is concluded, the average time in an easy case being only ten minutes. When the uterus is fixed by perimetritic indurations the organ is drawn forcibly downward, while the broad ligaments are clamped in sections and divided. If ovarian cysts or pus tubes are encountered, these are pulled down into the wound and incised outside of the peritoneal cavity, then clamped off. In complicated cases, when there is extreme fixation, Péan's method of removal by *morcellement* is employed. A catheter is left in the bladder until after the clamps are removed, *i. e.*, for thirty-six or forty-eight hours. Two or three vaginal douches are given daily after the latter have been removed. The patient is usually out of bed by the eighteenth day. The writer denies that there is any more danger of sepsis with the clamps than with ligatures, nor is there any added risk of injuring adjacent viscera. He has no fear of secondary hemorrhage if the clamps are left *in situ* for thirty-six hours, nor has he noted that the patients complain of more pain than when ligatures were used.