which agrees with Darboux's solution on putting

$$P = (u^2 - 1)/2,$$

$$Q = i(u^2 + 1)/2,$$

$$R = u,$$

for then we have $(l + im)u^2 + 2nu + (-l + im) = 0$

$$\text{or} \quad u = \frac{\pm r - n}{l + im} = \frac{l - im}{\pm r + n},$$

$$\text{also} \quad lP'' + mQ'' + nR'' = l + im,$$

$$lP' + mQ' + nR' = \pm r,$$

and on substitution these will give the solution quoted.

On some Solutions of $\nabla^2 v = 0$. By T. J. I'A. BROMWICH,
St. John's College, Cambridge. Received February 6th,
1899. Read February 9th, 1899.

Prof. Forsyth in his paper on this subject (*Mess. of Math.*, Vol. xxvii.) shows how to solve $\nabla^2 v = 0$ and allied equations when the solution of a certain functional equation is known. I recall briefly his results, stating them for three variables only, though, as he shows, the solutions hold for any number of variables.

He takes p, q, r three functions of a variable u such that

$$p^2 + q^2 + r^2 = 0,$$

and u is given by $au = px + qy + rz,$

where a is an arbitrary constant.

He then shows that we may take, as a solution of $\nabla^2 v = 0$,

$$v = f(u) + \frac{1}{\Delta} g(u),$$

where f, g are perfectly arbitrary functions and

$$\Delta = a - p'x - q'y - r'z.$$

Here, as throughout my work, accents indicate differential coefficients with respect to u .

It became clear to me in the course of the investigation contained in the preceding paper that solutions ought to exist of the type

$$v = Xf(u) + Yf'(u),$$

where X, Y are not necessarily functions of u alone. I now show how to effect a determination of the forms of X, Y .

On differentiation we have

$$\begin{aligned} \frac{\partial^2 v}{\partial x^2} = \frac{\partial^2 X}{\partial x^2} f + \left[2 \frac{\partial u}{\partial x} \frac{\partial X}{\partial x} + X \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 Y}{\partial x^2} \right] f' \\ + \left[X \left(\frac{\partial u}{\partial x} \right)^2 + 2 \frac{\partial u}{\partial x} \frac{\partial Y}{\partial x} + Y \frac{\partial^2 u}{\partial x^2} \right] f'' + Y \left(\frac{\partial u}{\partial x} \right)^2 f'''. \end{aligned}$$

Now, on addition of this and two similar terms, we get

$$\begin{aligned} \nabla^2 v = (\nabla^2 X) f + \left[2 \left(\frac{\partial u}{\partial x} \frac{\partial X}{\partial x} + \frac{\partial u}{\partial y} \frac{\partial X}{\partial y} + \frac{\partial u}{\partial z} \frac{\partial X}{\partial z} \right) + \nabla^2 Y \right] f' \\ + 2 \left[\frac{\partial u}{\partial x} \frac{\partial Y}{\partial x} + \frac{\partial u}{\partial y} \frac{\partial Y}{\partial y} + \frac{\partial u}{\partial z} \frac{\partial Y}{\partial z} \right] f'', \end{aligned}$$

for, as shown by Prof. Forsyth, we have

$$\nabla^2 u = 0$$

and

$$\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 + \left(\frac{\partial u}{\partial z} \right)^2 = 0.$$

Hence, to get a solution of the type indicated, we may take

$$\nabla^2 X = 0,$$

$$\nabla^2 Y + 2 \left(\frac{\partial u}{\partial x} \frac{\partial X}{\partial x} + \frac{\partial u}{\partial y} \frac{\partial X}{\partial y} + \frac{\partial u}{\partial z} \frac{\partial X}{\partial z} \right) = 0,$$

$$\frac{\partial u}{\partial x} \frac{\partial Y}{\partial x} + \frac{\partial u}{\partial y} \frac{\partial Y}{\partial y} + \frac{\partial u}{\partial z} \frac{\partial Y}{\partial z} = 0.$$

In the application originally contemplated, it seemed that Y would be a function of Δ only, and I restrict myself to this case.

Taking then

$$Y = F(\Delta),$$

we have that the third equation of condition is at once satisfied, for

$$\frac{\partial u}{\partial x} \frac{\partial \Delta}{\partial x} + \frac{\partial u}{\partial y} \frac{\partial \Delta}{\partial y} + \frac{\partial u}{\partial z} \frac{\partial \Delta}{\partial z} = 0.$$

Next $\nabla^2 Y = \left[\left(\frac{\partial \Delta}{\partial x} \right)^2 + \left(\frac{\partial \Delta}{\partial y} \right)^2 + \left(\frac{\partial \Delta}{\partial z} \right)^2 \right] F''(\Delta) + (\nabla^2 \Delta) F'(\Delta),$

and we have $-\frac{\partial \Delta}{\partial x} = p' + \frac{p}{\Delta} (xp'' + yq'' + zr''),$

on using the result $\frac{\partial u}{\partial x} = \frac{p}{\Delta}.$

Hence we have

$$\left(\frac{\partial \Delta}{\partial x} \right)^2 + \left(\frac{\partial \Delta}{\partial y} \right)^2 + \left(\frac{\partial \Delta}{\partial z} \right)^2 = p'^2 + q'^2 + r'^2,$$

and we find, on differentiating again and adding,

$$\begin{aligned} \nabla^2 \Delta &= -2 (pp'' + qq'' + rr'')/\Delta \\ &= 2 (p'^2 + q'^2 + r'^2)/\Delta, \end{aligned}$$

for

$$pp' + qq' + rr' = 0,$$

and, differentiating again, we find

$$(p'^2 + q'^2 + r'^2) + (pp'' + qq'' + rr'') = 0.$$

Thus $\nabla^2 Y = (p'^2 + q'^2 + r'^2) \left[F''(\Delta) + \frac{2}{\Delta} F'(\Delta) \right].$

Next, for a type to be taken as X , I use

$$X = xP + yQ + zR,$$

where P, Q, R are also functions of u only. We then have, on differentiation,

$$\begin{aligned} \frac{\partial X}{\partial x} &= P + (xP' + yQ' + zR') \frac{\partial u}{\partial x}, \\ \frac{\partial^2 X}{\partial x^2} &= 2P' \frac{\partial u}{\partial x} + (xP'' + yQ'' + zR'') \left(\frac{\partial u}{\partial x} \right)^2 \\ &\quad + (xP' + yQ' + zR') \frac{\partial^2 u}{\partial x^2}. \end{aligned}$$

Thus $\nabla^2 X = 2 \left(P' \frac{\partial u}{\partial x} + Q' \frac{\partial u}{\partial y} + R' \frac{\partial u}{\partial z} \right),$

the other terms vanishing as before, and this expression

$$= 2 (Pp' + Qq' + Rr')/\Delta.$$

Also
$$\frac{\partial X}{\partial x} \frac{\partial u}{\partial x} + \frac{\partial X}{\partial y} \frac{\partial u}{\partial y} + \frac{\partial X}{\partial z} \frac{\partial u}{\partial z} = P \frac{\partial u}{\partial x} + Q \frac{\partial u}{\partial y} + R \frac{\partial u}{\partial z}$$

$$= (Pp + Qq + Rr)/\Delta;$$

whence we have that the form assumed for v will give a solution provided

$$P'p + Q'q + R'r = 0$$

and $2(Pp + Qq + Rr) + (p'^2 + q'^2 + r'^2) [2F'(\Delta) + \Delta F''(\Delta)] = 0.$

Now Δ cannot be expressed as a function of u only, but $(Pp + Qq + Rr)$ and $(p'^2 + q'^2 + r'^2)$ both involve no other variable than u ; consequently we must have

$$\Delta F''(\Delta) + 2F'(\Delta) = 2A,$$

where A is a constant.

Hence
$$F(\Delta) = A\Delta + B + C/\Delta,$$

by integration, B, C being constants of integration. Thus we now have only to satisfy

$$P'p + Q'q + R'r = 0$$

and
$$Pp + Qq + Rr = -A(p'^2 + q'^2 + r'^2).$$

Differentiating the second of these and using the first, we find

$$Pp' + Qq' + Rr' = -2A(p'p'' + q'q'' + r'r'').$$

Now, since
$$p^2 + q^2 + r^2 = 0,$$

$$pp' + qq' + rr' = 0,$$

we have
$$\frac{qr' - q'r}{p} = \frac{rp' - r'p}{q} = \frac{pq' - p'q}{r} = \kappa, \text{ say,}$$

and then
$$p'^2 + q'^2 + r'^2 = [(r'p + \kappa q)^2 + (r'q - \kappa p)^2 + r^2 r'^2]/r^2$$

$$= -\kappa^2.$$

Hence the equations for P, Q, R are

$$Pp + Qq + Rr = A\kappa^2,$$

$$Pp' + Qq' + Rr' = 2A\kappa\kappa,$$

and
$$Px + Qy + Rz = X.$$

These give, on eliminating P, Q ,

$$\begin{vmatrix} p & q & A\kappa^2 - Rr \\ p' & q' & 2A\kappa\kappa' - Rr' \\ x & y & X - Rz \end{vmatrix} = 0$$

$$\text{or} \quad (X - Rz)(pq' - p'q) + (A\kappa^2 - Rr)(p'y - q'x) \\ + (2A\kappa\kappa' - Rr')(qx - py) = 0.$$

Here the coefficient of R is

$$-\begin{vmatrix} p & q & r \\ p' & q' & r' \\ x & y & z \end{vmatrix},$$

which is

$$-\kappa(px + qy + rz) = -\kappa au.$$

Hence we have

$$\kappa rX - \kappa auR + A\kappa^2(p'y - q'x) + 2A\kappa\kappa'(qx - py) = 0,$$

$$\text{or} \quad rX = auR + A\kappa(q'x - p'y) + 2A\kappa'(py - qx).$$

This solution for X is not symmetrical; a symmetrical form can be deduced by the addition of three such results, but their sum does not appear to simplify to any extent.

Hence a solution of $\nabla^2 v = 0$

$$\text{is} \quad v = \frac{1}{r} [auR + A\kappa(q'x - p'y) + 2A\kappa'(py - qx)] f(u) \\ + (A\Delta + B + O/\Delta) f'(u),$$

and it is clear that the terms in B, C are included in Prof. Forsyth's types of solution; further, since $uRf(u)/r$ is an arbitrary function of u , this term may be considered as also included in his solutions.

Thus the investigation indicates a solution of the new type

$$v = \left[\frac{\kappa}{r} (q'x - p'y) + 2 \frac{\kappa'}{r} (py - qx) \right] f(u) + \Delta f'(u).$$

Turning to the case worked out by Prof. Forsyth, we may take

$$p = (u^2 - 1)/2,$$

$$q = i(u^2 + 1)/2,$$

$$r = u,$$

and then

$$u = (a - z + \rho) / (x + iy),$$

where

$$\rho = \pm \sqrt{(a - z)^2 + x^2 + y^2}.$$

We find

$$\Delta = -\rho \quad \text{and} \quad \kappa = -i,$$

so that

$$v = (x + iy) f \left(\frac{a - z + \rho}{x + iy} \right) - \rho f' \left(\frac{a - z + \rho}{x + iy} \right). \quad (\text{A})$$

It will be seen that, if we differentiate with respect to x any of the results found, we shall get solutions of

$$\nabla^2 v = 0.*$$

Consider then
$$\frac{\partial}{\partial x} f(u) = f'(u) \frac{\partial u}{\partial x} = \frac{1}{\Delta} p f'(u).$$

But, since p is a function of u , this simply leads to Prof. Forsyth's second type of solution. Differentiating again, we have

$$\begin{aligned} \frac{\partial}{\partial x} \frac{f(u)}{\Delta} &= \frac{f'(u)}{\Delta} \frac{p}{\Delta} + \frac{f(u)}{\Delta^2} \left[p' + \frac{p}{\Delta} (xp'' + yq'' + zr'') \right] \\ &= \frac{1}{\Delta^2} \left[g'(u) \Delta + (xp'' + yq'' + zr'') g(u) \right], \end{aligned}$$

where

$$g(u) = p f'(u).$$

Now, on substituting the particular set of values for p, q, r given above, we find that this solution is of the form

$$\frac{1}{\rho^3} \left[(x + iy) g(u) - \rho g'(u) \right].$$

Consequently the type (A) is of the form

$$\rho^3 \frac{\partial^2}{\partial x^2} f(u)$$

in the particular case just referred to. Of course this type of solution could be anticipated by the general result that, if v be a solution of zero order, then $\rho^{2n+1} \frac{\partial^{n+1} v}{\partial x^{n+1}}$ is a solution of order n .

* Note added 15th April.—One of the referees remarks that Prof. Burnside has alluded to the series of solutions so found in a paper "On Equipotentials" in the volume of the *Messenger* which contains Prof. Forsyth's paper.

On trying to generalize this type of solution, it seemed probable that the form to use would be

$$\Delta^3 \frac{\partial^3}{\partial x^3} f(u),$$

but it will be found that this will only satisfy the conditions found above when

$$p'p'' + q'q'' + r'r'' = 0,$$

or when

$$p'^2 + q'^2 + r'^2 = \text{const.}$$

It will always be possible to satisfy this condition, for, if the values of p, q, r as originally found do not lead to the result, consider

$$\left[\frac{d}{du} (p\theta) \right]^2 + \left[\frac{d}{du} (q\theta) \right]^2 + \left[\frac{d}{du} (r\theta) \right]^2,$$

where θ is any function of u ; this will be seen to be

$$\theta^2 (p'^2 + q'^2 + r'^2),$$

and hence, if we take

$$\theta = i/\kappa,$$

we shall find that

$$p_1 = ip/\kappa, \quad q_1 = iq/\kappa, \quad r_1 = ir/\kappa$$

will satisfy

$$p_1'^2 + q_1'^2 + r_1'^2 = 1.$$

I shall now attempt to find in what cases $\Delta^{2n+1} \frac{\partial^{n+1}}{\partial x^{n+1}} f(u)$ is a solution of Laplace's equation. To attack this, it will be simplest to find the condition that

$$\nabla^2 (\Delta^m Z) = 0,$$

when

$$\nabla^2 Z = 0.$$

On performing the differentiation, we find that

$$(m+1) Z + 2\Delta \left(\frac{\partial \Delta}{\partial x} \frac{\partial Z}{\partial x} + \frac{\partial \Delta}{\partial y} \frac{\partial Z}{\partial y} + \frac{\partial \Delta}{\partial z} \frac{\partial Z}{\partial z} \right) = 0.$$

Now, when Z is $\frac{\partial^{n+1}}{\partial x^{n+1}} f(u)$, it will contain x explicitly and also implicitly in u, Δ ; in fact,

$$Z = \phi(x, y, z, u, \Delta),$$

and

$$\frac{\partial Z}{\partial x} = \frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial \phi}{\partial \Delta} \frac{\partial \Delta}{\partial x},$$

where $\frac{\partial \phi}{\partial x}$ is to be understood as derived only from the terms explicitly containing x .

On substituting this and the similar expressions in the condition obtained previously, we get

$$\frac{m+1}{2\Delta} \phi + \frac{\partial \phi}{\partial \Delta} + \left(\frac{\partial \Delta}{\partial x} \frac{\partial \phi}{\partial x} + \frac{\partial \Delta}{\partial y} \frac{\partial \phi}{\partial y} + \frac{\partial \Delta}{\partial z} \frac{\partial \phi}{\partial z} \right) = 0,$$

for
$$\frac{\partial u}{\partial x} \frac{\partial \Delta}{\partial x} + \frac{\partial u}{\partial y} \frac{\partial \Delta}{\partial y} + \frac{\partial u}{\partial z} \frac{\partial \Delta}{\partial z} = 0,$$

and
$$\left(\frac{\partial \Delta}{\partial x} \right)^2 + \left(\frac{\partial \Delta}{\partial y} \right)^2 + \left(\frac{\partial \Delta}{\partial z} \right)^2 = p'^2 + q'^2 + r'^2 = 1,$$

it being supposed that p, q, r are determined as just explained.

Now we have seen that

$$\frac{\partial^3}{\partial x^3} f(u) = \frac{g'(u)}{\Delta^3} + \frac{g(u)}{\Delta^3} (p''x + q''y + r''z),$$

so consider the next differential coefficient,

$$\begin{aligned} \phi = \frac{\partial^3}{\partial x^3} f(u) &= \frac{h''(u)}{\Delta^3} + \frac{3h'(u)}{\Delta^4} (p''x + q''y + r''z) \\ &\quad + \frac{h(u)}{\Delta^5} [3(p''x + q''y + r''z)^2 + (p'''x + q'''y + r'''z)\Delta], \end{aligned}$$

where we put

$$h(u) = pg(u),$$

and we now find that

$$\begin{aligned} -\frac{\partial \phi}{\partial \Delta} - \frac{\partial \Delta}{\partial x} \frac{\partial \phi}{\partial x} - \frac{\partial \Delta}{\partial y} \frac{\partial \phi}{\partial y} - \frac{\partial \Delta}{\partial z} \frac{\partial \phi}{\partial z} \\ = \frac{3h''(u)}{\Delta^4} + \frac{9h'(u)}{\Delta^5} (p''x + q''y + r''z) + \frac{9h(u)}{\Delta^5} (p''x + q''y + r''z)^2 \\ + \frac{h(u)}{\Delta^5} [4(p'''x + q'''y + r'''z) + \Delta(p'p''' + q'q''' + r'r''')], \end{aligned}$$

where we have used the fact that here

$$pp''' + qq''' + rr''' = 0,$$

for, since $p'^2 + q'^2 + r'^2 = 1 = -(pp'' + qq'' + rr'')$,

we have

$$p'p'' + q'q'' + r'r'' = 0$$

and $(p'p'' + pp''') + (q'q'' + qq''') + (r'r'' + rr''') = 0$,

giving $pp''' + qq''' + rr''' = 0$.

Returning to the condition that $\phi \Delta^m$ may be a solution of $\nabla^2 v = 0$, it will be seen that $m+1 = 6$ will satisfy the condition, provided

$$p'''x + q'''y + r'''z + \Delta (p'p''' + q'q''' + r'r''') = 0.$$

Hence $\frac{p'''}{p'} = \frac{q'''}{q'} = \frac{r'''}{r'} = p'p''' + q'q''' + r'r'''$

and $a(p'p''' + q'q''' + r'r''') = 0$,

Thus either $p''' = 0$, $q''' = 0$, $r''' = 0$; or we may take $a = 0$, and then $\frac{p'''}{p'} = \frac{q'''}{q'} = \frac{r'''}{r'}$ will satisfy all the conditions.

In the first case p, q, r are each quadratic functions of u , and here the most general values can be only linear functions of the particular set

$$p = (u^2 - 1)/2, \quad q = i(u^2 + 1)/2, \quad r = u;$$

and the solution of $\nabla^2 v = 0$ so found can be deduced from that particular case by a transformation of coordinates.

In the second case, when $a = 0$, the functions involved are all homogeneous in x, y, z , and we can apply the ordinary result of spherical harmonics, that, if Z be a solution of $\nabla^2 v = 0$, homogeneous in x, y, z of degree $-(n+1)$, then the corresponding solution of degree n is $(x^2 + y^2 + z^2)^{n+1} Z$.

We are thus led to the conclusion that the form $\Delta^{2n+1} \frac{\partial^{n+1}}{\partial x^{n+1}} f(u)$

for values of $n > 1$ is a solution of $\nabla^2 v = 0$ only when p, q, r are quadratic functions of u ; and that, in the particular case of $a = 0$, we may use the type $(x^2 + y^2 + z^2)^{n+1} \frac{\partial^{n+1}}{\partial x^{n+1}} f(u)$. It will be seen,

moreover, that the first case is virtually included in the second, for we can make a zero by shifting the origin, when p, q, r are only quadratic functions of u , and then

$$\Delta^2 = x^2 + y^2 + z^2.$$

Thus the only new form we can find in this way would seem to be the one first considered, namely,

$$v = \left[\frac{\kappa}{r} (q'x - p'y) + \frac{2\kappa'}{r} (py - qx) \right] f(u) + \Delta f'(u),$$

which is the same as

$$v = (p''x + q''y + r''z)f(u) + \Delta f'(u),$$

when we make

$$p'^2 + q'^2 + r'^2 = 1.$$

The other forms investigated will be the same as those given by Sommerfeld in a paper "On Diffraction," in the *Mathematische Annalen* (t. XLVII., 1896), in the form of contour integrals, where they are deduced by a stereographic projection.

The Irreducible Concomitants of any Number of Binary Quartics.

By A. YOUNG. Received and read February 9th, 1899.

The irreducible system is here arrived at by first finding the irreducible system of types and then the number of independent forms belonging to each type for a system of N quartics. Two concomitants are said to be of the same type when they can be obtained from the same form by polarization. For the purpose of discussing the system of types, a type is regarded as being of the first degree in the coefficients of each of the quartics concerned. The finiteness of the irreducible system of types has been established by Prof. Peano.* He proves that the complete system of concomitants for any number of binary n -ics may be obtained from the system for n n -ics by polarization alone; with the one possible exception of invariants of the type

$$\begin{vmatrix} A_0 & A_1 & \dots & A_n \\ B_0 & B_1 & \dots & B_n \\ \dots & \dots & \dots & \dots \\ K_0 & K_1 & \dots & K_n \end{vmatrix}.$$

In other words, every type of a binary n -ic which furnishes no irreducible form for n n -ics is reducible, with the possible exception just mentioned. It was with the help of this proposition that some of the reductions for the quartic were first arrived at; however, other

* *Atti di Torino*, t. XVII., p. 580.