safety, and at an early period, in a large number of cases.

I must not, however, forget that the title of this paper is Arthrectomy *v.* Excision, and if I have spoken much in favour of the plaintiff in this action, I would not have it thought that the defendant has no case. Excision of the knee shares with arthrectomy precisely the same advantages that are to be derived from our modern improvements in wound treatment, and the risks of septicæmia and pyæmia, or of the necessity of later amputation because of sup-puration, are greatly lessened. I know of no recent statistics on this point, but no one will question, I think, that the immediate results of excision have, with all other operations immensely improved in the last ten years. But no antiseptics, nor any healing by first intention, can lessen the chances of arrested growth in the limb when excision is performed in young people; and that operation, in my opinion, must be reserved for cases in which the knee has become ankylosed in a flexed position, or in which, on proceeding to the operation of arthrectomy, it is found that the disease of the bone is so extensive that mere scraping and gouging are insufficient for its removal. This was what happened in a case which I saw last week under the care of my colleague Mr. Pepper; and I operated myself on the same day on the 'knee of a man aged twenty-seven, with a history of disease of five years' standing, where the operation performed was neither an arthrectomy nor an excision, but a combination of the two. The surgeon must be prepared for this sort of thing, and it may even be that he will find conditions which contra-indicate the possibility of success from arthrectomy or from an arthrectomy excision, and amputation had better be done at once. Thus in one case—a lad of fourteen—there was found such a rarefying osteitis of the lower end of the femur that the femur broke three inches above the condyles during the manipulation entailed by the operation; and in another the bursa under the popliteus was found extensively diseased, and practically forming a large abscess cavity in the calf. Conditions such as these not only call for amputation, for which therefore the patient ought to be prepared: they show how much better it would be to prevent such things arising by early operative interference, when a mere arthrectomy may suffice. Surely everything points to the advisability of early operation, unless there be the most distinct evidence of improvement from rest. We do not want to have to do an operation at all like that of the old excision. Let us operate at a time when we shall have to sacrifice no more than is absolutely necessary for the removal of disease, when by such removal we shall place the tissues of the joint in a condition in which repair, once started, is sure to go on, when we can preserve the growing limb in its natural length, and, by improving the general health, shall fortify the patient against the possible inroads of future disease. These are inestimable benefits, and the more they are known the better will it be.

SACCHARIN.

By THOMAS STEVENSON, M.D., F.R.C.P. LOND.,
LECTURER ON FORENSIC MEDICINE AND ON CHEMISTRY AT
GUY'S HOSPITAL,
AND

L. C. WOOLDRIDGE, M.D., D.Sc., M.R.C.P. LOND.,
ASSISTANT PHYSICIAN AND CO-LECTURER ON PHYSIOLOGY AT
THE HOSPITAL.

IN view of the enormous increase in the use of saccharin for the purpose of sweetening foods, beverages, and medicines, we have been led to institute experiments, with the view of determining whether this substance is poisonous or not when given even in excessive quantities; and, if not poisonous under these or other conditions, whether its use in moderation interferes with the digestive processes, so as to render it advisable to forbid its use as a substitute for sugar. That saccharin is not a food is manifest; but there are so many circumstances under which the use of a sweetening agent in place of sugar is desirable, as to render it advisable to determine, if possible, whether the dietetic use of saccharin is innocuous.

1. As to the non-toxic nature of saccharin we have no

doubt. Reliable continental experimenters have admitted this, and we have confirmed their observations. To an underfed dog we gave daily for five days two grammes of saccharin (equal in sweetening power to more than a pound of sugar), in addition to food *ad libitum*. The animal increased in weight, and no inconvenient results were observed. We mixed large quantities of saccharin with the food of mice; they ate of the mixture *ad libitum*, freely, and for a considerable period, and in no single case was there any manifest action on the health of the animal.

2. As saccharin has decided antiseptic properties, and is capable, in sufficient quantities, of stopping the action of organised ferments, we ascertained its extra-corporeal action on the soluble ferments, and found that in respect of the peptic digestion of fibrin 0.1 per cent. of saccharin has no retarding influence, whilst 0.25 per cent. slows the process decidedly, and 1 per cent. greatly retards it: 0.1 per cent. of saccharin is the equivalent of 30 per cent. of sugar—an impossible dietetic quantity. The diastatic solution of starch was not hindered by 2 per cent. of saccharin. The ammoniacal fermentation of urine is retarded when saccharin is added to it, or when saccharin is taken—an important fact in clinical medicine. The putrefactive decomposition of a pancreatic digestive mixture we found was not prevented by the admixture of saccharin to the extent of 1 per cent.

Experiments on the extra-corporeal action of saccharin on the digestive ferments are, however, of no significance in determining the effects which saccharin would have when taken with food, since they involve the maintenance of a constant fairly strong solution of saccharin in the digestive medium—a condition which does not obtain when saccharin is taken, since this substance is quickly absorbed, and is excreted in the urine.

The following experiment was made for the purpose of exactly determining whether saccharin influences gastric digestion. Two similar dogs, after having fasted for thirty hours, were each fed with 300 grammes (about 11 oz.) of lean beefsteak, equally cut up and divided. The 300 grammes of steak contained 69 grammes (i.e., 23 per cent.) of dry solids. No. 1 dog weighed 21 lb., and had no saccharin; No. 2 dog weighed 15½ lb., and had one gramme of saccharin (equal in sweetening power to more than 8 oz. of sugar) with his meat. Five hours and a half after the meal the dogs were killed with chloroform, and the contents of the stomach and intestines carefully removed from each animal. The small intestine in each case was empty, except the duodenum which contained a small quantity of digestive mixture that had passed from the stomach; and at the lower end, just above the large intestine, there was material obviously of older duration. In each dog the stomach contained some undigested meat. The dry weight of the contents of the stomach and duodenum of No. 1 dog was 23 grammes, or 33 per cent. of the dry meat taken; whilst the similar dry weight in No. 2 dog (which had taken saccharin) was 21 grammes, or 30 per cent. of the dry meat taken. Obviously, the ingestion of a gramme of saccharin, equal in sweetening power to over 8 oz. of sugar, had not in the least interfered with the gastric digestion of the dog. One of us has, moreover, taken considerable quantities of saccharin daily without experiencing any ill effect. The saccharin used by us was "soluble saccharin," which is equal in power to about nine-tenths of its weight of commercial "pure saccharin." "Soluble saccharine" is as soluble in water as table salt, and when appropriately diluted is perhaps indistinguishable in taste from cane sugar, and is free from all the bitterness of flavour met with in the soluble saccharin first brought into the market.

Summary.—We conclude that (1) saccharin is quite innocuous when taken in quantities largely exceeding what would be taken in any ordinary dietary; (2) saccharin does not interfere with or impede the digestive processes when taken in any practicable quantity; and (3) our personal experience is that saccharin may be taken for an extended period without interfering with the digestive and other bodily functions. Hence there is no reason to think that its continued use is in any way harmful.

Guy's Hospital.

LADY DUFFERIN'S FUND.—It is stated that Lady Lansdowne, on her arrival in India, will take over the presidentship of this fund; and it is expected that, besides the capital, the balance in hand will amount to 30,000 rupees.