A CASE OF CONSERVATIVE CÆSAREAN OPERATION.

DOCTOR (*Centralblatt für Gynäk.*, 1893, No. 27) reports an interesting case in which the Cæsaean operation was done on a primipara, thirty-two years old, of undeveloped appearance. The height of the patient was 132 cm., her deformity consisting in an angular lumbo-sacral kyphosis at the fifth lumbar vertebra. Sacrum convex posteriorly. The transverse outlet was also much narrowed, two fingers being unable to find entrance between the ischia. The abdomen was opened when labor began. The uterus was brought out through the abdominal wound, an assistant grasping the uterine neck so as to compress it and hold it upright. The incision was made from the fundus to the neck, through the placenta, and the child quickly extracted. Severe hemorrhage was encountered from the placental site. The uterus was closed by silk sutures passed through its entire thickness about $\frac{3}{4}$ cm. apart, the abdominal sutures being of silkworm-gut. The strictest asepsis was maintained. Before the operation two injections of ergotine were employed, yet even with this and the manipulations, no active contraction of the uterus was observed; the retraction, however, was satisfactory, and the volume reduced to that of a recently delivered uterus. The total hemorrhage was not more than is found in the ordinary labor.

Recovery was prompt; little lochial discharge was present. The writer draws attention to the following points:

1. The operation was done at a most convenient time.
2. To arrest bleeding no elastic band was used, and no compression with the fingers.
3. The wound was closed with the simplest sutures. The arrest of hemorrhage must have been from uterine contraction.
4. Absolute asepsis.

A METHOD OF REDUCTION OF THE COMPLETELY RETROVERTED GRAVID UTERUS BY THE APPLICATION OF A PESSARY.

LEVAAT (*Le Mercredi Médical*, No. 30, 1893) describes a method for the replacement of the gravid uterus by manual means, the operation being that used for several years by Laroyenne. The manipulation consists in passing the hand between the uterus and promontory, avoiding all pressure upon the former, this being most easily done by following the lateral wall of the pelvis to the sacro-vertebral angle. When the hand has arrived here, the uterus is generally found partially enucleated from the promontory, but often the reduction is incomplete. The first stage having been successfully accomplished, the second consists in placing in position an appropriate pessary. This should have the form of a Hodge ring, with the posterior arc very large and of such a shape as not to press on any one point of the uterus. The ring should occupy the gutter made by the hand of the operator, between the uterus and sacrum. In these conditions, after a lapse of twenty-four hours or so, the reduction is complete. Two symptoms indicate complete restitution of the uterus, namely, cessation of the retention of urine and the restoration of the utero-placental souffle. The pessary may be withdrawn after four or five days from the time of the restitution of the uterus. The operation may require anæsthesia.

BRINGING DOWN THE FOOT IN BREECH PRESENTATIONS.

BONNAIRE, in a lecture on the above subject (*La Semaine Médicale*, 1893, No. 50), says: Cazeau observes in such cases that the great difficulty may be considered as being that a straight inflexible mass must pass through a curved canal little longer than itself (30 to 35 cm. against that of child, 20 cm.). Under favorable conditions of breech presentations with relaxed perineum, a very small child, or when the fetus is macerated, the delivery is easy; but when the opposite obtains, the birth is one of dystocia. In such cases it is useless to worry with filets or hooks, but remove the rigidity by taking from the fetal body its props—the legs. To do this one should draw down the anterior—the easier of access—and allow the other to glide across the belly. The body then regains its suppleness. The movement of extraction can be performed above the superior strait even, by compressing the leg, bending it on the thigh, extending the whole, and finally obtaining complete lowering of the foot. Once the leg and foot are lowered in the inferior segment of the uterus and vagina, the vertebral column of the infant ceases to be bridled laterally. The author would distinguish, besides the above manœuvre, a partial drawing down of the turned-up member—simply flexing the leg on the thigh. In the above methods prolapse of the cord must be borne in mind. Bringing down both feet is possible only if the presentation is above the pelvic excavation, or partially beyond and without the inferior strait. The manœuvre ought to be only used during labor, provided the cervix be sufficiently dilated to let in three fingers of the examining hand. The integrity of the bag of waters constitutes a contra-indication in the interests of the child. This method should not be used before the eighth month of pregnancy, or if the child be dead. It is also useless in cases in which great relaxation of the pelvic floor exists. Bringing down one foot risks less a prolapse of the cord, and is easier to perform.

PUERPERAL APHASIA.

CARRE (*Archives de Tocologie et de Gynécologie*, 1893, vol. xx., No. 7) reports four cases of puerperal aphasia.

In the first the patient presented a cardiac trouble, which developed during her second confinement, and perhaps contemporary with a phlebitis following her first labor. Under the influence of lively emotion, a clot from the heart entered the Sylvian artery and temporarily cut off the blood-supply. The embolism gradually disappeared and the centres recovered their functions, cerebral paresis remaining.

In the second case there was marked manifestation of embolism limited to the left hemisphere, and due to impoverishment of the blood. In this case, two weeks after confinement, the lachia and milk suddenly ceased. Thirty-six hours later complete aphasia set in, the only words which the patient could speak being "yes" and "no." Profound coma, succeeded by fever, and convulsions of the right side. These symptoms soon disappeared.

Case III. was a primipara with a normal delivery. Nineteen days after parturition she developed headache and otorrhœa, accompanied by fever and photophobia. There was aphasia of reception and transmission. These

symptoms disappeared gradually, but five months after the onset the patient still has observable weakness of memory and intelligence. The aphasia was due to thrombosis, favored by the puerperal state and provoked by inflammatory lesions of the middle ear.

In the fourth case, eight days after delivery there appeared right hemiplegia, with aphasia and inability to protrude the tongue. Temperature normal. Intelligence seemed to be preserved. No abdominal or cardiac complications. Six weeks after confinement death occurred. In this case labor had been easy, but lactation had been excessive; and in the absence of cardiac or rheumatic symptoms, the cerebral accidents are to be attributed rather to thrombosis than embolism.

In all there are sixteen cases of aphasia reported, and of these three proved fatal. In certain cases the aphasia may be nervous in origin, neuropathic, or hysterical. In other cases it depends on albuminuria or uræmia. In nearly one-half the cases it coexists with right hemiplegia, and depends on thrombosis or embolism.

Having occurred in one pregnancy, it is liable to re-occur in the next. It usually appears about the first week after delivery.

A CASE OF PUERPERAL BONE SOFTENING CURED BY CASTRATION.

HARAJEWICZ (*Wiener medizinische Presse*, 1893, No. 27) reports an interesting case of successful removal of the ovaries for the cure of puerperal bone-softening. The patient had had ten children, and was in her eleventh pregnancy when she began to have severe pains in the loins, back, and hips. Her twelfth pregnancy was attended by more severe and continuous suffering; all the bones of the body except the face became acutely tender, and coughing caused intense pain. No visceral disease could be found; the urine was normal. The deformity of the pelvis and skeleton generally was now marked; there was no ovarian tenderness, and no uterine disease. The operation was performed under strict antiseptic precautions; both ovaries were removed and found to be normal. Complete and permanent relief from pain at once ensued, and after three months the patient left for her home. The bones were hardened, but crooked. Should this condition develop during pregnancy, it is probably best to empty the uterus; and, if it continue, castration may follow, or a Cæsarean section may be made at once, the ovaries being removed at the same time.

TWO EPIDEMICS OF AFEBRILE ICTERUS NEONATORUM WITH HÆMOGLOBINURIA (WINCKEL'S DISEASE).

WOLCZYNSKI (*Internationale klinische Rundschau*, 1893, No. 28) reports the results of his experiences in two epidemics of the above. Numerous post-mortem examinations revealed subpleural extravasations in the lung parenchyma, with thrombi in the vessels. The same condition was to be found in the liver and spleen, the former being in a state of fatty degeneration. In all these thrombotic masses, or in effused blood, the characteristic bacilli were to be found. The kidney was the seat of a subcapsular extravasation; pigment masses and rods were also present. Cultures made from the above in living animals were followed by death in from twelve to seventy-two

hours. In the urine were found Teichmann's crystals, red corpuscles, coloring matter, and bile pigment.

The writer believes the disease to be the result of the introduction into the child's system of the bacterium *coli communis*; the mode of introduction in this instance was probably by washing the children's mouths with water from foul sources. The bacterium was found in the water of the well used, and the substitution of sterilized bottled water was followed by a cessation of cases.

In 1892 six children were attacked with the disease, and all died.

In 1893 the same number had the disease, with a mortality of five. The other case recovered.

BACTERIA IN HUMAN MILK.

RINGEL (*Münchener medicinische Wochenschrift*, 1893, No. 27) contributes the report of a series of investigations upon the bacteria found in human milk. Various observers had reported having found the staphylococcus aureus and albus, as well as the streptococcus, both in healthy milk and in that from mothers suffering from puerperal fever. Escherich examined 25 women. Of these, 24 specimens were sterile and 1 contained bacilli. He again examined 13 with puerperal fever, and found staphylococcus in 12; 4 being of the white and yellow variety intermixed; 8 the white only, and 1 of an uncertain form. Cohn and Neumann experimented on 43 cases of milk from healthy women, and found 36 containing staphylococcus albus, 1 staphylococcus aureus and pyogenes, 3 staphylococcus pyogenes albus and streptococcus pyogenes; in 2 all the above forms were united.

The writer made a series of investigations, drawing and using milk from the deeper parts of the breast only, the experiments being made under the strictest antiseptic precautions. The milk was taken from 12 healthy and 13 unhealthy patients.

The results were as follows: 3 specimens were sterile; 17 specimens contained staphylococcus pyogenes albus; 2 specimens contained staphylococcus pyogenes aureus; 1 specimen contained staphylococcus pyogenes albus and aureus; 2 specimens contained staphylococcus pyogenes albus and streptococcus pyogenes.

An examination of the mouths of nursing infants revealed corresponding bacteria in their secretions.

TREATMENT OF ECLAMPSIA.

CHARPENTIER (*Archives de Tocologie*, 1893, p. 509), in a collection of 454 cases from various sources, gives his results as follows: Children dead before or during labor 164, or 36.12 per cent.; maternal mortality 110, or 24.88 per cent. His conclusions are as follows:

1. Every pregnant woman who is albuminuric is exposed to eclampsia. Consequently, we should examine the urine of all women during the period of gestation, and if the least trace of albumin be found, they should at once be placed on an absolute milk diet. Milk is, above all, the best preventive of eclampsia.

2. Whenever one deals with an eclamptic, if she be strong, vigorous, and