The value of this case depends on the points to which attention has been directed; the inversion was complete; the position of the placenta and its large size, coupled with the absence of evidence of artificial production, tend to support the theory of Hennig to which reference has been made. The patient made an excellent recovery and there was no tendency to secondary chronic inversion.

Complete inversion of the uterus is extremely rare. Braun had never seen a case in 150,000 births. In Ramsbotham's tables of 50,000 births the accident is not even mentioned, and it seems doubtful if in 1847 it had even been recognised. From 1745 to 1880 at the Rotunda in 191,000 deliveries it had only once been observed.

Westerham.

ON BLOOD COAGULABILITY IN THE PUERPERAL STATE.

BY R. HINGSTON FOX, M.D. BRUX., M.R.C.P. LOND.,
PHYSICIAN TO THE FRIENDS' PROVIDENT INSTITUTION.

THE object of this paper is to record observations made on the coagulation time of the blood in the puerperal condition. The observations were made by Wright's method, modified by the author, and a correction has been introduced for temperature.

Method.—Sir A. E. Wright's apparatus is described and figured in THE LANCET of Oct. 14th, 1905, p. 1096. I have after experience dispensed with the mercury, the segmentation of the tube, and its hair-fine tapering end, and the method used by me is as follows:—

The apparatus consists of a series of capillary tubes, eight centimetres in length, and each of such calibre that five cubic millimetres of mercury fill it to the length of five centimetres; each tube is inserted, the smaller end outwards, into a short larger tube and fixed in its place in an air-tight manner by sealing wax. Rubber nipples to fit the tubes, a beaker of water standing on white paper, a spirit lamp, an accurate thermometer, and a piece of string are provided. Four of the tubes are taken in succession and nearly filled with blood from a puncture near the end of the finger or thumb. Care should be taken that the blood flows freely, as squeezing the finger is apt to cause lymph from the tissues to be mixed with the blood, leading to an increased coagulation time. Nor must the finger be ligatured, as this much lessens the coagulation period. The blood lying in the tube should not quite reach either end of it. Each tube as soon as filled is laid on the table in order by the side of the thermometer and the exact time of drawing the blood from the finger noted on paper. The finger is then wiped and the puncture closed, until after an interval of 30 seconds or more the next tube is applied and the time noted. When all four tubes have been filled the observer decides at what interval he will examine the first tube, commencing well under the expected coagulation period. A rubber nipple is applied and the blood in the tube expressed into the water. If the mass is readily diffused the next tube may be left for one or two minutes longer than the first. The process is repeated until either a defined wormlike clot is expelled, which retains its form for some seconds, or until the contents have become so dense that they are with difficulty expressed. This occurs in some cases apart from the formation of a red wormlike clot, the mass being partially diffusible in the water. The reason of this variation is probably the admixture of coagulable lymph. At present I reckon the coagulation time to be the time which elapses before either a wormlike clot is formed or else the tube is with difficulty emptied, whichever happens first. Should the nipple fail to expel the contents the string should be wound tightly around the lower part of the nipple and then persistent efforts in compressing it will generally succeed in emptying the tube. If this cannot be done the tube must be discarded. I have not found any means of cleansing a tube when once choked.

Sir A. E. Wright accounts the coagulation time to be completed before the first appearance of a shred of fibrin formation. This appearance is seen much earlier and I think may sometimes be accidental, due to the presence of a minute quantity of blood less freshly drawn or of skin tissue or other extraneous matter. It seems to me better to wait until the entire mass of blood is coagulated. As by this means we obtain a much longer coagulation period than Sir A. E. Wright's the figures are not comparable with his.

Correction for temperature.—Five observations in immediate succession were made on a healthy man, aged 22 years, whose pulse was 72, regular; arterial pressure 100 degrees (Hill and Barnard). The blood was known to have an over-average coagulation time. The tubes when filled with blood were placed in small test tubes, weighted with mercury, and plunged into a water bath of a capacity of about two litres. The water was maintained at a given temperature—viz., 40°, 50°, 60°, 70°, and 80° F. respectively—in the five observations. It had previously been ascertained that the temperature of the air within the test tubes at several readings accorded closely, in each case, with the temperature of the water in the bath. The following results were obtained. Coagulation time at 40° F., 16 minutes; at 50°, 10 minutes 45 seconds; at 60°, 8 minutes 30 seconds; at 70°, 7 minutes 30 seconds; at 80°, 6 minutes 40 seconds. On plotting these out to scale the coagulation time is seen to form a curve, the ordinates of time increasing greatly as the temperature descends towards freezing point, to which line the curve doubtless forms an asymptote. A curve has been found which closely approximates to this between the temperatures 40° to 80° F., but I have not yet obtained a formula for

reducing observations to the standard temperature of 60°. This correction must therefore be made from the diagram graphically.

Observations on blood coagulability in the puerperal state.—By the kindness of Dr. W. J. Gow, physician to Queen Charlotte's Lying-in Hospital, I have made observations on the blood of eight parturient women during the first week after delivery in the wards of that hospital. All were primiparæ; unmarried, except the patients in Cases 2 and 8; the labours were normal, but in Cases 1, 3, 5, 7, and 8 they were tedious; hæmorrhage was normal; all suckled their infants, who appeared to be healthy. In Case 4 there was a trace of albuminuria and the temperature rose to 100·2° F. on the second day, the pulse-rate being 108; milk on the fifth day was rather deficient. Gonorrhœa was present in Case 1. The observations were taken between the hours of 2.30 and 6 P.M. The patients were, of course, recumbent in their beds. The results, corrected for temperature, are as follows, the period required for complete coagulation being represented in minutes:—

	Years of age.	Minutes.	Minutes.
Case 1	26	First day ... 4	Fourth day ... 6½
" 2	25	" " ... 5	" " ... 8½
" 3	22	" " ... 5½	Fifth " ... 6½
" 4	21	" " ... 6½	" " ... 7½
" 5	28	Third " ... 6½	—
" 6	20	" " ... 6½	Seventh " ... 6½
" 7	23	" " ... 5½	" " ... 7½
" 8	23	Seventh " ... 7	—

The observations may also be summarised as under:—

First day	Four observations	4, 5, 5½, 6½ minutes.
Third day	Three	6½, 6½, 5½
Fourth day	Two	6½, 8½
Fifth day	Two	6½, 7½
Seventh day	Three	7, 6½, 7½

Comparing these figures with many other observations taken by the same method I find that the coagulation time immediately after delivery is below the normal but not to the same extent as is commonly supposed. For I have often found a much lower coagulation time in adults, as low as three minutes or less. Five and a quarter minutes, the average time on the first day after confinement, is not therefore very low. Perhaps under the careful management of labour in the hospital the hæmorrhage is less, and so the blood coagulability is less altered on this account. It will be noticed that in every case in which two successive observations were taken the coagulation time had increased during the interval. The excretion of calcium salts in the milk tending to reduce the coagulability of the blood was probably one cause of this increase.

An observation of the coagulation time of the blood after delivery might give useful warning: of the risk of thrombosis or embolism if the coagulation time were low, or of post-partum hæmorrhage if it were high. Appropriate treatment—for example, with citric acid on the one hand or with calcium lactate and salines on the other—would rectify the condition.

I made a few observations on some of these women during the latter part of pregnancy and subjoin them for what they are worth. So far as they go, they indicate a tendency to a low coagulation time, in several cases lower than after delivery.¹ The blood during pregnancy is probably rich in nutrient matter required for the growth of the foetus. According to Winckel, it has a somewhat diminished alkalescence and exhibits a more ready (*mässige*) leucocytosis. Bar and Daunay state that the density of the blood is diminished at the end of pregnancy but rises again after partus. My observations of the coagulation point were made under various conditions as to time of day and relation to exercise and to meals, contrasting with the even conditions attaching to those made in the wards of the hospital.

Observations on the Same Cases during Pregnancy.

Case 1	11 days before partus	4½ minutes.
" 2	18 " "	2½ "
" 3	31 " "	2½ "
" 6	54 " "	7½ "
" 7	47 " "	4½ "

I have not found any reference to the coagulability of the blood in pregnant or puerperal states in recent works on obstetrics, British, French, or German.

Weymouth-street, W.

¹ Compare Dr. Blair Bell's reference to the presence of large quantities of calcium salts in the systems of pregnant women, Brit. Med. Jour., April 20th, 1907, p. 921.