of the cast, seem to be: (1) the increase of the intermalleolar measurement; (2) the prominence of the heel; (3) the approximation of the malleoli to the plane of the sole; (4) the absence of fracture; (5) that the injury was not compound; and (6) the extreme ease with which reduction was effected.

NOTE ON A CASE OF RENAL CALCULUS.

By H. E. BRUCE PORTER, M.R.C.S. ENG.,
L.R.C.P. LOND.

THIS note upon a case of suppuration of the right kidney caused by a calculus which first gave rise to symptoms 12 years ago and which for the whole time was mistaken for sciatica may be interesting in the light of recent correspondence in THE LANCET upon sciatica.

The patient, a woman, aged 35 years, sent for me early in 1901. When asked what she complained of she seemed to make light of her trouble, replying, "Oh, it is only a touch of my old friend sciatica." As the patient was to my mind rather young to have old friends of such a nature my next query was, "When did you first have sciatica?" I was informed 12 years ago and that it was so bad that she was in bed for some weeks, her then medical attendant devoting much thought to the case and using every means in his power to cut short the attack. As the treatment had no effect on the disease she was sent to the German baths and spent the best part of three years seeking relief without much success. My next query was, "Please tell me about your present attack." The patient replied that she had lunched out and drove home in a dogcart; she felt rather out of sorts and during the night had a shivering fit which returned off and on for some hours. The pain was of a dull character in the right side, there was frequent micturition, and her temperature was 102° F. The right kidney seemed the most probable seat of trouble and it was felt to be enlarged and tender; there was no tenderness over the sciatic region. I now returned to the history of the sciatica of 12 years or so ago. The onset of this was sudden while out walking and, in the patient's words, she walked home doubled up with pain. This did not sound like sciatica. Everyone knows how a patient with this trouble walks—certainly not doubled up. The pain ran down the inside and outside of the thigh and at times was in the sole of the foot. It was not affected by the position of the leg—in fact, as she put it, the pain made her squirm. The first attack lasted, as I have said, off and on for three years and during this time she believed that the urine was examined; she had never noticed blood in it. For the last nine years she had always had an uncomfortable feeling in her right side and occasionally attacks of "sciatica" (?). During these attacks she had noticed the urine to be thick but took no particular notice beyond observing the fact.

The patient was married about four years ago and was pregnant for the first time 12 months later. When three months pregnant the pain in the right side became worse; she also had so much swelling of the hands that she could not wear her rings; she was given to understand that this was due to her pregnancy and her urine was not examined. The pain in the right loin during the later months of pregnancy was worse but was all put down to the uterus. Since the birth of her child she had had to give up many of her pursuits and from being active became the reverse. At times she was unable to bear anything tight around the waist. The lungs were quite healthy and there were no signs of wasting—in fact, considering the history the patient was very well nourished. There was no history of phthisis. I ordered the whole of the urine passed during the next 24 hours to be saved in one vessel, the clear portion to be poured off, and the sediment to be sent to me. This contained one-sixth albumin; a few pus cells were to be recognised under the microscope but no blood or crystals except urates. I came to the conclusion that the case was either calculous kidney or tuberculous disease of the kidney, that the attack 12 years ago had been renal colic, and that during pregnancy the change had become more rapid in the breaking down of renal tissue. I advised complete rest and also advised the husband to have a second opinion, pointing out that the urine ought to be examined by a bacteriologist to exclude as far as possible the tuberculous cause.

Dr. Archibald E. Garrod made an examination. In the urine of the second 24 hours there was more pus (about three drachms). He agreed with my view of the case and took a specimen of urine to town to have it examined for tubercle. It was found free from tubercle and though the negative result did not exclude the possibility of tuberculous disease it was an additional point as there was no evidence of trouble in the chest, and the history of what we were both agreed was an attack of renal colic 12 years previously made us hope the case would prove to be one of calculous kidney. The temperature varied between 102° F. at night and 100.4° in the morning. On Feb. 25th the patient's child being put on the bed pressed against the damaged kidney, the result being that the swelling was much greater. It could now be seen as an oblong tumour in the right side of the abdomen. On the morning of the 26th, as the pus was not coming away freely enough, hot fomentations were applied. On the 27th the pus was coming away more freely. Sir Frederick Treves came to see the patient to decide when he would advise operation. He confirmed by his examination of the case and history the view taken by Dr. Garrod and myself—i.e., that the trouble began with an attack of renal colic which was then taken for sciatica. He decided that it was a case for removing (when the active stage had passed and the temperature had become normal) what was a serious menace to life. Pus continued to come away and on May 6th the patient's temperature touched the normal line in the morning, rising to 99.4° at night. As it was necessary to have the patient in a surgical home in town for the operation I allowed her up for one hour on the 8th. The temperature went up to 99.6° in the evening. On the 9th the temperature was again normal. The patient was up for three hours. The temperature on going to bed was 101° but came down after she had been in bed about an hour. On the 11th I took the patient to town and she stood the journey well. On the 13th Sir Frederick Treves operated. The organ was removed from the front; an incision was made along the right semilunar line for a considerable extent. The other kidney was examined and found to be quite healthy; the ovaries were also felt and found to be healthy; the right kidney was adherent at the upper end so the lower was delivered first. It is not necessary to describe the various steps of the operation, so I will content myself with saying that the wound was closed without the insertion of a drainage-tube and the patient was taken back to bed. The kidney was a mere shell of pus; there was not a trace of renal tissue left. It contained at least 15 ounces of pus. The calculus which had evidently started the trouble had become quite disintegrated. The patient had a good night and the urine passed was quite healthy. The patient made a good recovery except for the delay caused by the formation of an abscess in connexion with the stump of the ureter. About ten months later the patient became pregnant and this pregnancy was free from the pains, &c., of the former one. The child was born at term. I think the case of interest as showing that in a young woman the symptoms caused by a renal calculus had given rise to the idea that she was suffering from sciatica. The patient is now in perfect health and certainly looks and feels years younger.

Windsor.

THE TREATMENT OF SCIATICA.

By J. MICHELL CLARKE, M.A., M.D. CANTAB.,
F.R.C.P. LOND.,

PROFESSOR OF PATHOLOGY, UNIVERSITY COLLEGE, BRISTOL; PHYSICIAN
TO THE BRISTOL GENERAL HOSPITAL.

My reason for writing this short paper is that I believe that the Weir-Mitchell method of treatment of sciatica by immobilisation of the affected limb by means of the long splint is not so generally employed as it deserves to be. It is true that there are references to the method in some of the text-books but they are brief, and although Dr. I. Burney Yeo gives a longer and more satisfactory description in the last edition of his "Manual of Medical Treatment," it does not seem to be generally known that obstinate and protracted cases of sciatica can be successfully dealt with in this way. It is not my intention to enter into the discussion of what is the exact nature of sciatica but to confine myself to a brief statement of my experience of this particular