rapid and uninterrupted, there being no repetition of the abnormal pulse condition.

Here I think the dyspnoea was the active cause, but so long as the arterial tension remained fairly good no paradoxical pulse showed itself; however, on the occasion of the tension dropping and the pulse becoming soft, then the dyspnoea was able to produce a definite effect, the result being the pulsus paradoxus.

CASE 2.—A man, aged 40 years, was admitted into hospital on Feb. 8th, 1899, suffering from broncho-pneumonia complicated by pericarditis with effusion. He had been ill for four days previously to admission. His condition was critical and he had an irregular temperature ranging between 99° and 104° F. The pulse-rate was 130 and the respirations were from 40 to 60 per minute. Under treatment he improved considerably up to Feb. 21st. From that date he slowly began to lose ground, his most prominent symptoms being marked restlessness, a very troublesome cough, considerable dyspnoea, and inability to sleep in the recumbent posture; towards the end he became delirious. His temperature remained a little above normal, the dyspnoea increased, and he died suddenly on March 26th. The definite improvement in the general condition up to Feb. 21st was associated with the gradual clearing up in the lungs. Following this, however, the pericardial condition and the cardiac changes accompanying it became the main factors in the case and were the direct cause of death. On March 13th I made the following note: "The patient is still suffering greatly from dyspnoea, cannot lie down, and is occasionally attacked by great difficulty of breathing, the number of respirations increasing to 70 per minute. During these attacks he becomes almost frantic. They are not associated with marked cyanosis. The præcordial dulness is increased in area, extending two inches to the right and five and three-quarter inches to the left of the middle line, together making a transverse dulness of seven and three-quarter inches. There is dulness in the first left interspace extending one and three-quarter inches from the middle line. No friction rub is to be felt on placing the hand over the præcordium, and only an indefinite cardiac pulsation. The apex beat cannot be seen or felt, and the heart sounds are practically inaudible. There is occasionally a short systolic murmur in the fourth and fifth spaces. There are râles to be heard in the chest, more particularly over the lower lobes, and a pleural friction rub over the right middle and lower lobes. There are definite signs of fluid at the right base, the dulness reaching as high as the ninth spine. The liver is eight and a quarter inches in nipple line and is enlarged and tender. The spleen measures four inches in long diameter. The veins of the neck and arms are tense and engorged, but there is no definite cyanosis." The left pupil was dilated and larger than the right, a condition which I have noticed in several cases of acute pericarditis with effusion. His radial pulse was soft, frequent, and varied rhythmically with the respiration.

Tracings Nos. 4, 5, 6, and 8 in the diagram show well the "pulsus paradoxus." They were, with the exception of No. 6, taken with the patient breathing quietly. During the time that No. 6 was taken the patient was asked to breathe more deeply. The five tracings all show a condition of low arterial tension, Nos. 4, 5, and 6 being distinctly hyperdicrotic. It is noticeable that where the paradoxical condition is most marked, as in No. 6, there the pulse is most hyperdicrotic.

On March 15th 63 ounces of clear fluid were withdrawn from the right pleural cavity. This was followed by considerable improvement in breathing and the patient passed a comfortable night. Pulse tracing No. 7 was taken on the evening of March 15th following the thoracentesis and shows a definite improvement. There is certainly a distinct respiratory curve but otherwise the tracing might pass for that of a normal pulse of low tension.

The above alteration in the pulse was undoubtedly brought about by the removal of the fluid from the chest and although the fluid did not re-accumulate, so far as can be judged by physical signs, the paradoxical condition of the pulse returned as markedly as ever.

Tracing No. 8 was taken on March 23rd, three days before the man's death; he was breathing quietly, though rapidly. The above tracings are all from the right radial artery; others were taken from the left radial artery presenting the same condition. It was impossible to obtain any tracings from the anterior tibial artery on account of the œdema about the ankles and feet. Unfortunately no necropsy was permitted.

I would like here to express my indebtedness to Dr. Nathan Raw, as it is through his kindness that I am able to publish these cases.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

HEMIATROPHY OF THE TONGUE.

BY L. A. PARRY, M.D., B.S. LOND., F.R.C.S. ENG.,
LATE HONORARY ANÆSTHETIST, NORTH-WEST LONDON HOSPITAL.

THE following case is of interest not so much because of the condition of the tongue as on account of the unusual cause to which it was due.

The patient, a man, aged about 50 years, complained that one side of his tongue was growing smaller. He had no other trouble whatever, but he could not quite understand why this should be, and so he had sought medical advice. The condition present was simply atrophy of one side of the tongue. The only point in his personal history bearing on the case was that some years previously he had had a serious injury to his head, resulting in fracture of the base of the skull. Now, the causes of an unilateral atrophy of the tongue are disease of the hypoglossal nucleus or disease of the same nerve. Supranuclear disease is not accompanied by wasting, or, at any rate, only to the slightest extent. Facial hemiatrophy is sometimes accompanied by hemiatrophy of the tongue, but here this need not be discussed, for there was no hemiatrophy of the face. Nuclear disease is almost always bilateral and is generally accompanied by disease of adjacent nuclei, giving rise to labial paralysis. But every now and then in chronic degenerations, such as locomotor ataxia or progressive paralysis, of the insane, this affection is unilateral. Here, too, other nuclei are generally affected and other symptoms characteristic of the disease (tabes or progressive paralysis) are present. If the lesion is below the nucleus it may be (1) within the medulla, and then there is paralysis of the leg and arm of the other side; or (2) outside the medulla, then the nerve may be damaged within the skull by meningitis or new growths, or outside the skull by wounds, cellulitis, tumours, neuritis, or caries of the highest cervical vertebræ. There are no symptoms in this case of tumours, meningitis, caries, and so on, and I think this hemiatrophy is due to injury of the nerve in its passage through the anterior condyloid foramen at the time of the fracture of the base of the skull.

Hove, Sussex.

OCULAR PHENOMENA ASSOCIATED WITH CHEYNE-STOKES RESPIRATION.

BY G. YOUNG EALES, M.R.C.S. ENG., L.R.C.P. AND
L.M. EDIN.,

LATE ASSISTANT DEMONSTRATOR OF ANATOMY, ST. GEORGE'S
HOSPITAL, S.W.

THE following case is, I think, worthy of record inasmuch as the ocular phenomena observed, so far as I have been able to ascertain, have not hitherto been placed on record.

A man, aged 61 years, short, dark, and thick-set, came under my observation two and a half years ago, when I found him to be suffering from a mitral systolic murmur, considerable hypertrophy of the left ventricle, and some amount of albuminuria. Dilatation of the ventricle gradually ensued and this was followed by pulmonary and hepatic congestion, increase of the albuminuria, and, finally, general anasarca, so that it may be said that the patient was more or less waterlogged for the last two years of his life. Eight weeks before his death uræmic convulsions set in and continued off and on down to ten days before the end, when he became suddenly hemiplegic, after which the convulsions ceased. The patient was now paralysed on the left side and had in addition loss of control over the sphincters. Three days before his death