ATROPHIC PARALYSIS OF THE UPPER EXTREMITIES.—Vierorth (*Dtsch. Arch. f. klin. Med.*, Bd. xxxi, Heft. 5, 6) calls attention to four well-defined affections, in the course of which the above-named condition was observed. 1. *Peripheral traumatic paralysis*. In this, the motor disturbance is strictly confined to the distribution of the affected nerve, and more or less marked muscular atrophy follows. Disturbances of sensibility are rarely absent in decided lesions, but do not always involve the whole distribution of the affected nerve, being often limited to a part only. Electrical examination exhibits different results according to the stage and severity of the affection. 2. *Peripheral neuritis*. The motor disturbance generally indicates the involvement of a definite nerve-distribution; certain muscles, however, may remain wholly or relatively unaffected, yet usually the involvement of one or more distinct nerves may be determined. Functional impairment usually precedes atrophy, though not always; both may go hand in hand. Disturbances of sensibility are generally in the distribution of the affected nerve, or confined to a portion of the same; sensibility may be normal, however, while disease of the muscles persists. Electrical examination may give very variable results, the different muscles also varying between themselves in electrical reaction; complete or partial "degeneration reaction" is the rule, however, when the disease is at its height. 3. *Progressive muscular atrophy*. The muscular affection may be more or less diffuse in its distribution, or, in some cases, entirely irregular, while in others it may be confined to definite nerve-tracts. It begins usually in the small muscles of the hands, exhibiting a complete parallelism between the development of the atrophy and the paresis. Sensibility remains unaffected. Usually partial, and occasionally complete, degeneration reaction can, at times, with difficulty be demonstrated, and in the latter stages reaction may disappear entirely. Fibrillary contractions are very frequent. 4. *Chronic poliomyelitis anterior*. The motor disturbance is not limited to definite nerve-tracts. In the common form paresis precedes atrophy; these relations may be changed: atrophy and paresis appearing at the same time. Decided disturbances of sensibility are absent. Complete or partial degenerative reaction is found.—Litten *Deutsch Medizinal-Zeitung*, Dec. 20, 1883.

THOMSEN'S DISEASE.—Prof. Westphal exhibited two cases of this disease before the Berlin medicinische Gessellschaft, in Jan., 1883. He gave this name to that rarely observed and peculiar disturbance of motility, consisting of a sort of tonic spasm, because the first information of this affection was described by Dr. Thomsen, and because his whole family were affected with it in a typical manner. The symptoms are, that along with an otherwise normal condition, all the voluntary movements are executed imperfectly or with difficulty, and combined with tetanic muscular contractions, which persist for some time after the execution

of the voluntary movement. These phenomena appear : 1. After a period of inactivity of the affected group of muscles. When, for example, the patient desires to raise himself from a chair, for which a certain amount of force is required, the muscles of the leg exhibit powerful contraction. 2. After forced movements. The biceps contracts strongly when the forearm is suddenly flexed ; extension of this member is only possible after a certain interval. 3. In complicated movements the same rigidity occurs, interfering with their regular execution. 4. According to Thomsen, merely thinking of a muscular movement is sufficient to provoke the spasm. Westphal's cases, however, did not exhibit this symptom. Cold favors the production of the phenomena. The passive movements are pretty free, even though certain muscular groups sometimes offer a resistance to their execution. All muscles supplied by the spinal nerves may be affected by this disease. The muscles themselves show such an excessive development that these patients exhibit an athlete's figure. This hypertrophy is not accompanied by a corresponding increase in function. On the contrary, it barely enables the normal efforts to be executed with a superfluous addition of force, while certain attempts fail, being, in fact, frustrated in consequence of this peculiar disturbance of function. The fingers are unaffected. The muscles innervated by the cranial nerves, on the other hand, do not escape, for muscles of the eyes and mouth may be affected. For example, when the eyes are closed the patient finds difficulty in opening them, or in bringing the lips into a proper position for laughing ; particularly at a low temperature. Sensation is normal, with, perhaps, the exception of occasional formication. Faradic and mechanical excitability is remarkably exalted. The application of an electrode to a muscle, or the stroke of a percussion hammer, produces an extraordinarily powerful and long-continued contraction. The skin and tendon reflexes are normal. The etiology of the disease has not yet been positively determined ; nor has the microscopical examination of an excised portion of muscle thrown any particular light on the cause of the affection. Westphal considers the most reasonable explanation to be, that it depends on an increase in the muscular tone, allowing weak stimuli to produce excessive excitation. The hereditary tendency of the affection is very noticeable. In Thomsen's family few escaped. One of the patients exhibited by Westphal is a nephew of Thomsen. Treatment has been thus far without effect. Aside from its scientific interest, this disease has also a practical side, as those affected may frequently be suspected of simulation, on account of their inability to execute complicated movements while they exhibit a magnificent muscular development. This is particularly true respecting military service, as happened to one of Thomsen's sons.—*Deutsche med. Wochenschr.*, Feb. 7, 1883, p. 86.

W. R. BIRDSALL, M.D.