MIDDLESEX HOSPITAL.

OPERATION FOR FISTULA IN ANO; INTERNAL PILES AND FISSURE OF THE ANUS IN A PATIENT FOUR MONTHS ADVANCED IN PREGNANCY; SATISFACTORY RESULT.

(Under the care of Mr. GEORGE LAWSON.)

ALTHOUGH it is a rule to avoid surgical operations during the period of gestation, yet occasionally patients present themselves when either the rapidity of an abnormal growth or intense suffering calls for immediate surgical interference, and such instances we have occasionally recorded in our "Mirror." In the case we now publish it was the intolerable pain which made the woman seek surgical relief, and express herself willing to undergo any risk to obtain it. A fistula, internal hæmorrhoids, and a fissure, together caused so much suffering that she could get ease in no position, and every evacuation of the bowels was attended with positive agony. Mr. Lawson considered that a continuance of the irritation was more likely ultimately to produce miscarriage than an operation under the influence of chloroform, which would speedily give her relief.

E. C—, aged thirty-four, married, a dressmaker, applied to the hospital on November 8th, on account of the continued and great pain suffered at the lower portion of the rectum, which, at each evacuation of the bowels, became, she said, almost intolerable. She was four months advanced in pregnancy.

On admission there was great fulness of the lower portion of the bowel and of the neighbouring tissues. On the left side was a fistula, which ran up by the side of the bowel for two inches, but did not communicate with it. There was also a fissure of the anus and some internal piles.

Nov. 11th.—The patient having been put under chloroform, Mr. Lawson placed a double ligature on a large internal pile, and then laid open the fistula into the bowel, evacuating at the same time the contents of a large abscess with which it communicated.

12th.—With the aid of opium she has passed a good night, and the operation has produced no apparent irritation of the uterus.

The patient progressed most favourably for the first four days, when she began to complain of pain and heat at the lower portion of the bowel.

19th.—The pain and inconvenience she has suffered the last few days is accounted for by a small abscess in each ischio-rectal fossa, both of which were freely opened.

From this date all went on well; the wounds speedily granulated and healed, pregnancy proceeded uninterruptedly, and the patient was discharged on Dec. 21st convalescent.

WESTMINSTER HOSPITAL.

REMARKABLE CONSTRICTION OF THE ENTIRE TRACHEA AND LARGE BRONCHI, WITH THICKENING OF THEIR WALLS, ASSOCIATED WITH FIBRINOUS GROWTHS IN THE LARYNX, WHICH WERE SEEN WITH THE LARYNGOSCOPE; TRACHEOTOMY; INABILITY TO GET IN A TUBE; FATAL RESULT.

(Under the care of Dr. GIBB and Mr. HOLTHOUSE.)

WE have never lost an opportunity of placing upon record in our "Hospital Mirror" any case that presented interesting and rare peculiarities, or that was likely to afford information or throw light upon doubtful points of pathology. Diseases of the larynx have been freely illustrated, more particularly to show the value of the laryngoscope in diagnosis. Lesions of the trachea, on the other hand, are comparatively seldom met with; and in the description of the difficulties of the operation of tracheotomy by many of our first authorities, the circumstance of constriction of the tube appears to have been wholly overlooked. Can it be that such a peculiarity has escaped observation, or do none of our numerous and pathologically rich museums possess such specimens? In the last volume of "Guy's Hospital Reports" are plates representing constriction of the trachea at one point from syphilitic disease, and contraction of a bronchial tube from a syphilitic ulcer, also at one point. These with others illustrate a valuable paper on the Syphilitic

Affections of Internal Organs by that indefatigable observer, Dr. Wilks. Although familiar with most of the specimens of laryngeal disease in the London museums, we do not remember having seen any specimen wherein the trachea was constricted from end to end, as in the following case, with extreme hypertrophy of its walls, at the expense of the natural calibre of the tube. We have in mind some four or five instances of isolated points of constriction, such as Dr. Wilks has described, but not of a general nature. Such a circumstance as that about to be described is sufficient to attract attention; but the interest of the case is increased by other peculiarities, which render it at present unique in the annals of medicine. The patient was admitted with a noisy respiration, depending upon smooth, shiny growths of the larynx, diagnosed during life with the laryngoscope, and which were supposed to give rise to attacks of dyspnoea. In one of these it was necessary to open the trachea, and, to the dismay of the surgeon, a tube could not be inserted. Here was an unexpected and unlooked for complication, sufficient to tax the resources of the most skilled. As not the smallest tube could be introduced, nor a catheter borne from the irritation it produced, the patient was left to breathe through the opening made, and succumbed the same night.

The particulars of the autopsy explain the cause of the difficulty, which was surmised when the tube could not be passed. The remarkable complications present in this case—laryngeal growths, constriction of the entire trachea and both larger bronchi, with great thickening of their walls—will long render it one of the most remarkable that has ever been placed upon record.

The subjoined notes were furnished by Mr. W. Gandy, house-physician to the hospital:—

C. R. M—, aged twenty, was admitted into Burdett ward on the 22nd December, 1863, under the care of Dr. Gibb. He had been previously a patient at the Hospital for Diseases of the Chest, Victoria Park, under the care of Dr. Thorowgood, who, recognising the presence of serious laryngeal mischief, sent him to Dr. Gibb. A difficulty of breathing and a stridulous noise had existed for twelve months, with cough and expectoration. He had lost his voice several times for weeks together, which he attributed to colds. On admission, there was severe dyspnoea, with a stridor or roughness on inspiration, some pain and severe constriction about the larynx, and a feeling of oppression at the upper third of the sternum, where he frequently placed his hand. His aspect was pale and wan; the features were drawn up with an anxious and careworn expression; he looked not more than sixteen, although twenty years old; was much emaciated; and his hands were long and thin, with clubbed fingers. He had a hard cough, with expectoration of a thick viscid mucus tinged with blood; the breathing was laboured, causing a peculiar croaking sound with each inspiration; at every paroxysm of coughing he had much pain in the lower part of the trachea, and great difficulty in expelling each pellet of mucus. There was harsh breathing in the apices of both lungs, and the breath sounds generally were remarkably feeble, as if the free entrance of air was somewhat obstructed. The laryngoscope revealed a partially pendent and lopsided epiglottis, with the presence of growths on the right side of the larynx above the true vocal cord. When the larynx was expanded, both of the true cords could be seen, and the voice, although feeble and somewhat hoarse, was quite audible. During forcible expiration these growths were prominent, and appeared to occupy the position of the right false vocal cord, extending forwards to the root of the epiglottis. They were seen by Mr. Gandy (the house-physician), Mr. Holthouse, Mr. Firth, and several of the pupils on various occasions; and the annexed woodcut gives the appearance they presented.

a, The growths occupying the position of the right false vocal cord. The left true vocal cord is seen on the opposite side. b, The lopsided epiglottis. c c, The arytenoid cartilages. d, The back of the tongue.

He was also examined at the hospital two days before his admission. No growths were seen below the cords, and the symptoms were believed to depend chiefly upon those present in the larynx, but they did not explain the feebleness of the

breath-sounds. The dyspnoea was so great at times as to oblige him always to remain in the prone position. He had no dysphagia. There was no history of syphilis, although the disease was suspected from ulceration and purulent secretion of the left nostril. He was ordered a mild pectoral mixture, and iodide of potassium thrice a day; a solution of tannin to the larynx; and nourishing diet.

He improved a little, but the dyspnoea was still urgent and the cough became distressing. New-year's day Dr. Gibb had appointed for removing the growths, but about two o'clock in the morning he was suddenly seized with extreme dyspnoea; so urgent, indeed, that Mr. Holthouse was sent for, and tracheotomy was performed at four o'clock; but all his efforts to get in a tube failed. A gum-elastic tube some inches long, and the size of a No. 10 catheter, could be introduced; but it caused the patient so much irritation that it was withdrawn. A portion of the front wall of the trachea was removed, and the patient was left. At the usual hour of the visit, fresh attempts failed to get in anything, and it was clear that the trachea was much constricted. Dr. Gibb observed that the air scarcely entered, and the breathing was quite inaudible over the bifurcation of the trachea and larger bronchi. The patient died at a quarter past nine in the evening, chiefly from asphyxia.

Autopsy, seventeen hours after death.—The lungs were engorged with blood, with a few scattered miliary tubercles here and there quite recently deposited; the liver had one or two small white nodules on the surface; the genital organs appeared natural, as well as the other viscera. The tongue, larynx, trachea, and bronchi were removed for careful examination. The oesophagus was healthy. The epiglottis, from its root to half way upwards on either side, was occupied by a number of flattish, fibrinous bodies, which partook of the character of warts; two or three, the size of small peas, smooth and round, were present on the right side of the larynx, involving the false vocal cord, and were those seen during life. The ventricles of the larynx were unobliterated, and the larynx was unobstructed below the vocal cords; the mucous folds everywhere, but especially the aryteno-epiglottic, were very loose. The trachea was greatly thickened, and its tubal diameter much contracted; the anterior walls about its middle were half an

inch thick, thinning upwards, but less so downwards. This thickening involved the right bronchus, and slightly the left. About half an inch from its commencement the contraction began, and below the wound its diameter was a quarter of an inch, and this continued nearly all the way to the bifurcation, where even the walls were two lines thick. The diameter of the left bronchus at its commencement was about two lines, and its lining membrane was intensely inflamed, with several ulcerated patches. The rings of the trachea could not be distinguished from within, for the whole of its interior was irregular and uneven from fibrinous deposit.

Medical Societies.

ROYAL MEDICAL & CHIRURGICAL SOCIETY.

TUESDAY, JAN. 12TH, 1864.

MR. PARTRIDGE, PRESIDENT.

A CASE IN WHICH THE LEFT OVARY WAS FOUND IN THE SAC OF AN OBLIQUE INGUINAL HERNIA, OCCURRING IN A YOUNG WOMAN; WITH REMARKS.

BY HOLMES COOTE, ESQ., F.R.C.S.,
SURGEON TO, AND LECTURER ON SURGERY AT, ST. BARTHOLOMEW'S HOSPITAL.

A YOUNG woman was brought into St. Bartholomew's Hospital with a swelling in the left groin, and suffering from the symptoms of strangulated hernia. In the course of a few hours the usual operation was performed, when the ovary and the Fallopian tube were found in the sac. A similar malposition of parts was subsequently noticed on the opposite side of the body. The left ovary was removed, some thickened omentum cut away, and the patient was put to bed; but the sickness and constipation continued, and she died four days after the operation. The cause of the sickness &c. was displacement of the stomach and transverse arch of the colon.

Mr. Coote raises two questions:—

1. Was the displacement of the ovaries congenital, or the consequence of the hernia? He inclines to the former opinion.
2. The woman stated that she had always menstruated regularly. Now, on the examination of the body, it was found that both ovaries were *well developed*, and that the formation of the Graafian vesicles was going on naturally; but the Fallopian tubes were quite impervious, the uterus was completely absent, and the vagina was a short canal—an inch and a half in length, and terminating in a thin membrane. She said that she had been menstruating the week before her admission; and some of the female attendants at the hospital noticed the usual marks, though faint, upon her dress. Are we to admit the possibility of menstruation under this abnormal condition of parts?

Mr. PARTRIDGE said it was a very interesting question to decide whether menstruation occurred, as was stated, under such circumstances as obtained in Mr. Coote's case, or whether it was merely a vicarious action. There was another question, of great importance in a moral point of view, which presented itself to surgeons in such cases. Were they justified in emasculating, as it were, a woman in whom the ovaries were thus involved? A case had lately come under his care in which a difficulty of this kind existed. The patient was a male child, with the parts of generation so imperfectly developed that it was mistaken for a female, and christened and educated as such. It was discussed whether the testicles should be removed. The surgeon in advance thought that they should, as their removal would be advantageous to the child in assisting it to keep up its assumed sex. Mr. Partridge decided, however, that the operation was not justifiable, and it was not resorted to. Mr. Partridge then referred to two cases in which the uterus was absent: the one was an unmarried, the other a married woman. In each the vagina was short, but the clitoris, ovaries, and breasts were fully developed. In neither of these cases had there been any menstruation.

Mr. CÆSAR HAWKINS observed that the paper before the Society opened two interesting questions—one in relation to the operation of hernia, and one of physiological importance. Now, in hernia scarcely two operations were found to be precisely similar. He had met with two cases in which the ovary was found in the hernial sac. In one of these the patient was an elderly woman, and died of peritonitis. The Fallopian tube and ovary had been for many years in the hernial sac, and by their position had produced an elongation of the greater portion

The trachea laid open from behind. *a*, The opening of the operation. *b b*, The walls greatly thickened, and drawn of the natural size.

Horizontal sections of the trachea in different parts of its course.