VIII. Herniotomy in Cases of Suspected Gangrene of the Intestine. By Dr. HELFERICH (Greifswald). It has been the custom, hitherto, in herniotomies to leave the gut suspected to be gangrenous remain in a suitable dressing outside the wound. This was accompanied by disadvantages; first, the hernial opening in the abdomen had to be enlarged to enable peristalsis to continue unobstructed; this incision amounted to a laparotomy. Thus the subsequent appearance of a large ventral hernia was favored by the operation. Again, if the much feared gangrene occurred the patient would have an artificial anus, necessitating subsequent operative interference. Helferich proposes to establish an intestinal anastomosis between that part of the intestine above the gangrenous gut and the part below it. When the anastomosis is completed, the intestine which has been drawn outside of the hernial ring for operation, is replaced. The part threatened with gangrene is allowed to remain outside the hernial opening. If gangrene occurs, no harm is done. The peristalsis of the intestine is maintained from the first moment. The author says the operation is simple, not at all dangerous, and may be performed in exactly similar manner to the operation of gastro-enterostomy. It may be performed with the aid of Senn's rings, but in the latter case the author fears that the gut, which is already in a reduced condition, may be further endangered by the pressure of rings. The anastomosis is established opposite the mesenteric attachment to the extent of about 4 cm. Author advocates sutures. The suspected gut is resected if it becomes gangrenous, without danger of peritonitis, and the intestinal wound readily closed, and with the established anastomosis peristalsis is completed. Should the intestine retain its vitality, then no harm is done by the operation of anastomosis; the adhesions to the abdominal peritoneum are loosened around the hernial opening, and the gut replaced into the abdominal cavity. Author has performed this operation in two cases. In one case success followed; in the other the patient died in collapse, and autopsy proved satisfactorily that the intestinal anastomosis was complete. The above operation is recommended for cases of threatened gangrene. It is questionable whether it will be of use in actual gangrene of the gut.—*Beilage zum Centr. f. Chir.*, No. 25, 1890.

HENRY KOPLIK (New York).