confined. The urine was generally of low specific gravity, rather copious, and contained less urea than usual; occasionally a trace of albumen was found. She was moderately intelligent, but there were signs of general weakness in the nervous system. Speech was slow, and somewhat laboured. She said she occasionally "loses herself" for a few seconds. The patient was very deaf; she could hear a loud voice, but not a watch pressed to her ear. The reflexes were normal. There was some general loss of cutaneous sensibility, and the floor felt soft as she walked. She could stand with her eyes shut. The hands felt cold, numbed, and painful. There were no marked signs in the heart and lungs. No thyroid could be felt. She was treated at first with iodide of potassium, under which she seemed to improve, and said she felt steadier. She was then given nitrate of pilocarpine in doses of one-eighth of a grain, which caused some slight moisture on the skin, and appeared to increase her comfort considerably, but of late she has somewhat relapsed, and I do not think she is much better than when I first saw her. She complains principally of severe occipital headache.

Leeds.

A CASE OF EPILEPTIC AUTOMATISM.

By SOLOMON C. SMITH, M.D.,
SURGEON TO THE HALIFAX INFIRMARY.

THE following is an interesting example of epileptic automatism:—

A young man has for four years been subject to attacks of petit mal. He does not fall, but seems to "fix" his eyes, and sometimes walks forwards, or sideways, at others sits down, and generally ends by going to sleep. There are neither convulsions nor frothing at the mouth, nor tongue biting. The only warning is that he suddenly begins to wonder whether he is going into an attack, and then he loses consciousness. He does not think he ever dropped anything out of his hand, but he has spilt tea out of a cup. In some of the attacks he will stop what he is doing and seem to "study" for a time, but in others he will go on doing what he was engaged in at the time.

A. On one occasion, while alone in the house, he had been playing the oboe, and was just turning over some music when he lost himself, and knew nothing more for an hour and a quarter (he happened to know the time); when consciousness returned he was lying on the sofa still alone in the house. He, however, must have, during the state of unconsciousness, put the oboe properly away in its case and taken out the reed, for it was uninjured in its place when he came to himself. Now, this taking the reed out of an oboe is an operation requiring very delicate manipulation.

B. Another day he was with several musical friends. I think he had been accompanying some of them on the piano. He was standing at the table turning over some pieces of music when they saw him go into an attack. While it was obvious to them that he was in a fit, for they had seen him before, he took up a piece of music, went to the piano, and played it; he was quite unconscious afterwards of having done this.

C. In his father's mill there is a machine for winding warps, which is not under the special care of anyone, for, when started, it works for some time by itself; but occasionally a warp breaks, and when this takes place anyone who happens to see the broken end of the warp hanging down stops the machine, "piecens"—i.e., joins—the ends, and starts the machine again. Now, one day his father and others saw him go into an attack ("fix" his eyes and walk forward); and while he was in it he passed this machine, in which a broken warp was at that moment hanging down. He stopped the machine—which is a peculiarly complex action, for it must not be stopped suddenly, but has to be let down or "slowed" by degrees—he tied the broken ends, and then started the machine again. All this he did while obviously quite unconscious, and then he passed into the usual drowsy ending of the attack.

These are good illustrations of the complexity of those automatic actions which often occur during epileptic unconsciousness. It is important, however, to recognise that they did not constitute the fit, but took place in spite of it; it was in no sense purposive epilepsy; the patient's actions

were purely automatic, occurring in response to stimuli reflexed through centres lower than those involved in the discharge. If this be limited to parts essential to higher consciousness only, and lower areas are left untouched, these, being freed from control, will, according to Dr. Hughlings Jackson, act automatically, sometimes with even hyperphysiological activity, in response to any stimuli which fall in their way, which may, as was possibly the case in A, be the remains of intentions already cerebrally formulated, or may be sense impressions received while in the unconscious condition, as was certainly the case in C. The apparent rationality of the actions is confirmatory of the opinion expressed in the recent Croonian lectures—that in the "dissolution" of epileptic discharge "the lower level of evolution remaining—the then highest level—is of some part of the highest centres," that the middle and lowest centres are of course in activity, "but are put in activity by what remains of the highest." A illustrates the great delicacy of touch and manipulation which may be exercised; B, that such an art as music may be practised, an art requiring the use of memory in a high degree; and C, that a sensory impression received during the condition of automatism may set up a fresh and most elaborate series of movements, which are performed with accuracy and perfect appropriateness. These have been spoken of as post-epileptic, the suggestion being that the automatism occurs after the paroxysm, but as somewhat similar phenomena (i.e., non-convulsive, apparently purposive acts) sometimes occur very early in the attack, even before a convulsion, we must believe that all that is necessary for their production is that discharge should occur in those nervous structures whose interlacements form the physical substratum of consciousness, while the reflex circles (or more truly mazes) required for the particular automatic actions of the case are intact, whether as the result of non-implication or of prior recovery is a matter of indifference, as it also is whether the discharge in the upper layers still goes on or not. There is this great difference, however, between the actions at the two extremes of the attack; in those which are post-epileptic external suggestions obviously play a part, so that the resulting phenomena vary in the different paroxysms of the same individual, but the early movements which take place at the commencement are apt to be repeated each time in nearly the same form, and so to seem really to constitute part of the fit itself. When, for example, a girl rushes into the street tearing at her dress and shrieking that she has rats inside her, and then falls into stupor, or when a man while conversing suddenly begins to undress, and this ends in a convulsion, and when in each case the same sequence of events recurs every time, the repetition of the same phenomena negatives their dependence on external and therefore varying stimuli, and suggests, wrongly as I think, that the action is part of the fit, the direct result of the discharging lesion. Notwithstanding the exactness with which it has been found possible to map out the convolutions into motor areas, it must not be concluded that either they, or the deeper parts with which they electrically correspond, are centres capable of initiating the movements which result from their stimulation, or that there exist centres at all for more elaborate acts; all we can say is that where cerebral activity results in certain movements, the impulses largely pass through certain corresponding areas. But these impulses come from no one part of the brain; they are partly sensory, partly reflex, partly volitional, and therefore imply activity of widespread regions, so that although irritation of a motor centre may produce a localised convulsion, it is impossible to conceive that a discharging lesion of any part of the brain could lead to the complex cerebration necessary for the production of the elaborate actions of epileptic automatism. These are, as Dr. H. Jackson has shown, the result of the activity of the remainder, which may have just been "set" to carry out some intention already cerebrally formulated at the moment of suspension of consciousness, or may be directed in certain grooves by sense impressions, either from without or from the viscera, or may be influenced by various subjective sensations produced by the discharge itself. In epileptic attacks consciousness departs in various ways, and its total abolition is often preceded by steps of less and less perfect relation to the environment—in other words, by a series of subjective sensations. When these develop slowly and are remembered they form the aura, but it is impossible to say how far into the fit they may extend, and they would be as likely to give rise to automatic action as would external sense impressions. Dr. Hughlings Jackson had a boy under his care who was

subject to fits beginning by a subjective sense of smell: "It was an interesting and important fact that he would hold his nose when the fit began. No doubt as his consciousness became obscured, 'he' believed the smell was of something actually outside. It is perhaps too strong an expression to say *he* believed, for where is the 'ego' when a man is unconscious, or partly so? We may say the smell developed an epileptic dream. I had a woman under my care who at the beginning of an attack had 'subjective' sensations of smell, who said, 'What a dreadful stink there is in the place!' She believed the smell was of something in the room."¹

In these cases the actions were obviously the result of the purely subjective sensation of smell, and in the boy's case went on into the unconsciousness; and it seems extremely probable that, when, as the first act of a fit, some one thing is always done, such as in the case of the man who takes his clothes off, or the girl who rushes out in terror, this is an automatic action started by some subjective sensation arising during the abrogation of consciousness; even the scream which so often heralds the attack may be caused by some great horror felt but not remembered. The theory, again so ably enforced by Dr. Jackson in the recent Croonian lectures, that the epilepsy is responsible for the convulsions and unconsciousness, but that the unfettered activity of the remainder of the brain is the cause of the post-epileptic phenomena, is of the greatest service in comprehending the disease. It perhaps simplifies things still further to look at all purposive acts, at whichever end of the attack, as due to the same cause.

Halifax.

NOTE ON THE TREATMENT OF CORYZA.

By G. E. DOBSON, M.A., M.B., F.R.S.,
SURGEON-MAJOR, ARMY MEDICAL DEPARTMENT.

THIS very troublesome complaint has scarcely received the attention it deserves, if we take into consideration the great number of sufferers and the serious laryngeal and pulmonary diseases to which it is too often a prelude. Amongst the host of remedies proposed for its abortive treatment, most of which are either of doubtful value or else difficult to procure or apply, or even dangerous to use, not one can be named of which it may be said that it is at every one's command, easy of application, unattended with danger, and really effective. No excuse is, therefore, needed in introducing to the notice of the profession the following simple yet thoroughly effective mode of treatment, which in my hands has never disappointed expectation. About a drachm of camphor, coarsely powdered, or shredded with a knife, is placed in an ordinary shaving jug, which is then half filled with boiling water. The patient, having made a paper cone (out of a sheet of brown paper or an old newspaper) large enough to surround his face by its wide extremity and the mouth of the jug by its narrow end, proceeds to respire freely, at each inhalation drawing the steam into his nostrils, and at each exhalation forcing it up against the outer surface of his nose and adjoining parts of the face. A twofold action is produced: the camphorated steam acts internally in a specific manner upon the whole extent of the mucous surfaces, and, externally, produces profuse diaphoresis of the skin covering the nose and sides of the face, there acting as a derivative from the inflamed Schneiderian membrane. The jug should be surrounded by a woollen cloth in order to prevent the water cooling, or, better, if a tin shaving-can be used, a small spirit lamp or heated iron may be placed beneath it, so as to maintain the heat of the water and the vaporisation of the camphor. The patient should continue his respirations (keeping the margins of the base of the paper cone closely applied round his face) from ten to twenty minutes, and this should be repeated three or four times in as many hours, till entire freedom from pain is experienced. Great relief is always felt even after the first application, and three or four usually effect a cure. Camphor, or some of its preparations, have, as is well known, been long in use in the treatment of colds, but the above-described method of employing it in conjunction with the vapour of water both as an internal and external application at the same time, has not, so far as I know, been previously brought to the notice of the profession, or, if

brought, has not been recognised in any general or special medical work. The mode of application is, however, all important; but as this is neither troublesome nor otherwise unpleasant to the patient, nor are the materials difficult to procure, camphor being everywhere a household drug, I believe that those who may give this treatment a trial will find it not only a simple, but also a most effective, remedy against coryza.

ON ACNE IN FIBRE-DRESSERS.

By CHARLES ATKIN, L.R.C.P. LOND., F.R.C.S.,
HOUSE-SURGEON TO THE SHEFFIELD INFIRMARY.

A DERMATITIS taking an acneform character, arising from an external irritant, is certainly uncommon. "Eczema," so called, in its various stages, is met with in many trades and occupations in all its varieties, from the "weeping vesicle" to the "leathery scaling skin"; but rashes of a tubercular form arising from external irritation of the hair-follicles have hitherto, as far as I know, been limited to those who work with tar, paraffin, benzoin, &c., or those cases where tar has been energetically used to cure some other form of skin disease, as psoriasis or chronic eczema. During last summer my attention was directed to a form of rash on the forearms and thighs of a certain class of workmen different from anything else I had ever seen in those situations. The first case met with was in a healthy-looking, intelligent man, aged twenty-four, who described himself as a fibre-dresser, at which special department of brush-making he had been employed for the last two years. His work consisted in dipping a fibre called "kitool" and cocoa-nut fibre into some special preparation of various oils, soaking it, and then tearing and pulling it out. Since he has been employed at this work he has suffered continually from a rash, which itches and is sometimes painful. His forearms were more or less covered with raised tubercles, discrete, and presenting all the stages from the "black point" of the closed sebaceous gland to the suppurating, tense, and painful nodule. There were no pustules on the backs of the hands, but the "black points" were universal, and could be extruded by pressure or picked out by a needle-point. Wherever pustules had been were well-marked scars. The hairs on the backs of the forearm were scanty, and could easily be pulled out with forceps, without any pain whatever. The same condition existed on the front of the thighs and shins, but more especially on the knees. There was no appearance of acne on the face, and only very scantily on the shoulders. Under the microscope the hairs treated with solution of potash were seen to be very swollen at their roots, with much adherent epithelium, evidently from proliferation of the lining of the hair-follicle. The man being otherwise healthy, with no appearance of struma, syphilis, congenital or acquired, or scorbutus, the rash was evidently connected with his work. On being questioned, he informed me that he continually got his trousers soaked in front in dipping the kitool, and that he worked with his coat off and his sleeves rolled up. The next question was, What was the cause of the rash—the fibre or the oil? The former, which is a kind of coarse dried grass, produces no local ill effects on those who simply sort and pick it, causing only some throat irritation from the quantity of dust arising from it; therefore the cause evidently lay in the liquid used. On procuring a sample of this I detected a smell of paraffin in it, and learnt on inquiry that it was not unusual to have this special oil mixture adulterated with this material. On visiting the place where the man worked, I found that all the men engaged in a similar occupation had acne more or less on their forearms, and that those who had been employed for some length of time were nearly absolutely hairless on those parts. Hebra says that when a rash is due to paraffin &c., it does not usually come from direct contact of the irritant with the skin, but by the person living in an atmosphere impregnated with its vapours; such, however, can hardly be the case here, for the rash is strictly localised to those parts which come in contact with, or are saturated by, the fluid, and also by the fact that it does not occur in those who only sort the fibre, though they are often at work in the same room. Again, it differs from acne caused by tar in that, Hebra says, the rash disappears on discontinuance of the irritation, leaving neither maculæ nor scars. Here the rash, though was told it very much

¹ West Riding Asylum Report, vol. v.