port wine naevus at a meeting of the Dermatological Society of London, in which treatment with radium bromide had been productive of decidedly encouraging results. The patient was a woman, aged twenty-six years, with a naevus covering the left cheek and the side of the nose. The two specimens of radium employed weighed 10 mgm. each, and were of the highest known radioactivity. Thirty-nine exposures, varying in duration from a half to one hour, were made at intervals of a week. At the time the patient was exhibited the naevus had entirely disappeared with the exception of a few small areas which had not yet been treated.

[The members of the Society before whom the case was presented expressed the opinion that the result of treatment was "distinctly encouraging." A photograph accompanies the report of the case, and if this picture shows the naevus at the close of the treatment, when the reporter states the mark had "entirely disappeared with the exception of a few small untreated areas," the result does not seem to the observer of the picture to be entirely satisfactory, as the photograph shows either part of the naevus remaining or scarring. If it be scarring, any treatment that will cause scarring in the treatment of naevi is to be deprecated.]

Thigenol.—SAALFELD (*Jour. mal. cut. et de syph.*, November, 1904) experimented with this new substance, put forward as a succedaneum to ichthyol. It is a composition of iodine and ichthyol, in which occurs 10 per cent. of sulphur in an organic state of combination. Thigenol possesses a dark-brown color and a syrupy consistence; is inodorous, almost insipid; soluble in water, alcohol, and glycerin. Applied to the skin it dries rapidly, and it does not stain the linen. It is a vaso-constrictor, is useful in inflammations, is drying, and favors resorption. It calms itching, as shown in 124 observations in eczema. In seborrhoeic inflammations in solution it was found especially useful. It is a mild remedy, and in the author's experience was of no value in psoriasis. In phlebitis, bandaging with thigenol, glycerin, and water, equal parts, gave very good results.

Treatment of Impetigo.—HONCAMP, of Berlin (*Monatshefte f. prak. Derm.*, Band xxxix., No. 12), gives a clinical note on the value of bismuth chinolin rhodionate (Oudim Edinger), which is applied as a powder after removal of crusts, with a wooden spatula, as metal instruments are to be avoided on account of ready oxidation. At the moment of application a slight burning is experienced, which soon passes away. This can be avoided by mixing the drug with equal parts of starch. Following the application, yellow, loose crusts form which become brownish. A bandage or covering is not required. The impetigo lesions heal under the crusts in three or four days.

Clinical Results Concerning Atoxyl.—BIRINGER (*Jour. mal. cut. et de Syph.*, November, 1904, p. 843) experimented with this new drug at the clinic for cutaneous diseases at Bonn. The drug is a white, crystalline, inodorous, almost insipid powder, soluble in 20 per cent. of hot water. The solution takes on a yellowish tint on cooling, and bases to take up about 2 per cent. of atoxyl dissolved. The toxicity is about forty times less strong than that of the other inorganic arsenical preparations. Forty-two patients received subcutaneous injections of 20 per cent.