Tabelle V.

Nr.	Rectascension			Declination		
	SH.	Reg.	SH. — Reg.	SH.	Mikr.	SH. — Mikr.
9	—30°861	—30°844	—0°017	—3' 53"57	—3' 53"59	+0"02
5	—26.810	—26.822	+0.012	+2 37.68	+2 37.99	—0.31
6	—22.115	—22.105	—0.010	+2 0.98	+2 0.99	—0.01
3	—20.739	—20.741	+0.002	+3 24.99	+3 24.95	+0.04
7	—20.241	—20.216	—0.025	+1 25.78	+1 25.79	—0.00
4	—20.188	—20.202	+0.014	+2 34.99	+2 34.95	+0.04
2	—16.020	—16.067	+0.046	+3 27.47	+3 27.58	—0.11
15	—9.360	—9.369	+0.008	—4 44.27	—4 44.41	+0.14
12	—7.515	—7.519	+0.004	—8 6.29	—8 6.50	+0.21
14	—1.489	—1.522	+0.033	—5 48.70	—5 48.76	+0.06
16	—1.338	—1.340	+0.002	—3 39.85	—3 39.83	—0.02
1	+0.008	+0.018	—0.010	+0 0.01	0 0.00	+0.01
13	+0.625	+0.614	+0.011	—8 7.30	—8 7.53	+0.23
17	+6.172	+6.165	+0.007	—4 2.70	—4 2.64	—0.06
18	+7.772	+7.751	+0.021	—4 25.16	—4 25.07	—0.09
19	+9.350	+9.330	+0.020	—3 11.68	—3 11.56	—0.12
25	+11.099	+11.076	+0.023	+3 8.71	+3 8.74	—0.03
23	+19.185	+19.187	—0.002	—0 32.71	—0 32.77	+0.06
26	+20.662	+20.697	—0.035	+2 0.01	+1 59.89	+0.12
27	+26.376	+26.337	+0.039	+0 31.30	+0 31.24	+0.06
21	+31.003	+31.051	—0.047	—4 58.12	—4 57.87	—0.25
32	+35.216	+35.223	—0.007	+6 11.33	+6 11.38	—0.04
34	+37.293	+37.311	—0.018	+3 42.43	+3 42.52	—0.09
33	+41.864	+41.851	+0.013	+3 48.64	+3 48.73	—0.09
31	+44.765	+44.796	—0.031	+9 45.89	+9 45.58	+0.31
29	+45.096	+45.098	—0.002	+2 45.04	+2 45.19	—0.15
22	+53.500	+53.521	—0.021	—1 50.92	—1 50.77	—0.15
30	+60.872	+60.889	—0.017	+6 47.03	+6 46.91	+0.12

Wegen der Nullcorrectionen ist jetzt nicht mehr der Stern Nr. 1, sondern der Punkt $\alpha = 18^h 21^m 47^s.07$, $\delta = +6^\circ 25' 19''.5$ als Anfang der Coordinaten zu betrachten.

XI.

Es sind noch zwei nicht zum Sternhaufen GC. 4410 gehörige Sterne ausgemessen worden, und zwar BD. +6°37'52

Utrecht, Juli 1897.

(9^m2) auf vier, und BD. +6°37'44 (9^m2) auf einer Platte. Dieselben hatten ursprünglich als Anhaltsterne dienen sollen, doch bin ich davon aus verschiedenen Gründen zurückgekommen. Nach gehöriger Reduction finde ich für 1895.0:

BD. +6°37'52 $\alpha = 18^h 21^m 16^s.19$ $\delta = +6^\circ 47' 45''.75$
+6.3744 18 20 28.38 +6 35 53.2

A. A. Nyland.

On the Cause of the Failure of Photography to depict delicate phenomena which are projected into a luminous field.

By T. F. F. See.

In two previous articles (A. N. 3438), I have discussed the effect of air currents and air waves upon the definition of telescopes, and have shown that these obstacles set a practical limit to visual work, and that the limit is relatively contracted with very large apertures. These atmospheric causes give rise to a large number of partial images, one resulting from each wave visible in the field, and their superposition produces one imperfect composite image; rapid internal change of this multitude of points or luminous blot

is the inevitable result of the constant motion of the currents. The wave motion has the effect of shifting the positions of the partial images at insensible intervals by measurable quantities, varying, according to some measures of Mr. Douglass and the writer, by the amount of from 0".5 to 2".0, even when the seeing is good enough to do fairly satisfactorily work on moderately difficult objects. As the eye catches by glimpses every combination of the partial images which gives good definition, the observer is enabled to detect

by uninterrupted watching very minute and very delicate phenomena which are not continuously visible. This result is due to the fact that the effect of the light upon the retina is not cumulative, but each instant produces its own impression, and the mind without any special effort readily determines what is permanent and what is transitory.

The photographic plate on the other hand has no such power of utilizing those instants of good seeing, when the partial images are well superposed, and hence by taking the impression of all the momentary irregularities due to the waves, obtains an enlarged and blurred image, which, being a composite photograph of many successive changing images, can give nothing but the average result for the whole time of exposure. We have remarked that the partial images of a moderately small star range from time to time over an area of 0.5 to $2''$, and hence it is clear that phenomena smaller than say $0.5''$, could not be taken by photography, even if there were no spreading of the light on the plate, no bodily motion of the whole image, and the motion of the telescope were perfect. Add to this impossibility of photographing objects having smaller diameters than $0.5''$ the well known spreading of the light on the film, and it does not seem possible that anything requiring even a moderate exposure could be detected in a luminous field (where the contrast is not very great) when the diameter of the image is less than about $1''$.

It is well known that the images of the smaller stars are much enlarged on the plates even when the driving of the telescope is perfect; the cause of this result, aside from the spreading of the light on the film, is undoubtedly the air waves, which continually change the shape and relative positions of the partial images in the focal plane. When images are produced by an indefinite number of neighboring points, the result on the photographic plate can be nothing but a mass of blurred light, in which practically all contrast is obscured. It is for these reasons, I think, that previous efforts to photograph the surfaces of the planets have failed, and from the known character of the causes at work, it does not seem probable that any considerable improvement in this direction can ever be effected.

Thus the quality of cumulative impression possessed by the photograph but not by the eye, and justly counted a gain for the former in detecting faint objects on a dark background, where the contrast is complete, in the case of objects projected in a luminous field, where images from different portions overlap, under the disturbing influence of the waves, leads to an inevitable failure, because of the cumulative action on the film, in addition to the old difficulty of spreading. When effort has been made to photograph small strongly marked spots in a luminous background with a moderately short exposure, a blurred and indistinct image has been the result; if the object were faint as well as small no trace whatever could be obtained. In this connection it is interesting to recall the historical fact that no satisfactory photographs have yet been made of the surfaces of planets so brilliant and easy even as Jupiter and

Saturn, while all effort to detect the canals or moderate sized markings on Mars have failed utterly. It is also worth noting that even the best photographs of the moon with the greatest telescopes are easily surpassed as regards sharpness and multiplicity of detail by the visual image of a good six-inch telescope. And in the case of the moon the conditions for successful photography are the most favorable; for the required exposure is very short, and the atmospheric difficulties are reduced almost to a minimum. When the photographs are taken with a large aperture, the blurring effects of the atmospheric disturbances are of course greatly increased. In view of these obstacles it is hardly necessary to remark on the absolute impossibility of detecting by photography the markings recently discovered on Mercury and Venus and on Jupiter's Satellites. In the case of Mercury and Venus the difficulties already mentioned would be enormously augmented by the additional obstacle of the light of the sky during the day, the only time when these planets can be studied. Moreover, since the lines upon their surfaces are fine and delicate and the surrounding surfaces are very bright, they could scarcely be photographed even in a dark sky under the best atmospheric conditions.

In closing these remarks it may be pointed out that for the reasons mentioned above photography is equally incapable of recording small and delicate irregularities in bright nebulae; though when the phenomena are not very small and the neighboring portions are not of very unequal brightness, an advantage is possessed by the plate in its accumulative power during a long exposure. If the curves or lines in the nebulae be very small or delicate, as frequently happens, the wave motion from the moderately bright region about them would cause total obliteration of detail on the resulting photograph; besides in long exposure the brighter portions come out too bright and the fainter portions too faint. In conversing with Sir William Huggins about the delineation of double nebulae, while in London during 1892, he expressed the opinion that faithful eye drawings of a conscientious and skillful observer would be better than any photographs that could be obtained. This opinion based upon the long experience of so great an authority, and the additional theoretical difficulties discussed above, convince me that some of the ideas heretofore current — that almost all celestial phenomena could be successfully photographed — will have to be abandoned. Judging by the past history of planetary and nebular photography it does not seem probable that plates of long exposure could ever disclose small or delicate details projected in a luminous field, because the ever-present waves in the atmosphere greatly obscure or entirely obliterate all contrast of neighboring phenomena. Thus it is likely to happen, as has often been the case before, that for the study of different classes of observational phenomena different methods will be required; and it seems fairly certain that the eye-method will always have a large field in the delineation of the more minute and delicate phenomena which engage the attention of practical astronomers.