best equipped universities, and the granting of this higher degree would serve to stimulate higher ideals and better preparation in the physicians all over the country entering the practice of the specialties. Finally, it is the duty of the faculties of our university medical schools to solve this problem by standardizing the work required in the preparation of the specialist. Our great need is not for more specialists, but for properly trained specialists.

THE PRESENT STATUS OF PHOTOTHERAPY.*

JAY FRANK SCHAMBERG, M.D.

PHILADELPHIA.

Since the epoch-making researches of Finsen, light therapy has been accorded an established place in the treatment of disease. Finsen and his disciples have conclusively proved the value of this agent in the cure of one of the most disfiguring and distressing of cutaneous maladies—lupus vulgaris. Not every patient with lupus vulgaris can be cured by phototherapy, but this treatment produces a larger number of successes and effects better cosmetic results than any other known procedure. The best results are obtained when all of the details of treatment are carried out in accordance with the teachings of Finsen. In the light institute of Copenhagen, which city has become the mecca of lupus patients, better results are obtained than elsewhere by reason of the skill of the physicians and the experience of the nurses. Treatment with the high power (50 to 80 amperes) Finsen arc lamps is not at all adapted to office use because of the size of the apparatus, the cost of the current, the duration of the individual treatments and the imperative services of a trained assistant. The same reasons militate against the successful operation of such lamps even in hospitals. In the city of Philadelphia there are not enough lupus patients to keep a large lamp going without great cost to the institution or to the patients. We have a large lamp in one of our hospitals in Philadelphia (Fig. 1), but it is only occasionally used. I have been told that no New York hospital is now actually using the large Finsen apparatus.

Many efforts have been made to improve the construction of lamps so as to lessen the cost of operation and the duration of the exposures. With the possible exception of the Finsen-Reyn lamp, these efforts have been unsuccessful. Lamps of the Lortet-Genoud and London Hospital type (Fig. 2) and the iron electrode lamps treat a larger surface and cause more rapid reaction, but are decidedly lacking in penetrative effect.

THE BASIC PRINCIPLES OF LUPUS PHOTOTHERAPY.

The rationale of light treatment in lupus vulgaris is based on three main propositions: (1) The property of certain rays of the spectrum to destroy micro-organismal life. (2) Their ability to penetrate the skin. (3) Their power to produce reactive structural changes. Success in the light treatment of lupus depends, therefore, on the ability of the rays to penetrate, to kill bacteria and to inflame.

The destructive effect of light on bacteria was proved as far back as 1876 by two English workers, Downes and Blunt, in an investigation most remarkable for its accuracy of detail. The results were confirmed by the researches of Arloing of Lyons in 1885. Numerous in-

vestigators, including Finsen, Bang, Bie, Busck, Freund, Strebel and others, have further proved the bactericidal influence of light.

Bang showed that unconcentrated light from a 30 ampere arc at a distance of 30 cm. will kill a surface layer of tubercle bacilli in six minutes.

The strongest bactericidal sun's rays are absorbed in the air, and thus an arc lamp has been found superior.

Dieudonné observed that bacteria were killed in half an hour by direct sunlight, in six hours by diffused daylight, in eight hours by electric arc light of 900 candle power, and in eleven hours by the electric incandescent light.

Other investigators have demonstrated that concentrated sunlight will check germ growth in one minute and kill bacteria in from five to seven minutes. Concentrated electric light checks growth of bacteria in from four to five minutes and kills in from fifteen to twenty minutes. Arc light from metal electrodes and the electric spark kill micro-organisms in from five to forty seconds.

Von Jansen demonstrated that a piece of skin inserted between the light source and the bacteria did not prevent their death when the thickness of the skin was 1.2 mm. and the exposure one and one-fourth hours. With 1.5 mm. thickness bacteria were no longer killed. Bacteria were weakened at a distance of 4 mm. in the tissues.

Busck killed pneumococci inoculated into the cornea of rabbits with light, and Nagelschmid destroyed tubercle bacilli experimentally inoculated into the skin of a guinea-pig.

On the other hand, Klingmüller and Halberstädter claim that the Finsen light does not kill the tubercle bacilli deep in the tissues. They exposed pieces of lupus tissue for ten minutes to the Finsen lamp and then inoculated them into the peritoneal cavity of guinea-pigs. That the bacilli were alive was evidenced by the fact that tuberculosis resulted.

It has been demonstrated that the bactericidal effect of light is largely exerted by the blue, violet and ultra-violet rays. The most scientific experiments on this point were carried out by Marshall Ward, who broke up light by passing it through a rock-crystal prism and allowed the spectrum to be cast on an inoculated solid culture medium. He found that the bacteria-killing effect began first in the transition between green and blue, and was strongest on the border between blue and violet. Especially when electric light was employed the bactericidal influence of the light extended far into the ultra-violet region. The dominant ability of the ultra-violet rays to kill bacteria, which was first demonstrated by Ward, was independently determined by Finsen three years later.

More recent studies by Bie² have established the important fact that the blue-violet end of the spectrum does not exclusively possess bactericidal properties. He says: "I found that with the application of a sufficiently strong light all portions of the spectrum were able to restrain the growth of bacteria or to kill them. The red rays possess the weakest power; this power then increases as one approaches the ultra-violet end of the spectrum, where it is strongest." Four per cent. of the bactericidal effect was ascribed to the red, yellow and green rays, while the blue, violet and ultra-violet rays exercised about 96 per cent. of the effect. Red, orange,

* Read in the Section on Cutaneous Medicine and Surgery of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

1. Deutsche med. Wochschr., 1905, No. 14.

2. Die Anwendung des Lichts in Der Medizin, Wiesbaden, 1905.

yellow and green act only after long exposures. Pure red requires one and one-half hours to produce a determinable effect on bacteria.

Some difference of opinion exists as to the manner in which the tubercle bacilli in the skin in lupus are destroyed. Finsen held that their death is due to the exclusive effect of the ultra-violet rays. This view is not universally concurred in, many German workers contending that they are destroyed by changes induced in the tissues by the light. Roux established the fact that light exerts an important influence on nutrient media. He showed that Arloing was in error in regarding certain spores treated with light as dead, and proved that they would grow when transplanted to a fresh unexposed medium.

Richardson demonstrated that the sterilization of urine by exposure to light in the presence of air is due to the formation of hydrogen peroxid. Dieudonné confirmed this observation and agreed with Richardson that the ability of light to kill bacteria is due to the formation in culture media of hydrogen peroxid. Kruse, who carried out the same experiments, did not regard the results as due alone to change in the media, but also to a direct action of light on the bacteria. Bie admitted that hydrogen peroxid is formed in many nutrient media on exposure to light, but claimed that bacterial death is more rapid in media in which no hydrogen peroxid can possibly form.

Downes and Blunt early suggested that the destruction of bacteria by illumination depended on an oxidation process and required the presence of free oxygen. Bie's experiments led him to the conclusion that the destruction of the bacteria is not an oxidation process in the sense that the presence of oxygen is a necessary condition. The stronger the light in the blue violet and ultra-violet rays the less does its bactericidal effect depend on the presence of oxygen, and *vice versa*, the fewer the chemical rays that the light contains the less able is it to kill bacteria without the aid of oxygen.

The researches of Hertel³ are most interesting in this connection. Hertel claims that the short-waved rays of light (chemical rays) cause a splitting off of oxygen from the blood and cell plasma, much in the same way as oxygen is freed from plants by light, and that this process affects both the tissues and the bacteria.

PENETRATION OF THE RAYS OF SPECTRUM.

It has been shown that the penetrating power of the various rays of the spectrum is in inverse proportion to their bactericidal and chemical power. The red rays are the most penetrating and the ultra-violet the least.

Most of the efforts in phototherapy in the past have been devoted to securing deep penetration of the chemically active rays. Within the past few years the attempt has been made to increase the chemical activity of the penetrating rays. In 1900 Tappeiner and Raab⁴ made the important discovery that infusoria can be sensitized to the action of certain rays of light by the addition to their nutrient media of certain chemical agents.

This work was confirmed by Dreyer⁵ working in the Finsen Light Institute in Copenhagen. Dreyer demonstrated that the addition to the nutrient medium of a small quantity of erythrosin renders bacteria sensitive to the action of the yellow and green rays of light. The germs which ordinarily are not influenced or but slightly

influenced are rendered sensitive much in the same manner that a bromid of silver plate is sensitized to these rays. The microbes thus sensitized may be killed by the passage through moderately thick layers of skin of light from the usually employed sources owing to the increased action of the highly penetrating yellow and yellowish green rays. The *Bacillus prodigiosus*, protected by a layer of skin 1.25 mm., was not killed by exposure to light for eleven hours; after being sensitized death occurred after twenty minutes exposure.

Animal tissues, such as frog's tongue, rabbit's ear, human skin, etc., were also rendered sensitive to the action of the yellow and green rays.

In the treatment of lupus the exposures are greatly shortened by sensitizing the tissues.

Later Tappeiner and Jesionek⁶ employed, before exposing to light, a 5 per cent. eosin solution as a paint and also subcutaneously in the treatment of superficial epithelioma, lupus, flat condylomata, tuberculous ulcer, etc., with alleged good results.

Lenkei⁷ found that blue and yellow rays penetrated the skin and connective tissue to a depth of from 1 to 2 cm. When no muscular layer was present traces of the blue rays penetrated to a depth of over 3 cm. A small portion of the light penetrated to 5 or 6 cm. even through the muscular layer, but this was exclusively yellow rays. The red rays could not be examined because the ordinary light, sensitive substances are not changed by them, but Lenkei is certain that these last rays penetrate into the depths in larger quantities than the yellow. Five per cent. of the light consisted of blue rays and 80 per cent. of yellow and green.

Finsen and Jansen have compared the penetrating power of light from various lamps, namely, the Finsen, the Finsen-Reyn, the Lortet-Genoud and the Bang. They found that the two latter are distinctly inferior to the first two lamps when deep penetration was required.

It appears to me that these tests, which are based on the influence of light on silver salts, are open to error. It by no means follows that the chemical effect of rays of light on photographic plates is a measure either of the bactericidal power of the rays or of their ability to produce inflammatory changes in the tissues. I have found that an ordinary sixteen-candle-power incandescent lamp with a reflector will emit light which will pass through the skin, subcutaneous tissue and mucous membrane of the cheek and affect a photographic film in the mouth in thirty seconds (Fig. 9).

On the other hand, a light extremely rich in chemical rays (the uvioi lamp) was unable to produce any effect in five minutes.

Photographic plates may be affected by the yellow and even the red rays, and these rays are the most penetrating.

Busck⁸ found that red, orange and yellow rays passed through his hand and affected a photographic plate in five seconds. The blue, violet and ultra-violet rays from the same light source produced no effect on the plate in ten minutes. This will illustrate the comparative penetrability of the various rays.

HISTOLOGIC CHANGES.

The histologic changes effected in the skin by the action of light have been carefully studied by a number of

3. Ztschr. f. Allg. Physiol., iv-v.
4. Münch. med. Wochschr., 1900.
5. Derm. Ztschr., 1903, x, No. 6.

6. Münch. med. Wochschr., 1903, No. 47.
7. Die Durchdringungsfähigkeit der blauen und gelben Lichtstrahlen durch tierische Gewebe, von W. Lenkei-Almadi Orvosi Hetilap, 1906, No. 27.
8. Mitteilung aus Finsen's Lys Institut, iv, 1903.

observers, including Moeller,⁹ Sack,¹⁰ McLeod,¹¹ Schmidt and Marcuse,¹² Zieler,¹³ Meirowsky¹⁴ and others.

The findings are in general agreed on. There is a pronounced dilatation of the superficial and deep cutaneous blood vessels with exudation of leucocytes. Sack believes the first change to be a proliferation of the endothelial lining of the blood vessels. Meirowsky, working in Unna's laboratory, states that the light stimulates the epithelial cells and leads to nuclear division. After intense exposures degenerative changes in the epidermis may take place and the dead cells may be cast off through the formation of blebs.

The light also causes an increase in the number of connective tissue cells and a swelling of the collagen. At the height of the process the rete mucosum is thickened. Unna says that light makes the skin dense and harder, in that under the influence of the light the protoplasm is reduced to keratin. Moeller demonstrated the fact that light caused a hyperplasia of the epidermis and an abnormal cornification.

THE INDICATIONS FOR THE USE OF PHOTOTHERAPY.

The treatment of lupus vulgaris represents the dominant field of usefulness of light therapy. Nevertheless, there are other diseases in which light may be employed with advantage even though it does not supplant other methods.

Light would doubtless have been more used as an auxiliary therapeutic measure in many cutaneous diseases had not the suddenly established reputation of the x-rays cast a shadow of obscurity over the milder acting agent. The ability of the x-rays to produce rapid and profound changes in cells and tissues constitutes at the same time the advantage and the disadvantage of that treatment. When judiciously and carefully applied in properly selected cases the x-rays certainly represent a wonderful addition to our therapeutic armamentarium; but improperly used they are capable of doing harm, often irremediable harm. Light energy, on the other hand, while much slower and much milder in its effect, is a perfectly safe remedy. Patients themselves have discovered this difference and will often permit one to use light therapy while refusing Roentgenization.

The supplementary list of diseases in which phototherapy has been alleged to be of value includes lupus erythematosus, alopecia areata, acne, acne rosacea, flat vascular nevi, certain forms of eczema, leg ulcers, etc.

LUPUS ERYTHEMATOSUS.

In this disease, one of the most refractory of dermatoses, phototherapy may often be used with the happiest of results. Finsen's first report included thirty-one cases with eleven cures and ten patients still under treatment. Since then a larger percentage of cures has been obtained by him. Leredde and Pautrier mention twenty-three cases with eleven cures, many partial cures and three failures. Morris and Dore treated eleven cases with great improvement in seven. Hyde, Montgomery and Ormsby report a total of nineteen cases, with five cures, nine patients greatly improved, three with but slight improvement and one made worse. I have had several cures in lupus erythematosus of the vascular

type; it is in this particular form in which phototherapy is apt to be useful. The x-rays are often valuable in the hypertrophic variety, whereas they are apt to aggravate the condition in the vascular type.

ALOPECIA AREATA.

This occupies a place among the diseases in which light therapy is indicated. Finsen, Forchhammer, Jersild, Leredde and Pautrier, Sabouraud, Hyde, Montgomery and Ormsby, Kromayer and others have reported successes. Professor Kromayer of Berlin has had such remarkable results that special reference to them is deemed valuable:

Kromayer¹⁵ uses an arc light with iron electrodes, without pressure lens; a current of from 15 to 25 amperes is employed. The exposures range from one-half minute to ten minutes for each area, at a distance of from four to ten centimeters. In some cases Kromayer claims that a regeneration of the hair is evident after the first exposure. On the second day new hair points are at times visible. At first this was not believed possible, but was confirmed by independent examinations. Kromayer treated thirty-three cases of alopecia areata, which he divided into two classes: The milder ones with a hair loss of less than half of the head, and the severe cases with hair loss greater than this or complete. The result in the thirteen milder cases was throughout good; nearly all were completely cured. In three, small bald areas were still present; in a few, relapses were noticed. Of the twenty cases of the second group thirteen were total alopecia and the other seven approached total alopecia in various degrees. Of these twenty patients only five were treated with negative results, only one of which was not a total alopecia; hair growth was stimulated, but the hair either ceased to grow further or fell out.

Summing up his results, Kromayer claims twenty-eight successes out of thirty-five cases. In the milder cases the results were all satisfactory. In the severer group there were five failures among twenty cases. Many of these cases had lasted for years and had been treated with all sorts of methods without results. One patient, for instance, was a woman of 50 years of age whose hair loss was almost total and had dated back twenty-five years. After twenty-seven treatments the hair was restored to normal condition except over the temples, where it is a little thin.

Kromayer concludes that the iron light at the present time is the best and surest remedy in alopecia areata.

The stimulating effect of the chemical rays of light on hair growth has been repeatedly observed in Copenhagen, where the bare arms of the nurses giving Finsen light treatment have exhibited increased pilary development.

HIGH-POWER INCANDESCENT LAMPS.

The much advertised high-power incandescent lamps have been viewed skeptically and not with much favor by dermatologists in general. The skepticism has doubtless been based on preconceived theoretical considerations. Subsequent experience may prove the lack of faith to have been well based, but in the light of newer researches as to the effectiveness of all rays of the spectrum, and as to the efficacy of heat rays, trial of such light sources by properly qualified and conservative men would be desirable. It will be recalled that Bie found that with the application of sufficiently strong light all

9. Bibliotheca Medica, Part D, 11.

10. Muench. med. Wochschr., July, 1902.

11. Brit. Med. Jour., Oct., 25, 1902.

12. Arch. f. Derm., 1903, p. 323.

13. Derm. Ztschr., 1906, No. 1.

14. Monats. f. Prakt. Derm., April, 1906, xlii.

15. Monats. f. Prakt. Derm., July 1, 1905.

portions of the spectrum were able to restrain the growth of bacteria or to kill them.

Audry and Krösing¹⁶ have been enabled to heal venereal sores rapidly with heat rays. Holländer claims good results in lupus and chancres with his hot air treatment. Scholtz¹⁷ believes that the heat rays in the Finsen treatment of lupus are by no means without effect. He is of the opinion that the chemical rays are absorbed in the superficial layers of the skin and that the death of the bacilli in the deeper layers is due to the heat rays.

MERCURY VAPOR LAMPS.

In 1892 and 1896 Dr. Leo Aarons,¹⁸ a noted German electrician, demonstrated that a continuous electrical current passed through a vacuum tube containing a certain quantity of mercury is capable of creating an arc and of generating an intensely bright light emitted by the incandescent mercury vapor. Peter Cooper Hewitt of New York improved this lamp and made it commercially practical. This light contains few or no red rays and is extremely rich in blue, violet and ultra-violet rays. The latter, however, are largely absorbed by the glass of the tube itself. To obviate this absorption of the most chemically active rays Dr. E. Zehimmer, working in Schott's glass works, produced a glass (said to be a barium phosphate-chrome glass) which is pervious to the ultra-violet frequencies and which, according to Schott, permits by far the greater portion of the short waves originated in the glass tube to emerge. These tubes, which are made of varying lengths, must be tilted to make the arc and to start the light.

The lamp made by Schott is called the "uvio" lamp, the name being a convenient contraction of the term ultra-violet. Another mercury vapor lamp made of quartz with a water-cooling jacket is made by Heräus of Hamburg. For several years these lamps have been in use in Germany.

Axmann¹⁹ treated with the uvio lamp thirteen cases of leg ulcer with cures after a treatment of some weeks; some of these cases had lasted years.

Acute and weeping eczema, after a ten-minute treatment were alleged by him to be so far improved on the following day as to require no further treatment. Acne, furunculosis and alopecia areata are also said to have been cured.

Hans Meyer²⁰ found that the uvio light acted on photographic paper through a piece of epidermis of a thickness of 0.5 mm. in three-fourths of a second at a distance of 1.5 mm. (The epidermis was taken from bullous lesions.)

Keller determined that the uvio light had a bactericidal influence, but acts only superficially. Plates exposed immediately after inoculation were affected, but later growth was not influenced.

Meyer had no success in ringworm of the skin, but secured favorable results in tinea versicolor. He believes that the uvio light, like the iron light, constitutes a method that surpasses others in the treatment of alopecia areata.

Meyer noted, as did Kromayer with the iron light, that fine hair tips begin to be visible after one treatment. That the light is the exclusive factor may be

determined by comparing areas treated by the light with those that are not. He obtained good results in severe acne indurata. The acne nodules under the influence of the hyperemia undergo resorption.

In a case of bullous dermatitis herpetiformis Meyer obtained an apparent cure. The blebs dried up in a very short time under the influence of the light erythema. Excellent results were also obtained in the prurigo of Hebra.

Kromayer²¹ considers the following indications for the use of the mercury vapor lamp (the author employed the quartz lamp made by Heräus): Lupus and superficial epithelioma, vascular nevus, alopecia areata, acne rosacea, eczema, rebellious syphilitic affections (psoriasis linguæ) and ulcers that are often refractory to ordinary general and local treatment. Kromayer obtained successful and sometimes brilliant results in isolated cases rebellious to all other methods.

Assfalg of Frankfurt²² used the uvio lamp for one year. He believes that it is necessary to produce a light reaction in order to obtain therapeutic results. He treated three patients with alopecia areata and cured them with an average of fifteen sittings of from one-half to one hour in duration. The cases had lasted for at least a year and the patients had been treated in various ways without result. In several cases he saw extension of patches of alopecia areata on the border at the same time that new hair was growing in the center. He believes the hair returns more quickly under uvio treatment.

Acne was found to be very favorably influenced. Assfalg preferring the light treatment to the x-rays. A case of furunculosis of the entire back and chest of four years' duration and treated with various remedies was completely cured after ten treatments of from one-half to three-fourths hour in duration. There was no return within nine months. Two patients with chronic seborrhoeic eczema were completely cured after five and eight treatments respectively. Two patients with acne rosacea were considerably improved after three treatments and are still under treatment. One is cured now for six months. A number of patients with chronic eczema, dry and moist, have also been cured.

Two patients with varicose ulcers were treated. One, the ulcer the size of a half-dollar, was healed after six irradiations, the other, still under treatment, was much improved after three treatments.

Gastou and Nicholas²³ report several patients treated with the Heräus quartz lamp; patients with lupus erythematosus, nodular tuberculid and glandular swelling were cured. The writers prophesied a great future for the lamp.

The uvio lamp has been employed in the Philadelphia Polyclinic for about a year. The breaking of the lamps interrupted the treatments, and some of the patients were obliged to be treated by other therapeutic measures. I have also used the uvio lamp in my office for some months. I have had constructed a tilting lamp-holder on a stand to receive two short tubes run in series and so arranged that the lamps could be used horizontally or vertically (Fig. 6). Above the lamps is a reflector covered with aluminum paint.

With this lamp a light reaction may be produced after five minutes' exposure at 4 centimeters distance. Several times, after ten minutes' illumination at from 4 to 8

16. Arch. f. Derm., xliii.

17. Berlin klin. Wochschr., 1904, No. 18.

18. Wiedemann's Annalen, 1892 and 1896.

19. Deutsch. med. Wochschr., 1906, p. 583.

20. Einige Erfahrungen mit der Uvio Lampe, Med. Klinik., September, 1906, p. 990.

21. Deutsch. med. Wochschr., 1906, No. x.

22. Muench. med. Wochschr., Oct. 9, 1906.

23. Bull. Med., 1905, No. 57, p. 668.

cm. distance, I have produced an erythema with some puffing of the eyelids. The erythema largely disappears at the end of twenty-four hours. The parasiticide properties of this lamp in my experience appear to be rather feeble. Petri dish agar cultures of the *Staphylococcus aureus*, *Bacillus coli communis* and the *Bacillus capsulatus aerogenes* were not killed after twelve minutes' exposure at 4 cm. with the Petri dish lid off. The cultures were about twenty-four hours old and it is possible that the effect of the light was insufficiently penetrating to kill the bacteria growing down into the medium.

I have used the light in alopecia areata, acne, eczema, lupus vulgaris and leg ulcers, but in entirely too few cases to justify any definite conclusions. I will briefly narrate the results and state my impressions:

CASE 1.—A. G., an Italian boy, aged 12, had an extensive alopecia of 14 months' duration. A partial regrowth at one time took place, but was followed by a subsequent loss of hair.

Fig. 1.—Large Finsen lamp in use at the Stetson Hospital, Philadelphia.

The usual local remedies were used without much effect. About fifteen light treatments of ten minutes' duration, at a distance of from 4 to 6 inches, were given in the course of from six to eight weeks. The hair soon began to grow and is now fully restored over the entire scalp except at the nape of the neck, which received no illumination (Figs. 3 and 4). This fact strongly suggests the actual curative influence of the rays in this case.*

CASE 2.—J. F., aged 12, a robust blond boy with a total alopecia of nine months' duration. The usual treatment was entirely without avail. Under uviole exposures a fine down appeared, which subsequently vanished. Fourteen treatments were given, but without any successful result.

CASE 3.—M. R., aged 21, a young woman with a half dollar sized patch of alopecia in the right temporal region of two weeks' duration. Three light treatments of from 15 to 20

minutes were given. There was extremely rapid restoration of hair. A stimulating antiseptic wash had been used and the cause of the unusually prompt result can not be definitely stated.

CASE 4.—In connection with these cases of alopecia areata, and bearing on Kromayer's successful results, it is proper to note a failure in the case of a 12-year-old girl with a total alopecia of ten years' duration. The girl was treated in the light laboratory of the Philadelphia Polyclinic with the London Hospital lamp. Thirty-two twenty-minute exposures were given without result. The accompanying illustration (Fig. 5) shows the light reaction in areas covered by the lens.

CASE 5.—T. S., man, aged 69, with damaged heart and kidneys and arteriosclerosis, had an extensive almost universal erythematous eczema of many years' standing. There was considerable thickening and lichenification. Itching was severe. Internal and local treatment of varied character had given no relief. The itching and infiltration were distinctly lessened by the light treatment. On the breakage of the tubes, x-ray treatment was substituted with more benefit than that derived

Fig. 2.—London Hospital lamp in use at the Philadelphia Polyclinic.

from the light treatments. The patient is now vastly improved.

CASE 6.—M. M., a girl, aged 9, with a severe papular eczema of the face, hands and arms of several years' duration. Uviole treatments of fifteen minutes' duration were given twice a week. After four treatments, the itching was markedly diminished and the eruption greatly improved. Treatment was discontinued on account of breakage of apparatus.

Several cases of acne were treated, but inasmuch as these were private patients sole reliance was not placed on the light treatment. In non-suppurative papular or nodular lesions rapid resorption is effected by ten minute light-exposures. The influence of the light is much more prompt but less permanent than that of the x-rays.

In several leg ulcers treated granulation was markedly

* The patient has remained well now for five months.

stimulated by a few treatments. One severe case appeared to be on the road to cure, when the breakage of apparatus caused substitution of mild x-ray treatment, with immediate aggravation and extension of the ulcers.

In this connection the observations of Strebel are pertinent:

Strebel showed that ultra-violet irradiation of open

Fig. 3.—Patient in Case 1. Alopecia areata of 14 months' duration cured with the uvioi lamp.

Fig. 4.—Showing restoration of hair.

wounds causes rapid granulation and skinning over. He explained this by the favorable action of the ultra-violet light on the blood vessels and trophic nerves. He recommended ultra-violet light in the treatment of leg ulcers.

Axman found uvioi light of great value in leg ulcers.

In summing up my impressions as to the value of the mercury vapor lamp I should say that it appears most capable of accomplishing good in alopecia areata, leg ulcers and certain forms of eczema. The light is rich in chemical rays, but lacks deep penetration. These lamps have the advantage over others that a broad volume of light is emitted which can be conveniently applied over large areas. Another very pronounced advantage is that the grade of light erythema desired may be produced and even predetermined by the distance and duration of the exposure. The histologic changes produced by light have already been referred to; that a dilatation of blood vessels with exudation of fluid and corpuscles and regressive changes in pathologically altered cells occurs after the production of a distinct light erythema is evidence that in properly-selected cases such treatment may prove beneficial.

Such phototherapeutic measures are not to be regarded as panaceas but as aids to other approved agents in the treatment of certain cutaneous diseases.

DISCUSSION.

Dr. W. S. GOTTHEIL, New York City, said that in a corner of his waiting room there stands a big apparatus containing

Fig. 5.—Patient in Case 4.—Alopecia areata of 10 years' duration treated without success with the London Hospital lamp. Areas of light erythema shown.

a 60-ampere arc light—the American form of the Finsen apparatus—unused for many months. He employed it persistently for a long time, and did get results, some of which have been published; but he hardly uses it at all now. His experience can probably be paralleled by some of the members present. He knows of one case, at least, in which a very costly Finsen apparatus was imported, and, when he last heard of it, it was still unpacked, owing to the amount of space that it required, and the time that its use would consume. In Copenhagen it may be possible to give daily treatments of an hour each for months at a time, but Dr. Gottheil's experience is that patients here will neither submit to it nor pay for it. Furthermore, the applicability of the method is limited. In alopecia areata he has obtained no results that were not attainable by simpler means. Some patients with lupus erythematosus were greatly improved and apparently cured, but some relapsed, and he has come to believe that one can never tell when this treacherous disease is ended. In lupus vulgaris the x-ray and ordinary mechanical methods of treatment are generally preferred. The Copenhagen results, according to all reports, are the best that have yet been attained

in this disease, but there must be great simplification of the method and shortening of the time required before it can be used extensively here. So far as penetration of the body by light rays is concerned, Dr. Gottheil proved some years ago that a sufficiently powerful beam of light can be made to penetrate the entire thickness of the human body and print on a photographic plate. All the rays, even the ultra violet, can be made to pass through the surface of the body and reach the internal organs. A necessary condition, however, is to have a light beam of sufficiently high candle power; 50 or 100 candle-power incandescents are toys, not efficient therapeutic agents. A 30,000 candle-power arc ray, concentrated as much as possible, and cooled, from a 60-ampere current, is as small a light as should be used for action on anything deeper than the epidermis.

DR. L. DUNCAN BULKLEY, New York City, said that several years ago he went to Copenhagen, and, after a careful investigation of the Finsen method of treatment, became enthusiastic regarding it, and purchased one of the lamps. He had it shipped to New York, properly set up and wired from the street, with a special attachment to allow water to run through the lens. That was two or three years ago. He has used it in two cases of lupus, and for the past year or two it has remained untouched. He is still strongly in favor of the

visited Copenhagen and spent some time at the Finsen Light Institute, and saw many patients treated. He was impressed with the great care with which Finsen's assistants employed this method and by the benefit that was derived from the treatment, especially in lupus erythematosus and nevus. After getting better acquainted with some of the assistants, he learned that the consensus of opinion seemed to be that the best results are obtained in lupus erythematosus and vascular nevi. Like Dr. Bulkley, he became enthusiastic regarding the treatment, and made arrangements to have instruments and nurses sent to this country. He afterward visited Berlin and saw Finsen's apparatus employed, but the important details of the technic, such as the proper pressure and focus, were not observed. The results obtained were indifferent, and he believes the operators regarded it as an indifferent method. He then visited the clinic of Lange, who was carrying out two series of experiments, one by the well-known surgical methods, and another series by the Finsen light, but did not remain long enough to see the results of these experiments. He afterward saw a report in print that the light method has no special

Fig. 6.—Method of using two uviole lamps in series.

Copenhagen method of treatment, and if a bad case of lupus vulgaris, with lesions about the face, should come to him tomorrow, he would advise the patient to go to Copenhagen. Finsen's results, as shown at the dermatologic congresses, both in Paris and Berlin, were very brilliant, and Dr. Bulkley said that Dr. Hyde of Chicago is very enthusiastic regarding them. In Copenhagen Dr. Bulkley saw 150 patients treated, men and women who had had the disease for ten, twenty and even thirty years, and the results were simply superb. By the method pursued there the lens must be held over one little spot for an hour at a time, and afterward a salve is applied. A corps of nurses makes these applications, and some of the patients have had as many as 600 or 800 of them, the treatment sometimes continuing over a period of from three to five years. The results obtained are better than he has ever seen by any other method of treatment, but he thinks that if the x-ray is used rightly, excellent results can be obtained in lupus. It is not so satisfactory, however, as the method followed in Copenhagen, and it is apt to result in scarring and burns. He has had little experience with the other forms of light.

DR. W. T. CORLETT, Cleveland, said that four years ago he

Fig. 7.—Condition of redness two days after a radiation of 15 minutes' duration with the uviole lamp. One of the squares was kept under water during radiation in order to ascertain if the heating of the skin by the lamp during radiation had any particular effect. The two places did not differ in any respect. (Schott.)

Fig. 8.—Shadowgram of a triangular piece of lead on a photographic film in the mouth, the cheek in contact with the pressure lens of the large Finsen lamp (50 amperes current).

Fig. 9.—Shadowgram of a perforated piece of lead on a photographic film in the mouth, the cheek being one inch from a 16-candle power incandescent lamp, backed by a parabolic reflector. Exposure 30 seconds.

advantages over the ordinary surgical methods that Lange was using. In Paris he found that the Finsen light treatment is applied in an indifferent manner, but in London it is given as in Copenhagen, under specially instructed nurses. In short, the result of his tour of investigation made him somewhat skeptical as to any special advantage the Finsen light possesses over other less complicated methods, and he gave up the idea of ordering a lamp.

DR. J. FRANK WALLIS, Norristown, Pa., showed a hood, made of sheet iron, with an aluminum reflector, and which is attached to the ordinary arc lamp. From this lamp he can obtain intensely blue or violet rays, and with it he has from time to time treated small acne lesions. One advantage of

this lamp is that the nearer the carbon, the stronger is the blue light. In addition to acne lesions, he has used it for small abscesses and sebaceous glands. There are also some cases of eczema that react very quickly to this light.

DR. M. L. HEIDINGSFELD, Cincinnati, said that he has used a Heraeus quartz mercury lamp with the greatest satisfaction. He said that, in his opinion, it possesses decided advantages over other instruments used in phototherapy, particularly in the furuncular type of acne and in the treatment of small abscesses, on which it exerts an influence superior to any other form of light therapy that he has ever tried. In many of these cases he has used the x -ray and the London Hospital lamp, but mostly with indifferent results. He has also used a Kromayer quartz mercury lamp, and he has found the effects of this and the Heraeus instruments rather remarkable. They emit practically no heat at a distance of from six to eight inches, and, after an exposure of three or four minutes, they give rise to a decided bullous reaction. The effect of these instruments on certain forms of skin affections is striking and is almost remarkable in indurated and pustular acne; chronic forms of eczema and dermatitis, particularly infectious types, will be favorably influenced by a few applications of this form of light treatment. Cases of folliculitis and kerion, which resisted all forms of simple measures and rather prolonged x -ray treatment, yielded almost completely to one thorough application of this form of light energy. Some cases of lupus erythematosus, which baffle all other forms of treatment, x -ray and topical applications, have yielded very rapidly to this form of treatment. Others, however, have not been favorably influenced. Unlike Dr. Schamberg, he has not obtained any favorable influence from this form of treatment in alopecia areata, but it may be due to the fact that he did not push the treatment. In other respects his results do not differ materially from those of Dr. Schamberg, and he heartily endorses all the latter's favorable comments concerning this form of treatment.

DR. J. H. MUDGETT, Philadelphia, said that he has had considerable personal experience with light therapy, and he does not see how so many physicians, especially dermatologists, are so skeptical regarding it, and wonders whether or not they have any right to form an opinion about something they have never tried. With phototherapy it must be the same as with anything else; one must have experience with it before any opinion can be formed regarding it. Dr. Mudgett had five cases of ulcers of the leg last October which he treated successfully with an ordinary 500-candle power incandescent light, making hour exposures, and repeating the treatment every day and every other day. Some of them took two or three months to heal, but they were all long-standing cases—as long as ten years. In addition to the phototherapy he only used ordinary antiseptic applications, such as dermatologists recommend. In conclusion he suggested that dermatologists give this method of treatment a fair trial, in comparison with other methods.

DR. JAY F. SCHAMBERG, Philadelphia, said he believes that it is pretty well agreed that the treatment of lupus vulgaris remains the dominant field for phototherapy. Nevertheless, there are other dermatoses in which light treatment may be found useful. There is a distinction between phototherapy and actinotherapy. The blue, violet and ultra violet rays are not the only frequencies capable of influencing tissues. It has been shown that there is real efficacy in heat rays in certain conditions. The high-power incandescent lamps should be given a trial by competent and conservative workers. Extravagant claims have been made for various forms of light treatment which can not be indorsed. It has been shown that the penetrative power of light rays is in inverse proportion to their bactericidal effect.

Tainted Air.—The man who is 60 years old has spent 20 of those years in his bedroom. If his mind has been saturated with the old-time delusion that night air is dangerous, he has probably proceeded to make it a hundred-fold more dangerous by breathing over and over again the poisons that have been thrown off from human lungs.—*Good Health.*

BLASTOMYCOSIS IN AN INFANT.*

J. B. KESSLER, M.D.

Lecturer on Dermatology and Clinical Dermatology, State University of Iowa.
IOWA CITY, IOWA.

This is a peculiar disease, from a geographic standpoint. Its home so far seems to be confined to America, and only in a few localities, largely in the Mississippi valley, if we also include the city of Chicago. The forms on the Pacific coast seem to differ from those found in other localities in the United States and are named protozoic and coccidioidal.

In cases of protozoan infection the organism develops by endogenous spore formation and not by budding,

Fig. 1.—Blastomycosis. E. J., at 8 months, showing lesion on right cheek.

Fig. 2.—Blastomycosis. E. J., at 8 months, showing lesion on left cheek.

while in blastomycosis the only way of development in the tissue is by budding. The cultures also differ somewhat from those obtained in blastomycosis.

The first case reported from Boston was in the latter part of 1906. There were no recognized cases in England up to 1903; a doubtful one was reported from Madrid the same year. I have seen three cases in the past eight years, one of them somewhat doubtful.

I am indebted to Dr. W. W. Jones for referring to

* Read in the Section on Cutaneous Medicine and Surgery of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.