AN EXPERIMENTAL STUDY OF SLEEP*

BY BORIS SIDIS, A.M., PH.D., M.D., BROOKLINE, MASS.

CHAPTER XI

Motor Reactions and the Nature of Sleep

BEFORE we conclude the account of the experiments it may be of interest to make a brief statement of a statistical inquiry in regard to our mode of going to sleep. It is usually supposed that the course and mode of our activities are due to voluntary decisions, that they are irregular, changeable with mood and caprice as our whimsical will is. We are free agents and anything we decide to do we can do in any old way; we do it just as we please and it does not matter to us what the course and arrangement be, provided the will wills it so. As a matter of fact we are far more creatures of habit and instinct than of reason and will. This holds especially true of our fundamental reactions, such as the induction of sleep. If we ask the ordinary person on which side he goes to sleep or on which side he falls asleep most comfortably, the question puzzles him at first. At the first blush it seems to him he can go to sleep on either side; it does not matter. When the question is repeated to him and made clear and he begins to think about it, he soon finds out that he has a definite way of going to sleep. Should he try to go to sleep in a somewhat different manner, he would find it very difficult to fall asleep. We have definite ways of reactions which are due either to habit or to the structural arrangement and mode of function of our organism.

Manaceine in studying motor reactions in sleep has found that right-handed people react in their sleep with the left hand, while left-handed react with their right hand. I have similarly experimented on human subjects and could not confirm Manaceine's results. In my experiments on infants I found that in early infancy the reactions in sleep differ but little from that of waking state. The reactions are *undifferentiated*, but I found that later on in the course

* Begun in April-May number.

of the child's development, the motor reactions mostly conform to Pflüger's laws of reflex action.¹ To quote from my notes:

Boy two weeks old; sound asleep. I tickled his left nostril: bilateral reaction, both hands are thrown up at once. When the tickling is more intense or more persistent and prolonged motor reactions in both legs are induced. The same holds true in the case of the tickling of the right nostril, the reactions are bilateral and become more diffused bilaterally, if the stimulus is more prolonged or more intense.

The baby reacted in the same way in the waking state, so that there was no difference in the reactions of the two states.

When the baby is fast asleep, I tickle the right temple, the response is a slight reaction with the left. This is really accidental. On other similar occasions the reaction is of the right hand and sometimes the reaction is of both hands and also of both legs, according to the intensity or to the cumulative effect of the stimulation.

I invariably found that a slight stimulation given the first time produced a reaction, while a second and third stimulation of the same intensity often produces *no* result.

PFLÜGERS LAWS:

¹(I) Law of unilateral reflexes.

If peripheral stimulations cause contraction in only half of the body, the contraction always occurs on the same side as the stimulus, and in general those muscles contract whose nerves arise from that segment of the cord, to which the irritated sensory nerve belongs.

(II) Law of reflex symmetry.

If the effects of stimulating a sensory nerve upon one side extend to the other side, only such motor fibres are called into activity which correspond with those which are already excited on the stimulated side.

(III) Law of unequal contraction on the two sides.

If the contraction is unequal on the two sides, the stronger reflex is always on the side of stimulation.

(IV) Law of reflex irradiation.

(1) When stimulation of a cerebral nerve causes reflex contraction, the motor nerve concerned is invariably either in the same level as the sensory nerve, or it is further downward toward the *medulla oblongata*. (2) When stimulation of a spinal nerve causes reflex contraction beyond its own segment, irradiation always takes place toward the *medulla oblongata*.

(V) The law of three locations of reflex contractions.

Upon stimulation of a sensory nerve, reflexes can occur in only three parts of the body. These are:

- (a) at the level of the stimulated nerve.
- (b) in parts innervated from the *medulla oblongata*.
- (c) in the whole body.

Boy six weeks old. Tickling of the left ear excites at first movements in the head, in the trunk and then in both hands; when the stimulation is continued, the whole body becomes involved in the reaction, then the left hand is raised. The same holds true in the case of the right ear,—the movements are first in the head, then spread to the trunk, then to the lower extremities, then only to the side specially stimulated. In other words the reaction is at first *diffused* and then only becomes more special,—the hand is raised corresponding to the side stimulated. This mode of reaction holds true both in waking and sleeping states.

Boy twelve weeks old. I observed an interesting phenomenon which may be described as that of *associated movements*. The infant has both hands in his mouth. When one hand is taken away from the mouth the other goes off *simultaneously*, as if an invisible power has pulled it away at the same time.

Boy thirteen weeks old. When he was asleep, I tickled his left temple, he reacted with the left hand. Tickled his right temple, reacted with the left hand again. When the stimulus became more intense and prolonged, he moved his head, left hand, and also the right hand.

I tickled the left foot when the boy was asleep, both hands reacted, the left hand having the start; I tickled the right foot, both hands reacted, the *left* hand having the start again.

Tickled the *left* temple; moved *right* hand; tickled *right* temple moved *right* hand again. *The reaction is indeterminate*,—sometimes it is bilateral and sometimes crossed. The reactions do not conform to Pflüger's law. On the whole *bilateral reactions predominate*.

When the child was asleep, I passed a pencil over the whole length of the left side of his face,—he reacted with the left hand first and then started with both. I passed the pencil over the right side of the face, he threw up both hands, the right having the lead.

Now and then one can begin to observe the manifestation of Pflüger's law. When the child was asleep, I tickled his right foot; he drew away the foot. When the stimulation is slight, the right foot alone reacted; when the

stimuli became summated he also reacted with the right hand. The same holds true, when the left foot is tickled,—on slight stimulations the left foot is drawn away; on continuous stimulation the left hand also reacts.

Reactions of four months' old babies and older conformed more and more to Pflüger's law.

Experiments on older children give results of like character. Boy of seven years; when asleep, I tickled the right side of his face, the right hand reacted; I tickled the left side of face, the left hand reacted. When stimulations became summated or more intense the whole trunk reacted.

In experimenting on adults in the waking state the results are somewhat uncertain. When touching with a moistened object the right side of the face, for instance, and telling the subject to wipe it *immediately*, sometimes the right and sometimes the left hand reacts. One can often observe that when the subject reacts to sudden irritations, Pflüger's law holds good; and sometimes one can observe reactions with the left hand alone, the *right hand being used for more differentiated movements*.

Experiments on adults, when asleep, almost entirely conform to Pflüger's law of reflex reaction, especially is this the case, if the person has his hands free or sleeps on his back. In cases of hypnoidal states and hypnosis, the reactions are somewhat modified. If the subject happens to react with one hand, he *keeps on reacting with it*. Thus when the subject is touched with a wet or irritating object and is told to react, if it be the right side, he reacts with the right hand. If now the left side is similarly stimulated, the *right hand keeps on reacting crossing to the left side*. This takes place, even if there is a long series of stimulations,—the subject reacts with the hand he first starts to react with. This may be largely due to subconscious suggestion. I observed similar reactions in very suggestible subjects even in the waking state.

Manaceine's results are not confirmed by me, and still there is some truth in Manaceine's statement. Now I have suspected for some time that if the condition of limitation of voluntary movements is one of the important factors in the induction of sleep, we should expect that it

would not be a matter of indifference on which side we rest. We should expect that, if the right side is the more active, that the limitation of the voluntary movements would be more marked on that side. In carrying on my experiments on children and adults when in their sleeping states, I had occasion to observe that there was a *definite course in the motor reactions*, in the process of falling asleep. There is method in sleep. Some people go to sleep only on their back and find it difficult to fall asleep otherwise, while others who go to sleep on their side and who form the greater majority *always go to sleep on the same side*; there are very few who can fall asleep indifferently on either side. Moreover, my observations have shown me that by far the majority of *right-handed people go to sleep on the right side, while left-handed people go to sleep on their left side*. I further verified this interesting fact by statistical inquiry among my patients as well as among Harvard students. Some of the right-handed people who go to sleep on the right side may after some time turn to their left to change position, while others keep on sleeping on the same side through the whole night. The majority change position, the right-handed to the left and the left-handed to the right. More than *seventy-five per cent of right-handed people* have given records to the effect that they sleep on the *right side* or rather *fall asleep* on that side. Of the left-handed persons, I find only *one out of ten* who falls asleep on the right side. One case is specially interesting to quote: "Up to my seventh year I slept on my right side and I was right-handed. At about the age of seven I met with an accident, I was run over by a team and my right side was injured so that I could not use the limbs of the right side. I used my left hand only, I began to sleep on my left side. This I did up to my fifteenth year. I then began to practice with my right hand too and am now ambidextrous. I sleep now on either side. I use both hands."

Some claim that the reason why they sleep on the right side is because of the dreams produced by the pressure of the heart when sleeping on the left; but it is interesting to observe that others sleeping on their left write in their accounts that they cannot sleep on their right side, because

of the bad dreams, restlessness and nightmares produced. Some claim that it is simply a matter of habit, but the very fact that habit should give such a large percentage of right-handed, sleeping on the right side, and of left-handed, sleeping on their left side, points to a fundamental condition of our functioning activities and mode of rest. This condition we have found in the variability of our consciousness, by the ceaseless variations in the incoming sensory impressions from sense organs, muscles, joints and various other organic activities. The kinæsthetic sensations coming from the motor activities of our muscles and joints are very important in this respect. To bring about a state of rest and sleep, we must have the condition of monotony and limitation of voluntary movements, hence we can well see the reason why right-handed people whose right side is more active limit that side, while left-handed people prefer to limit the left side most active in them. We can also partly account for Manaceine's results, namely that in sleep, right-handed people react with their left hand which remains free as they sleep on the right side, while left-handed people react in sleep with their right hand, because they sleep on their left side. Manaceine's explanation is pointless in referring these reactions to the activities of the right and left hemispheres. According to Manaceine the left hemisphere of right-handed people is fatigued and hence in sleep the right hemisphere and the left hand have the ascendancy. This explanation is fanciful and does not agree with facts. For when the experiments are carried out rigorously, *the reactions are found to conform, with the exception of very young infants, to Pflüger's law.*

Thus the experiments and observations made on lower animals, infants, children and adults all point in one direction, they point to the fundamental conditions of sleep, to monotony and limitation of voluntary movements. Taking as my motto the dictum "*hypotheses non fingo*" I strictly followed the logic of facts. Sleep is not so much due to merely cutting off sensory impressions, be they intense or faint, as to the *monotony* of sensory impressions which in fact may even be intense and numerous. It is the invariability of sensory impressions that reduces the organism to the passive state which we experience as sleep.

PART II
THEORETICAL
CHAPTER XII

Cell-energy, Threshold, "Stimulus-exhaustion" and Sleep

WE may advance in a tentative way the following theory which should be regarded as a provisional hypothesis of the causation of sleep.

One of the main characteristics of living protoplasm is its adaptability to the conditions of the external environment. External stimuli give rise to reactions of adjustment on the part of living substance. This property known in physiology as irritability is specially characteristic of all living matter or of what Huxley so aptly describes as "the physical basis of life." Verworn defines the irritability of living substance "as its capacity of reacting to changes in its environment by changes in the equilibrium of its matter and its energy." In other words, living tissue responds to external stimulations with some discharge of energy. The form of the discharge depends on the peculiar protoplasmic structure, according as it is muscle, gland, nerve cell or but slightly differentiated protoplasm, such as amoeba or bacterium. The character of the reaction to stimuli depends on the state of organization of the living tissue.

The delicacy of response of living matter to external stimuli has its limit. Very weak stimulations do not call forth any reaction. Living tissue can only be set into activity by stimuli of certain intensity. If the stimulus falls below that intensity, the protoplasm does not react. This holds true of all cells, from the simplest bacterium and infusorium to the most highly differentiated muscle-cell or neuron. The minimal intensity below which the stimulus remains ineffective is regarded as the threshold of stimulation. As Howell puts it: "A stimulus too weak to give a response with a motor nerve is usually designated in physiology as subliminal; a similar stimulus with sensory nerves is frequently expressed by the equivalent term of subliminal, that is below the threshold, so a stimulus just strong enough to provoke a perceptible reaction is the

minimal stimulus for efferent nerves and the threshold stimulus for sensory nerves." "Exceedingly feeble stimuli" says Landois "are without effect. The degree of intensity of stimulation that originates the first trace of sensation is called the threshold of sensation or the threshold value." The same is more clearly put by Verworn:

"Let us imagine an organism or part of an organism, *e.g.*, a muscle, under conditions in which no stimulus affects it, and let us bring to bear upon it a stimulus, *e.g.*, the galvanic current, which varies in intensity from zero upward and can be graded easily and delicately. Then we should expect the muscle to exhibit phenomena of stimulation, *i.e.*, to perform a contraction, as soon as the intensity is increased above 0. But this does not happen. The intensity can be increased considerably before the muscle performs even the slightest twitch. Only when the intensity has reached a certain degree does the muscle respond with a contraction; from here on the contraction is never wanting, and up to a certain degree becomes more energetic the more the intensity is increased. The stimulus, therefore, begins to operate only at a certain intensity, and this point is termed the *threshold of stimulation*. Below the threshold the stimulus is ineffective; above it the effect increases with the increasing intensity of stimulus. For the different forms of living substance the value of the threshold is very different. Thus, nerve-fibres are put into activity by extremely feeble galvanic stimuli, while *Amæba* demands very strong currents. The same is true of all other varieties of stimuli in relation to the various forms of living substance."

Psychologically we may agree with Stout that "the point at which it (the stimulus) is just indistinguishable,—so that the least increase would make it distinguishable is called stimulus-threshold."

Kulpe's definition is short; "the just noticeable stimulus is technically termed the stimulus-threshold (*die Reizschwelle*)."

With the increase of stimulations the irritability of the living substance diminishes, the threshold rises. The same minimal stimulus will no longer bring about a reaction, the stimulus must be increased in intensity before any effect

can take place. Perhaps the best account of it is given by Verworn in his "General Physiology."

"If a living object be stimulated by long-continued, oft-repeated, or very strong stimuli, after some time it passes into the condition of fatigue. The general characteristic of fatigue is a gradual decrease of the irritability of the living substance. This is expressed especially in the fact that with increasing fatigue, the intensity of the stimulus remaining the same, the result of the stimulation becomes constantly less.

"We have already become acquainted with some examples of this fact in considering galvanic stimulation. If a constant current of average strength be passed through an *Actinosphaerium*, at the moment of making, there begin to appear at the anode marked phenomena of contraction. The protoplasm of the pseudopodia flows centripetally until the latter are drawn in. Then the walls of the vacuoles break; and a granular disintegration of the protoplasm results, which proceeds constantly farther from the kathode during the passage of the current. This disintegration, beginning with great energy, becomes slower and less extensive the longer the current flows, and after some time is at a complete standstill. This means that the living substance of the *Actinosphaerium* becomes fatigued in the course of continual stimulation, and decreases in irritability; hence the stimulus, which at first induced pronounced phenomena of disintegration, later produces no reaction at all. *Pelomyxa* is fatigued still more rapidly than *Actinosphaerium*. Stimulation for a few seconds is sufficient to make individuals of this genus wholly non-irritable to currents of equal intensity; a much greater intensity is then required to call out the same reaction."

The principle of variability of stimulation is of great importance in the reaction of nerve-tissue. When the stimulus remains invariable, both in intensity and quality, no reaction follows. This is clearly brought out in experiments on nerve-tissue. "The electrical current" says Landois,¹ "exerts its strongest irritant effects upon a nerve at the time of its entrance into the nerve and at the time

¹ Text-book of Physiology.

of its disappearance. In like manner any rapid increase or decrease of the current passing through a nerve has a strong irritant effect. If on the other hand, the current be allowed to pass gradually into the nerve trunk or to disappear, or the current passing through the nerve be gradually increased or diminished, the visible signs of nerve irritation are much less marked. In general, the stimulation is most pronounced the more rapid the current-variation within the nerve, that is the more suddenly the strength of the current passing through the nerve is increased or diminished." This holds true in the case of the nerve, in which as Bowditch has shown there is little or no fatigue; where fatigue is present the principle of variability becomes a factor of the utmost consequence. The principle of variability of stimulation plays a very important rôle in cells in general, and in nerve-cells in particular where fatigue easily sets in and the threshold is raised with the continuation of stimulation and with the successive discharges of cell energy. Variability of stimulation and fatigue influences the fluctuations of thresholds.

It may be well to bring here the recent work of Sherrington in regard to the relation of neuron threshold and the discharge of neuron energy. Sherrington points out that among the characteristic differences between conduction in nerve-trunks and in reflex arcs (where nerve-cells are interpolated, we find in the latter "Irreversibility of direction instead of reversibility as in nerve-trunks, fatigability in contrast with the comparative unfatigability of nerve-trunks," and "much greater variability of the threshold of stimulus than in nerve-trunks." Sherrington points out that "in certain cases, especially in Invertebrata, observation (Apathy, Bethe, etc.), indicates that many nerve cells are actually continuous one with another. It is noteworthy that in several of these cases the irreversibility of direction of conduction which is characteristic of spinal reflex arcs is not demonstrable; thus the nerve net in some cases *e.g.* Medusa, exhibits reversible conduction (Romanes, Nagel, Bethe and others). But in the neuron-chains of the gray centred system of vertebrates histology on the whole furnishes evidence that a surface of separation does

exist between neuron and neuron. And the evidence of Wallerian secondary degeneration is clear in showing that that process observes strictly a boundary between neuron and neuron and does not transgress it. It seems therefore likely that the nexus between neuron and neuron in the reflex arc, at least in the spinal arc of the vertebræ involves a surface of separation between neuron and neuron." "In view therefore of the probable importance physiologically of this mode of nexus between neuron and neuron it is convenient to have a term for it. The term introduced has been *synapse*" "At each synapse there is a neuron threshold. At each synapse a small quantity of energy, freed in transmission, acts as a releasing force to a fresh store of energy not along a homogeneous train of conducting material as in a nerve-fibre pure and simple, but across a barrier which whether lower or higher is always to some extent a barrier."¹

Assuming then the principle of variability of stimulation of neuron energy and neuron threshold in regard to the liberation of neuron energy, principles advanced in my previous works, we can return to our subject under discussion, namely sleep. We may regard *sleep as a reaction of protoplasm* and as such we may express it in terms of *neuron threshold and neuron energy liberation*.

If a series of stimulations are kept up without variation, the sensory threshold gradually rises and finally the stimuli fall out of consciousness, they fail to awaken the psycho-physiological systems which have responded to the same stimulations before, because of the rise of the thresholds. This psycho-physiological law, characteristic of all tissues and psycho-neural systems, underlies the phenomena of sleep. As the cell or the neuron keeps on reacting to stimulations the disposable physiological energy becomes lowered and there is greater economy in the liberation of cellular or of neuron-energy. The cell or neuron does not respond to the same intensity of stimulus with the same amount of energy. In other words, *the threshold rises*. Should the stimulation keep on acting without variation, both as to quantity and quality, the threshold rises so high that the stimulus can no longer step over the threshold and

¹ Sherrington, "The Integrative Action of the Nervous System."

can no longer call forth any reaction in the cell. We say that the cell or the neuron is fatigued or exhausted. Really *this is not so much due to the exhaustion of the cell as to the exhaustion of the stimulus*, — it is the stimulus that has exhausted itself. Vary the stimulus in quantity or in quality and the cell or neuron reacts once more.' We may possibly best describe this general physiological fact of cellular reaction by the term of "*stimulus-exhaustion*." By repetition the stimulus exhausts itself and can no longer call forth a reaction in the cell, although the cell may otherwise possess a large amount of disposable energy.¹

As far as the particular stimulus is concerned the cell may be regarded as asleep. The stimuli are withdrawn. Putting the same general law in different terms we may say that *sleep is produced by monotony*.

The cell or the neuron may be regarded as a reservoir of energy. With the great biologist, Strassburger, we may describe the cell as "*energid*." Conceived in terms of energy the cell possesses various levels of energy. The

¹ This may be related to the interesting experiments referred to by Sherrington:

"When the scratch-reflex elicited from a spot of skin is fatigued, the fatigue holds for that spot, but does not for the reflex as obtained from the surrounding skin. The reflex is then tired out to stimuli at that spot easily obtainable by stimulation two or more centimetres (half an inch) away. This is seen with either mechanical or electrical stimuli."

(*The Integrative Action of the Nervous System*, Sherrington)

² "The singleness of action from moment to moment thus assured is a keystone in the construction of the individual whose unity it is the specific office of the nervous system to perfect. Releasing forces acting on the brain from moment to moment shut out from activity whole regions of the nervous system, as they conversely call vast other regions into play."

"An arc under long continuous stimulation of its receptor tends even when it holds the common path, to retain its hold less well. Other arcs can then more readily dispossess it. A stimulus to a fresh arc has, in virtue of its mere freshness, a better chance of capturing the common path."

"This waning of a reflex under long-maintained excitation is one of the many phenomena that pass in physiology under the name 'fatigue.' Its place of incidence lies at the synapse. It seems a process elaborated and preserved in the selective evolution of the neural machinery. It prevents long continuous possession of a common path by any one reflex of considerable intensity. It favours the receptors taking turn about. It helps to ensure serial variety of reaction. The organism, to be successful in a million-sided environment, must in its reactions be many-sided."

(Sherrington, address British Association for the Advancement of Science, 1904).

main levels of cellular energy have been described by me in former works as *dynamic*, *static* and *organic*.

By *dynamic* energy is meant that portion of energy which the cell can dispose of in its relations and reactions to the stimuli of the external environment.

By *static* energy is designated that portion of energy which is used only for the life maintenance of the cell, both in relation to other cells and to its own molecular constitution. It is the energy requisite to keep up its morphological organization and internal physiological functions.

By *organic* energy is meant the energy contained in the tissues of the dead cell not as yet decomposed into its inorganic constituents.

The levels of cellular energy are not different in kind. They merely represent progressive phases or stages of the same process of cellular activity.

In its relations with the external environment, the cell does not utilize the whole of its dynamic energy. A large amount of it lies fallow, so to say, and remains inaccessible to the ordinary stimulations of the external environment. This amount of unused energy may be termed *stored, reserve energy*.¹ For the cell stores up energy as it stores food, in order to be able to meet the various emergencies that may arise in the course of its relations with the external world. In respect to storing up energy the law of stimulus-exhaustion may be regarded as a safeguard to the cell. The same qualitative stimulus cannot draw more than its allotted portion of energy. If the stimulation continues and attempts to draw more than its share the door, so to say, closes and the stimulus knocks in vain against a locked door. *The threshold rises with each successive stimulation*. When the *maximum* amount is drawn, there is no longer response to that particular stimulus, or rather to say, to that particular qualitative stimulation of a definite intensity. In regard to it the cell no longer reacts, — it is asleep. In order to respond again to the particular stimulus of a certain intensity and quality, the cell must recuperate its special form of energy. When this recuperation is effected, the cell is once more ready to react to the given stimulus.

¹ See my *Studies in Psychopathology*, Boston Medical and Surgical Journal, 1907.

The liberation of cellular energy in response to external stimuli may be regarded as the *active waking states*, while the restitution or passive rest-states of the cell, during which the threshold is raised and cellular recuperation takes place, may be regarded as *sleeping states*.

From the standpoint of cell metabolism, waking states are correlative with *katabolic* changes, while sleeping states are correlative with *anabolic* conditions. When the cell is active and awake, it liberates energy; when it sleeps, it stores energy. This process of storing up energy is going on all the time to some extent. If a stimulus of a certain intensity and quality keeps on liberating energy, it may arrive after some time at the *maximum*, when the threshold rises so high that the cell no longer reacts.¹ In respect to that stimulus the katabolic, active, waking state ceases and is replaced by the anabolic sleeping states. The cell may thus be awake to all other stimulations, but in regard to the special stimulus it is asleep. We may say that the cell is *asleep partially*. When the liberation of energy has reached the *maximum* point, and the thresholds are raised

¹ "Whenever by a stimulus applied to an irritable substance, the potential energy there stored up is liberated the following phenomena may be observed: (1) A so-called latent period of variable duration during which no effects of stimulation are manifest; (2) A very brief period during which the effect of stimulation reaches a maximum; (3) A period of continued stimulation during which the effect diminishes in consequence of the using up of the substance containing the potential energy—i.e., a period of fatigue; (4) A period after the stimulation has ceased in which the effect slowly passes away.

The curve drawn by a muscle in tetanic contraction, as shown in the diagram, illustrates this phenomenon. Thus, if AD represents the duration of the stimulation, AB indicates the latent period, BC the period of contraction, CD the period of fatigue under stimulation, and DE the after-effect of stimulation showing itself as a slow relaxation. When light falls upon the retina corresponding phenomena are to be observed."

An American Text-Book of Physiology, William H. Howell, Vol. II, p. 343.

in regard to most or to all the ordinary stimuli of the external environment, the cell may be said to be *fully asleep*.

Should however, the liberation of energy by various stimuli keep on and begin to liberate the reserve energy levels and then pass over into the levels of static energy, the cell falls into a *pathological waking state*.

We may represent the relation of the different levels and their correlative states by the following diagram.

What is true of the cell holds also good of systems of cells and neurons constituting a multicellular organism. The condition of monotony and the law of stimulus-exhaustion play a very important rôle in the various changes and adaptations of life-activity. *Stimuli which have exhausted themselves by their monotony drop out and are replaced by new ones until the whole round of stimuli is gone through and the organism ceases to respond to its external environment, — falls asleep.*

The rise of threshold after stimulation holds true in the whole domain of biological activities. If the gastrocnemius muscle of the frog, for instance, is stimulated, say by an electric current, the muscle with successive stimulations responds less readily with contraction and this becomes more evident with the onset of fatigue. Pffefer in a series of extremely interesting experiments has shown that spermatozoids of ferns are attracted by malic acid, the progressive response of the attraction of the cell requiring a constant increase of the degree of the concentration of the acid, the increment of stimulations bearing within certain limits a constant ratio to the total stimulus. *The threshold rises with each successive stimulation.*

In the sphere of sensation we find the same rise of threshold. We are all acquainted with the fact that an additional candle or lamp, for instance, in well lighted room does not produce the same sensory effect as when brought into a more or less dark room. An electric light in the sun is scarcely perceptible. An additional ounce to a lifted pound does not feel as heavy as when raised by itself. A sound added to another sound or to a noise sounds less loud than when appearing isolated or when the same sound is breaking silence. The same relation holds true in the case of other senses.

Organic Energy Static Energy Reserve dynamic Energy free dynamic Energy

~~Restitution of Energy~~

~~Liberation of Energy~~

~~Waking States~~

~~Sleeping States~~

Pathological Physiological

Death

Morbid
Reactions

Psychomotor
Reactions to
unusual changes
of external environment

Psychomotor
Reactions to
habitual
environment

This same truth is still more clearly brought out in the fact that if we take a certain stimulus as a unit giving rise to a definite sensation, then as we progressively ascend and add more and more units of the same stimulus, the qualitative intensity of the sensation is far from rising proportionately. If we take for instance, the weight of an ounce as our unit of stimulation, then the successive moments of unit-stimulations, that is of ounces, will not give rise to as distinct and similar sensations as did the initial stimulation. The second ounce will give a sensation fainter than the first one and the third fainter than the second and so on until a point is reached, when the sensation of an additional ounce will not at all be appreciated, will dwindle away and almost reach the zero-point. In the same way, if the pressure of a gramme is excited in the hand, successive increments of grammes will not in equal degree increase the sensory effect. The additional increments of grammes, though they are equal units of stimulation, give rise to fainter and fainter sensations, until finally all sensory appreciation of the added unit fades away and disappears. If the hand is immersed in water say at the freezing point, an addition of ten degrees will be perceptibly appreciated, while successive increments of ten degrees each will be felt less and less and finally will not be noticed and will become difficult to detect. In short the threshold rises with each successive stimulation.

To bring about a sensory response of an already stimulated sense-organ the intensity of the stimulus must be relatively increased. This is what constitutes Weber's law. The continuous progressive sensory response of a sense-organ requires a constant increase of stimulation which within certain limits, bears a constant ratio to the total stimulus.¹ This law is sometimes summed up by

¹ By noting for each strength of stimulus the addition required to evoke a just perceptible alteration of sensation, a series of quantities is obtained expressing the law according to which sensation alters when stimulation is increased. This expression is the so-called "law" of Weber. It says that a given stimulus is perceived less when added to a large than to a small one, or that an addition to a large stimulus is perceived less than an addition to a small one, unless it, relatively to the stimulus, is as great. The "law" may be phrased variously in physiological theory. It may be interpreted as not physiological at all, but psychological. The

physiologists and psychologists in the statement that the stimulus rises in a geometrical and corresponding sensation in an arithmetical progression, or as Fechner expresses it, "the sensation varies as the logarithm of the stimulus." We are probably nearer the truth, if we limit ourselves to the statement that "the increase of the stimulus necessary to produce a just perceptible difference in the sensation bears a constant ratio to the total stimulus." In short, *activity raises the threshold*. The phenomena fall under the law of stimulus-exhaustion. Each reaction to an external stimulus raises the threshold of that stimulus and with its repetition becomes finally excluded, temporarily, at least, from the cycle of *living* stimuli, that is, such as are capable of bringing about reactions and adjustments to the external environment. *The excluded stimulus is ignored, becomes non-existent for the organism. The organism is asleep for that stimulus.*

In the course of the daily activity of the individual the thresholds of sensitivity and of sensori-motor reactions in response to the stimuli of ordinary life rise gradually, the stimuli fall below the threshold of living reactions. The mass of stimuli of the external world constituting the external environment of the individual drops out of the life-cycle of the individual and for the time being ceases to exist for him. *The individual is asleep, asleep for that environment.* In other words the stimuli of the external world by their continuous action have become monotonous, have raised the thresholds and have become excluded from the life existence of the individual who is now deeply asleep for that given environment.

disproportion between increment of stimulus and increment of sensation may take place in purely psychological events and processes. Wundt is of that opinion. He points to the wide occurrence of such a ratio in all psychical activity as outcome of the relativity inherent in every conscious process. Waller finds the response in a nerve trunk directly stimulated, as judged by action current, increase much more nearly directly as the increase of external stimulus than does the response from muscle when nerve is stimulated, or from nerve when retina is adequately stimulated. Waller's evidence seems to point to the law being in part a function of the nerve-cell endings; probably, therefore, applicable to synapses as to motor plates. Delbœuf considers the law an expression of ever-increasing proportion of loss of effect in the central nervous system, due to "fatigue." (Sherrington in Schäfer's *Text-Book of Physiology*, Vol. II, p. 931.)

Should an appreciable variation occur in the environment at about the time when the organism is ready to fall asleep *the external world becomes again a living reality*. The individual becomes alive and wide awake to what is going on around him. If an appreciable change in the total mass or in some stimuli constituting the environment takes place, the individual may be awakened out of his sleep. Should the variation occur in some of the stimuli, their thresholds become lowered and the individual, though asleep in regard to all else, is awake in regard to them. This partial waking state is the soil in which dreams develop most luxuriantly. *We do not dream, when we are asleep, we dream, when we are awake.*¹

The individual may be asleep to all else and still be awake to special stimuli whose thresholds are very low. The mother watching over her babe, the nurse, the physician attending their patients may be sound asleep to loud noises, but are alive and wide awake to the slightest changes in their charges. They have lost touch with the whole external world, but the ones cared for still have a firm hold on them. High as our bridges are raised for the whole world, they are low for the ones we care for.

Let an appreciable change occur in the stimuli to which we are awake, an amount of energy immediately is drawn from the reserve store, the thresholds barring the entrance to the host of intruders are lowered and we are once more in communication with the external world, — our sleep is gone and we are wide awake. *We can thus be wide awake in the very depth of our sleep*. The possibility of shaking off the grip of sleep under appropriate circumstances and especially the highly significant fact of wakefulness in the very depths of sleep form insuperable difficulties for all those plausible, apparently scientific theories of sleep, theories based on circulation, engorgement, anemia or hyperæmia, narcosis and autointoxication. We can watch in our sleep, count the flow of time and awake at the right moment. The fact that we can sleep and still watch and keep awake in regard to special objects and particular persons shows that sleep is not a matter of blood circulation

¹ My work on dreams will be published in a separate essay.

or of intoxication by accumulation of waste products in the system. Sleep is not a disease, not a pathological process due to the accumulation of toxic products in the brain or in the system generally. Sleep is not of those fatal uncontrollable onsets characteristic of the morbid processes, such as epilepsy, sunstroke, or apoplexy, nor is sleep a sort of narcolepsy. Sleep is not a disease, it is not, as the chemical speculators would have it, a kind of narcosis of the system by the poisons of fatigue products. *Sleep is not pathological, it is essentially physiological in character.* We do not go to sleep, because we are poisoned and wake up when the poisons are eliminated. We go to sleep at our own sweet will and can wake at pleasure. We can wake at any moment and can even be awake in the very depths of our sleep. Sleep is not an abnormal condition, it is a normal state. Like the waking states, *sleep-states are part and parcel of the life-existence of the individual.* Waking and sleeping are intimately related, — they are two different manifestations of one and the same life-process, — one is as normal and healthy as the other. One cannot help agreeing with Clapèrede's biological view that sleep is a positive function of the organism, that sleep belongs to the fundamental instincts. As Clapèrede forcibly puts it: "Le sommeil est une fonction de défense, un instinct qui a pour but, en frappant l'animal d'inertie, de l'empêcher de parvenir au stade d'épuisement: ce n'est pas parce que nous sommes intoxiqués, ou épuisés, que nous dormons, mais nous dormons pour ne pas l'être."

If we look at the matter from a psychological standpoint, we may say that sleep is a *rise of the thresholds of mental aggregates or of moments consciousness.* I have pointed out in a former work¹ of mine that to minimize the expenditure of neuron energy and reach the minimum of consciousness constitutes the tendency of psychomotor life. We can fully realize the importance of this tendency, if we regard it from a teleological point of view. In the struggle for existence or in the economical system of competition of modern life the saving of unnecessary expenditure, where only possible, is of the highest consequence. Organisms

¹ Multiple personality.

that can best effect such an economy of energy will be best fitted to survive. Those organisms that are enabled to reduce to its *minimum* the friction and loss of neuron energy have the advantage of possessing at their disposal a greater amount of energy to cope with new circumstances, with novel conditions, and react better and in a more favorable way when confronted with changes in their environment. This economizing becomes absolutely indispensable in the life-existence of higher organisms, the environment of which is highly complex. *The reduction of psychomotor activity to the least amount of psychophysiological expenditure, in other words, to the minimum of consciousness is the law of psychomotor life in general and of the highest representatives of that life in particular.*

I have further pointed out in the same work that this economizing is by no means an endless process, there is a certain minimum of consciousness beyond which psychic states cannot pass. This minimum of consciousness once reached, must remain stationary, for a fall below it is the arrest of the activity of the mental aggregate. In other words, there is a certain minimum below which consciousness cannot be reduced with impunity. Reduce the consciousness of the total psychic aggregate by lowering the sensibility of its constituents and the whole mental system will cease to function. Now under the conditions of monotony and limitation of psychomotor activity the *moment-threshold rises until the psychic minimum vanishes and the organism is asleep.*

Putting the matter again in teleological terms we may say that we go to sleep when we relinquish our hold on the relations of our external environment. We fall asleep when our consciousness is fagged, when we wish no longer to enter into communication with the external world, when we lose interest in our surroundings. When our interests in external existence fags and fades away we go to sleep. When our interests in the external world cease we draw up the bridges, so to say, interrupt all external communication, as far as it is possible, and become isolated in our own fortress and repair to our own world of organic activity and inner dream life. We fall asleep when the vital interests

in external being have fallen into the background of consciousness; we awake when those interests are aroused. When the struggle for existence ceases we repair to our castle and battlements. Sleep is the interruption of our intercourse with the external world; — it is the laying down of our arms for a respite in the struggle of life. Sleep is a truce with the world. When all psychomotor reactions to the stimuli of the external environment cease, we sleep. We sleep, because we are no longer interested to take an active part in the battle of life. From a teleological standpoint we may say that sleep is a dismissal of the external world with all its vicissitudes, troubles and pains. We cease to desire, we cease to react, and we sleep and dream in peace.

And when evening descended from heaven above,
And the earth was all rest, and the air was all love,
And delight, tho' less bright, was far more deep
And the day's veil fell from the world of sleep,

And the beasts, and the birds, and the insects were drowned
In an ocean of dreams without a sound;
Whose waves never mark, tho' they ever impress
The light sand which paves it, consciousness.

—*Shelley.*

CHAPTER XIII

Motor Consciousness and Sleep

THE condition of limitation of voluntary movements plays an important rôle in the induction of sleep states. This is due to the fact that motor consciousness not only forms the main body of our mental activity, but also that ideo-motor life is more subject to changes from slight stimulations than is our purely sensory life. Motor elements are highly plastic and modifiable; they enter readily into ever new combinations. From a biological standpoint one can realize the importance of the great modifiability displayed by sensori-motor and ideo-motor elements, since in the adaptation of the organism to its environment it is these elements that are mainly employed in reaction to stimuli of the external world. From the standpoint of adaptation a slight difference of sensory experience may give a widely different and highly complex motor reaction.

Psychomotor processes form the most important and largest portion of mental life. With the exception of man, all the representatives of the animal kingdom, from the lowest to the highest forms, represent but different stages in the evolution of sensori-motor life. The great majority of mankind still leads a life closely allied to animal sensori-motor states. Instance the delight of children in their plays, and the all absorbing interest of college students in their baseball and football games. Even in the highest and most developed forms of mental activity motor ideas and representations are by far the most predominant. Without motor elements ideational life is arrested. It is these sensori-motor and ideo-motor elements that constitute the "stream, the flow, the current" of our mental life. Motor elements enter freely and easily into combinations with all other elements of mental life.

This ease and high plasticity of ideo-motor elements is specially well brought out in hypnosis. Sensori-motor and ideo-motor suggestions are taken before purely sensory suggestions; paralysis, catalepsy, contracture, all kinds of

motor and kinæsthetic illusions and hallucinations are easily induced even in the very light stages, while it is only in the deeper stages that changes of ideo-sensory elements can be brought about. The induction of purely ideo-sensory illusions, hallucinations or delusions, positive and negative, can only be effected in the very deep stages of hypnosis. The freedom in forming new combinations and associations makes the suggestion of motor ideas and representations highly effective.

Throughout the scale of animal life, from the lowest to the highest forms, intelligence is intimately related to the degree of development of the muscular system and to the delicacy of motor adjustments. Among the lower forms of life the Cephalopods are well equipped with powerful muscular arms capable of executing a great variety of vigorous movements. Now the Cephalopods also possess a more highly developed nervous system with a higher grade of mental functions than the rest of the Mollusca. The great activity of ants and bees is notorious and their instinctive psychic life is the richest among the Arthropoda. Note the great variety of motor adjustments of the beaver and also of the intelligence that goes along with it. Birds possessed of a high degree of activity and motor adaptability are also the most intelligent of their kind, such for instance as the different species of talking birds. Notice the activity and great agility of the fox and also the unusual cunning for which the animal is so celebrated in song and fable. The suppleness of the dog, his quick reactions to stimulations, the resources of his motor adjustments and the great extent of his modifiability under changing conditions are all well known and along with them goes a high degree of mental activity. Of all the Mammals the Quadrumana are the most active, the most imitative, and with the exception of man, they are also the most intelligent. When we come to man we cannot help admiring the high complexity and extreme delicacy of his motor adjustments. Most marvellous however is the human hand, that divine organ which gives shape and form to works of art, to all outward visible manifestations of civilization. The great artists and thinkers of antiquity held the human hand in

great veneration. One of the great Greek philosophers did not even hesitate to declare that man's superiority over the brute was due to his hand. Finally in the wonderfully delicate motor adjustments of speech we find clearly illustrated the intimate relation between motor and psychic activities.

Experiments prove the same truth of the importance of motor elements in our mental life. If a series of syllables or numbers is given to memorize, after one reading, five out of ten can be remembered, though with some difficulty; but if the syllables or numbers are written down at the same time, though not looked at during the writing, a far greater percentage such as six or seven syllables can be remembered. If the motor elements in a train of ideas are suppressed, the order of the series becomes confused and even totally destroyed, showing that the motor ideas are important links in trains of association of ideas. *Motor elements form the nucleus of consciousness.*

Biologically regarded, voluntary activity, will, is the power of the organism to adjust itself to the conditions of the external environment. In its last psychological analysis voluntary activity, will, consists of ideo-motor elements, of various modes of adaptations. Will, consisting of kinæsthetic elements, constitutes the active subject of personality and individuality. If this be realized, then the vital importance of motor consciousness cannot be too highly overrated. *Motor consciousness is at the very heart of personality. We are what we can accomplish.* Extreme variability and adaptability of reactions to environment are the main characteristic traits of intelligence, will, personality.

The readiness of psycho-motor elements and group to enter into ever new combinations gives rise to the formation of a great wealth of associations which help to make the labile psycho-motor groups and systems easy of recall. In fact it may be said that *the ease of recall is proportionate to the mass of associated kinæsthetic elements.* The great modifiability and variability of systems of motor elements requisite in the adaptation of the organism to the varying conditions of life, to its environment, make the ever greater

instability of motor elements an imperative necessity in the struggle for existence.

Forming the predominant elements, both as to intensity and mass, of the most complex, relatively stable, though ceaselessly shifting groups and systems, constituting the highly developed organization of mental life, the motor elements, presentative and representative, are also the first to become involved in the process of dissociation. In the various forms of nervous and mental diseases, under different conditions of intoxication and auto-intoxication, in the traumas caused by shock, physical and psychic, the delicate movements of adjustment are the first to become affected; dissociations of systems of motor elements are first to occur.

The instability of psycho-motor elements may be brought in relation with the fact of the early affection of muscular and kinæsthetic sensitivity and with the predominance of sensori-motor over purely sensory symptoms, so frequently occurring in the course of nervous diseases. With this may be correlated the significant fact referred to by Mosso that "all substances which slowly destroy the organism must produce phenomena analogous to those of curari since the motor nerves. . . . have less vitality than the sensory." It may also be observed in passing that cellular kinoplasm with the "kinocentrum," the centrosome and its archoplasmic structures, possibly the most primitive organoids of the cell, similarly manifest a high degree of variability and instability.

Motor elements may be regarded as the labile constituents of consciousness; — they become easily and frequently dissociated and dropped into the subconscious; but for that very reason they are also very easily reproduced or regenerated. In this respect motor elements follow the general biological law of organic regeneration: Organs that are easily and frequently lost in the struggle for existence are also easily regenerated, as for instance, the legs and claws of Crustacea or the tentacles of the starfish and the octopus. Dissociated systems of motor elements often become regenerated and under pathological conditions, when synthesis is impossible, they may recur with great persistence giving rise to the most uncontrollable types of

insistent ideas and impulses and to various forms of so-called "psychic epilepsy," especially of the motor type, closely mimicking typical organic epilepsy. Such dissociated, subconscious systems, like rudimentary, aborted organs, the appendix, for instance, may often prove very injurious to the organism. The recurrence of such subconscious, submerged, dissociated systems has its parallel in the biological phenomena of reversion or atavism.

The fact that psycho-motor elements enter easily into combinations and form extensive associative systems, makes them easy of recall and hence persistent in memory. From an educational standpoint one realizes the importance of this fact. Children learn things best not by abstract notions, not by looking at objects and hearing of things, but by acting out whatever is taught them. Motor consciousness is more vital than sensory. In the training of the mentally defective the best method followed is that of motor instruction, — to get at the meaning of things by means of action, even if it be automatic at first. As a matter of fact even the normal and well balanced mind gets at the meaning of things by handling them, by having the attributes and qualities of the objects and processes to be learned acted out. *Acting forms the greater part of man's life.*

Forming as motor consciousness does the very heart of mental activity, we can well realize the paramount importance of the condition of limitation of voluntary movements. By limiting the motor activity of the organism we impoverish its mental processes and lower the heart-beat of mental life. The active nucleus of psycho-motor reactions becomes passive, the organism becomes disabled in its response to the stimulations of the external environment, the thresholds rise and the organism is no longer in relation with the external world. When motor activity with its concomitant motor-consciousness becomes lowered, restricted, and fades away, the organism becomes necessarily passive and passes into sleep.

Thus monotony and limitation of voluntary movements work in one direction, — they tend to raise the thresholds of psychomotor reactions, they coöperate in the induction of sleep. These conditions are usually brought about natu-

rally in the course of the daily or nightly life activity of the individual organism, or the conditions may be produced artificially. In both cases the result is the same, — the organism falls asleep. In the life of higher animals the two cases may often combine. When the individual has fagged out his life interests in his active relations with his surroundings, when the stimuli have become monotonous to him and his activity with its correlative motor consciousness has become limited and lowered, he makes artificial arrangements for the intensification of the conditions of monotony and limitation of voluntary activity. He seeks for a dark nook, closes his eyes to exclude as much as possible all extraneous disturbing stimulations and tries to lie down quietly and comfortably, restricts his voluntary movements, breaks his connection with the external world and goes to sleep. *The organism falls into sleep, when the thresholds rise, wakes and rises when the thresholds fall.*

Looked at from a purely physiological standpoint and expressed in terms of energy, of the "cell-energy," sleep may be regarded as the onset of fatigue, as the onset of exhaustion of the levels of dynamic energy in response to external stimuli. The law of stimulus-exhaustion comes here into play as we have indicated in our discussion of cell-energy. Each particular stimulus has its maximum amount of energy which can be drawn upon, under ordinary conditions of daily life. As the special stimulus approaches its limit, it works under greater and greater difficulties, draws less and less energy and finally ceases to awaken any response,—the threshold is raised to its maximum and the organism, as far as that special stimulus is concerned, is no longer awake,—the organism is asleep. In the course of its daily activity the same takes place in regard to most of the objects, to most or to all of the stimuli that constitute the external world of the organism. The stimuli of the external world have drawn all that was permitted to them on their bank accounts, so to say, and the account for the time being is closed. Nothing more is permitted to go out. No stimulus of ordinary life is permitted to draw over and above a certain amount. There must always be ready capital for unusual situations, for emergencies. When the stimuli have drawn their due

and the organism is left with its reserve energy, — liberation of energy with its accompanying waking states ceases. The organism is no longer awake to the stimuli and is asleep.

As in the waking states the katabolic processes predominate, so in sleep the reverse processes, the anabolic, take the upper hand. The organism begins to recuperate its losses and fills up the accounts drawn upon by the stimuli of the external environment, when in active relation with them. With the increase of the income of energy the raised thresholds begin to fall until a point is reached when the stimuli once more overstep the lowered thresholds and once more gain access to the stores of life-energy, — the organism awakes and enters into active relations with the external environment.

Regarded then from various standpoints, sleep is a rise of moments-thresholds under conditions of monotony and limitation of voluntary movements. In this respect sleep strongly contrasts with hypnosis. In hypnosis the individual is specially accessible to any kind of suggestions coming from the external world, the psycho-motor reactions are greatly lightened and are released by the suggestion or external stimulus with great facility, far greater than in the waking state. This great facility is often expressed by the statement that in hypnosis the inhibitions are removed. What specially characterizes *hypnosis* is the *fact of fall of thresholds* present in individuals, with a predisposition to states of dissociation; in sleep, on the contrary we have found from our study, the general characteristic is the *rise of psycho-motor thresholds*. In passing from the waking state into the subwaking hypnoidal state the individual may either pass into hypnosis with its dissociated states and lowered psycho-motor thresholds or may go into sleep with raised psycho-motor thresholds. *The process of redistribution of thresholds takes place in the intermediary, hypnoidal states. When the redistribution of thresholds in the hypnoidal states brings about a fall of thresholds due to predisposition to and further cultivation of dissociations, the result is hypnosis; when the redistribution in the hypnoidal states brings about a rise of thresholds, the result is sleep.*

Biologically regarded, sleep is as much an instinct as hunger or sex.

Phylogenetically and ontogenetically, the sleep-states of higher animals are developed out of undifferentiated, intermediary, subwaking, hypnoidal-like states found in the resting states of the lower representatives of animal life. The hypnoidal state is the primitive rest state out of which sleep arises. Briefly put, the hypnoidal state is the germ of sleep.

Physiologically and psychologically regarded, sleep is an actively induced passive state in relation to the external environment; the psycho-physiological systems have their thresholds raised in relation to external stimulations; the rise of threshold is induced by a mass of impressions possessing little or no variability, by limitation or by relative withdrawal of stimulations, or what is the same, by monotony of stimulations and by limitation of voluntary movements.

BIBLIOGRAPHY

- ABERCROMBIE, Inquiries.
- ACHASHEFF, K., Kasuistik Narkolepsii. Med. Sbornik, Warsaw. 1896.
- ACKERRMAN, Versuch einer Physischen Darstellung der Lebenskräfte.
- BALDWIN, Senses and Intellect.
- BALLET, Sommeil Pathologique, Rev. de Med. 1882.
- BARTHEZ, Nouveaux Elements de la Science de l'Homme.
- BAUCHARD, Lectures on auto intoxication in Disease.
- BENJAMIN, Über den Schlaf Allg. Z.f. Psychiatrie. 1898.
- BERGER & Loewy, Etat des yeux pendant le sommeil, J. de l'Anat. 1898.
- Etat des yeux pendant le sommeil, Int. des B n. 1902.
- BERGSON, Le Rêve, Bull. Inst. Psych. int. Mai. 1901.
- BERNARD, Cl. La Science Experim. 1878.
- BIGELOW, Mystery of Sleep, Harper. 1897.
- BINNS, Anatomy of Sleep. 1843.
- BINZ, Ueber den Traum. 1878.
- BLODET, Observation de sommeil lethargique. Gaz. hebdomadaire, October, 1864.
- BLODGETT, A. N., A Case of Narcolepsy. Am. J. of Med. Sc. August, 1906.
- BOIRAC, La Veille continuee dans le sommeil.
- BORGHIERINI, Schlafählicher Zustand bei Thieren, Neur. Centrbl. 1891.
- BOUCHARD, Leçons sur les auto intoxications. 1887.
- BREISACKER, Zur Physiologie des Schlafes Du Bois's Arch. 1891.
- BREUER UND FREUD, Studien ueber Hysterie.
- BRODIE, BENJAMIN, Psychological Inquiries.
- BRODMAN, Plethysmographische Studien am Menschen, J.f.Ps. & Neur. 1902.
- BROWN, Compound Fractures of Cranium, Am. J. Med. 1860.
- BROWN SEQUARD, Leçons sur les nerfs vaso-moteurs. 1872.
- Le sommeil, Arch. de Phys. 1889.
- BRUNELLI, Il letargo. Riv. ital di Sc. nat. 1903.
- BURDACH Physiologie.
- CAMP, CARL D., Morbid Sleepiness, The J. of Abnormal Psych. May, 1907.

- CAMUSET, Apropos de la narcolepsie, *Gaz. de Hop.* 1880.
- CAPPIE, J., Causation of Sleep.
The Intracranial Circulation.
- CARPENTER, Physiology.
- CHARCOT, M., Attaque du sommeil, *Gaz. de Hop.* 1888.
- CHAVIGNY, M., Narcolepsie, *Soc. Med. de Hop. de Lyon.* 1904.
- CHESLEY, Does the Mind Ever Sleep?
- CLAPEREDÉ, ED., Esquisse d'une Théorie Biologique du Sommeil. *Archives de Psychologie*, February, March, 1904.
- CORNING, Brain Rest. N. Y. 1885.
- COUSINS, WARD, A Case of Profound Sleep, *Med. Times.* April 18, 1863.
- COX, Sleep and Dream.
- CZERMAK, J. N., Nachweis echter hypnotischer Erscheinungen bei Thieren.
- CZERNY, Z. Zur., u. c. entniss des phys. Schlafes, *Jahrb. f. Kindheilk.* 1896.
- DADDI Sulle alterazioni degli el. d. sys. ner. nell'insonnia experim. *Riv. di Pat. ner. e ment.* III.
- DANA, On Morbid Drowsiness, *The J. of Ner. Men. Dis.* 1884.
- DANILEWSKY, Ueber die Hemmungen, *Pflüger's Archiv Bd. XXIV.* 1881.
Recherches physiologique sur l'hypnotisme des animaux.
Beobachtungen und Versuchen über hypnotisierten Zust. bei Thieren, *Pflüger's Archiv.* 1873.
- DE BEER, Der Schlaf der Fische, *N. Wien. Tag.* 1895.
- DEBOECK & VERHOEGEN, Contr. à l'étude de la circ. cerebr., *J. de Med. Bruxelles.* 1890.
- DELBOEF, Le Sommeil et les rêves.
- DEMOOR, La plasticité des neurones et le mécanisme du sommeil, *Bull. de la Soc. d'Anthr. de Bruxelles*, XV. 1896.
- DE SANCTIS & NEYROZ, Exp. Invest. conc. the depth of sleep, *Ps R.* 1902.
Die Träume.
- DEVAUX, E., Théorie osmotique du sommeil, *Arch. Gen. de Med.* 1906.
- DICKSON, Essays on Sleep.

- DIDRICHSON, V. K., K voprosu o nervnoi bessonnitsi.
- DRSDOW, V. J., Morbus hypnoticus, Arch. f. Psych. u. Nervenkrankheiten, Bd. 13.
- DUBOIS, R., Etude sur le mécanisme de la thermogénèse et du sommeil Ann. de l'Univ. de Lyon XXV. 1896.
Le centre du sommeil, C. R. Soc. Biol. 1901.
Sommeil naturel par autonarcose carbonique, C. R. S. Biol. 1901.
- DUGAS, Le Sommeil.
- DURHAM, The Physiology of Sleep, Guy's Hosp. Rep. 1860.
- DUVAL, Théorie histologique du sommeil, Soc. de Biol. 1895.
- EDWARDS, J. Report of a case of Somnolency, Lancet V. I. 1884.
- ERRERA, Le mécanisme du sommeil, Bull. Soc. Anthr. Bruxelles. 1895.
Pourquoi dormons nous? Rev. sc. July, 1887.
Théorie toxique du sommeil, Soc. de biol. 1891.
- EWEN, A Case of Narcolepsy, Bost. Med. and Sur. J. June, 1893.
- FECHNER, Elemente der Psychophysik.
- FERE, CH., Le Sommeil paraxystique, Semaine Medicale. 1893.
Crises de courbature, Annales Med. Psych. 1888.
Note sur l'apathie epileptique, Rev. de Med. 1891.
Note sur la narcolepsie epileptique, Rev. de Med. 1898.
- FISCHER, F., Epileptoid Schlafzustande, Arch. f Psych u Nervkr. Bd. 8.
- FLEMING, Note on the Induction of Sleep, Br. & For. Med. Rev.
- DE FLEURY, Grands symptômes neurasthéniques.
- FLINT, Text Book of Physiology.
- FOOT, A. W., Narcolepsy, Dublin J. of Med. Sc. V. 82.
- FOREL, Der Hypnotismus.
Observations sur le sommeil du loir, Rev. de l'Hyp. 1887.
- FORNERIS, Du Role de la Glande thyroide, Gazette Sarda. 1858
- FOSTER, A New Standpoint in Sleep Theories Am., J. of Psych. 12. 1900.
- FRANCOSCHI, F., Il sonno patologico, Riv. di Pat. ner. e mentale V. 9
- FRANCK, Recherches sur les mouvements du cerveau. J. de l'anat. 1877.

- FREUD, Ueber den Traum.
- FRIEDMAN, M. Ueber die nichtepiletischen Absenzen, Deut. Arch. f. Nervenheilk. 1906.
- GAIRDNER, British Med. Jour. Oct. 30, 1875.
- GALTON, Inquiries.
- GAY, J., Sleeping Girls. Lancet. July 3, 1880.
- GELIEAU, De la narcolepsie, Gaz. de Hop. 1880. Las
- GILBERT, J. A., Loss of Sleep, Am. Medicine, IX, 1902.
- Gilles de la Tourette, Des attaques du sommeil hysterique.
- GIRONDEA, De la circulation cerebrale.
- HAFFNER, Schlafen u. Traumen.
- HAGEN, Psychologie und Psychiatrie.
- HAMMOND, On Sleep.
- HANSEMAN, Unters, ueber d. Winterschlaforgan, Engelm., Arch. f. Phys. 1904.
- HARTLEY, Observations on Man.
- HAYMAN, Sleeping Girl of Turville. Lancet. June 12, 1880.
- HEERWAGEN, Statistische Unters, ueber Traume und Schlaf.
- HENKE, Hypothese uber den Schlaf, Z. F. R. Med. Bd. XIV.
- HENNINGS, J. Ch. Von den Traumen.
- HEUBEL, Abhängigkeit des wachen Gehirnzustandes von ausseren Erregungch, Pflüger's Arch. XIV. 1877.
- HILDEBRANDT, DER TRAUM.
- HILL, The Physiology and Pathology of Cerebral Circulation.
- HOLLAND, HENRY, Chapters in Mental Physiology.
- HORWATH, Betr. z. Lehre uber den Winterschlaf, Verh. d. phs. medes. Würzburg. 1878.
- HOWELL, A Contribution to the Theory of Sleep, J. of Exp. Med. 1897.
- HUGHLINGS, JACKSON, Med. Times and Gazette. Dec. 6, 1862.
- JACOBY, Falling Asleep while at Work, N. Y. Med. Rec. 1890.
- JAMES, Psychology, V. I.
- JAMIESON, Edinburgh Med. Jour. July, 1871.
- KIRCHER, A, Ars magica.
- KIRCHOFF, Psychiatrie.
- Kohlschütter, Mechanik des Schlafes, Z. f. rat. Med. 1869.
Messungen der Festigkeit des Schlafes. 1862.
- KULPE, Outlines of Psychology.
- KUSSMAUL AND TEUNER, Molesch. Untersuch. 1857.
- LAJOUX, Découverte de la cause du sommeil naturel.

- LAMACK, Quelques cas de narcolepsie, *Rev. des Med.* 1897.
- LANDOIS, *Test Book of Physiology.*
- LANGLET, *Physiologie du sommeil.*
- LASEGUE, L'appetit du sommeil, *Gaz. de Hop.* January, 1882.
- LAUHSEN, Schlaf & Schlaflosigkeit, *Berl. kl. Woch.* 1897.
- LEGRAND, Un cas de narcolepsie, *France Med.* 1888.
- LEHMAN, Die Korpelichen Aeusserungen Psychischer Zustände.
- LEVIE, Somnolence, *Arch. Gen. de Med.* 1896.
- LEWES, *Physiology of Common Life.*
- LIEBAULT, Streifzuge in das Gebiet der Passiven Zustände, des Schelfes und des Träume *Z. J. Hyp.* 1893.
- LOMBARD, Varieties of the Normal Knee Jerk, *Am. J. of Ps.* 1887.
- LONEY, D. W., An Obscure Case, *Med. Record.* June, 1888.
- LORAND, Myxoedema & maladie du sommeil, *Presse Medicale.* 1905.
- MACARIO, Du sommeil.
- MACCORMACK, A Case of Narcolepsy, *Brain.* 1889.
- MACFARLANE, Dreaming.
- MACKOWITZ, V., Z. Kasuistic der Schlaftrunkenheit, *Arch. f. Krim. Anthropol.* 1903.
- MACNISH, *Philosophy of Sleep.*
- MADDEN, T. M., On Lethargy or Trance, *Dublin J. of Med. Sc.* 1881.
- MAILLARD, *Etude sur le sommeil.*
- MAKENZIE, Circulation of the Blood during Sleep, *J. of M. Sc.* 1891.
- MANACEINE, Son kak tret zhisni tsheloveka.
- Manaceine, Le sommeil.
- MARDUEL, Note sur un cas de maladie du sommeil, *Lyon Med.* 1872.
- MARSHAL HALL, *Practical Observations and Suggestions in Medicine.*
- MARTIN, L. DE ST., Influence du sommeil, *C. R. Ac. des Sc.* 1888.
- MATAS, Morbid Somnolence, *New Orleans Med. and Surg. J.* January, 1884.
- MAURY, Le sommeil.
- MAUTHNER, Pathologie u. Physiologie des Schlafes., *W. Med. Woch.* 1890.

- MAYS, Ueber die Bewegungen des Menscl, Gehirns, Virchow's Arch. Bd. 88.
- MENDEL, Ueber Anfälle von Einschlafen, Lyon Med. 1872.
- MICHELSON, Unters. über die Tiefe des Schlafes, Psych Arb. II. 1897.
- MICHELSON, Die Tiefe des Schlafes.
- MOLL, Der Hypnotismus.
- MONNINGHOFF u. Piesbergen, Messungen ueber die Tiefe des Schlafes, Z. f. Biologie. 1883.
- MOORE, C. A., On Going to Sleep.
- MORGAN, SIR T. C., Sketches of the Philosophy of Life.
- MORTIMER, GRANVILLE, Sleep and Sleeplessness.
- MOSSO, Fear.
Sulla circolazione del sangue nel cervello. 1880.
- MULLER, J. Handbuch der Physiologie.
- MUMFORD, Waking, Sleeping and Dreaming.
- MYERS, Human Personality.
- NICARD, Le sommeil normal.
- OBERSTEINER, Zur Theorie des Schlafes, Z. f. Psychiatrie. 1872.
- OPPENHEIM, Lehrbuch der Nervenkrankheiten.
- OPPENHEIMER, Physiologie des Schlafes, Du Bois' Archiv. 1902.
- PARISH, E., Ueber die Trugwahrnehmung.
- PARMENTIER, De la Forme Narcoleptique, Arch. Gen. de Med. 1891.
- PATRICK AND GILBERT, On the Effects of Loss of Sleep, Psych. Rev. III.
- PETTENKOFER U VOIT, Ueber die Kohlensäuerausscheidung und Sauerstoffaufnahme während des Wachens und Schlafens, Sitzb. d. Kngl. Ak. d. Wiss. zu München. 1866-67.
- PFLÜGER, Theorie des Schlafes, Arch. f. Phys. V. 10. 1875.
- PICTET, A., Obs. sur le sommeil chez les insects, Arch. de Ps. 1904.
- PILECZ, Ueber die psych. Erscheinungen des Schlafes, W. Med. Woch. 1891.
- PLAYFAIR, On Sleep, North. J. of Med. 1844.
- POPOV, M. N., K patologii Bessonitzi, Archiv. Psychiat. 1897.
- PORTER, K. H., Narcolepsy.
- KRAEPELIN, Psychiatrie.

- PREYER, Ueber eine Wirkung der Angst bei Thieren, C. f. Med. Wiss. 1873.
Die Cataplexie und der thierische Hypnotismus.
Ueber die Ursache des Schlafes. 1877.
- PREVOST, Sur le sommeil.
- PRINCE, MORTON, The Dissociation of Personality.
- PUPIN, Le Neurone.
- PURKINJE, Wagner's Handwörterbuch der Physiologie. III. 1846.
- RACHEL, Tissue-Metamorphosis and Sleep, Med. Record. 1883.
- RADESTOCK, Schlaf und Traum.
- RAY, Mental Hygiene.
- RAYNER, H., Sleep in Relation to Narcotics, J. M. Sc. July, 1902.
- REGIS, Mental Medicine.
- RENAUDIN, Sur l'Influence Pathologique de l'Insomnie, Annals Medico Psychologiques T. III.
- RICH, B., Ueber den Physiologischen und pathologischen Schlaf.
- RICHARDSON, The Phenomena of Sleep.
- ROBERT, Der Traum.
- ROSENBAUM, Warum müssen wir schlafen? 1892.
- ROUSSEAU, Observ. de narcolepsie, Encephale. 1881.
- RUMMO, E. FERRANINI, La circulation du sang dans le cerveau del homme pendant le sommeil.
- SAHLMEN, Ein fall von Schlafsucht, Berl. kil. Wochensch. 1881.
- SAJO. Der Schlaf der Insekten, Prometheus VII. 1896.
- SALMON, A. Sull origine del sonno. 1905.
- SCHLEICH, Die Neuroglia u. ihre Beziehung z. Schlaf, Z. f. päd. Ps. 1904.
- SCHOLZ, Sleep and Dream.
- SERGUEYEFF, Physiologie de la veille et du sommeil.
- SIDIS, B., The Psychology of Suggestion.
Psychopathological Researches.
Multiple Personality.
Studies in Psychopathology, B. Med. and S. J. March, 1907.
Hallucinations, Psych. Review. 1904.
- SHERRINGTON, The Integrative Action of the Nervous System, Rep. Br. Assoc. for the Adv. of Science. 1904.
- SIMPSON, The sleep of Molluscs, Pop. Sc. Monthl. I. 1894.

- SKERRIT AND STEWART, A Case of Potracted Sleep, *Br. Med. J.* 1898.
- SOCA, Sommeil, *Nouv. Ic. de la Salp.* 1900.
- SOLLIER, Genese et Nature de l'Hysterie. 1897.
- SOMMER, Neue Theorie des Schlafes, *Z. f. rat. Med.* 1868.
- SOUKHANOFF, S., A Case of Pathological Sleep, *The J. of Men. Patholo.* October, 1902.
- SPITTA, Schlaf und Traumbzustande.
- STEFANOWSKA, Examen du cerveau de souris endormis, *J. de N.* 1900.
- STORRING, G., Vorlesungen ueber Psychopathologie.
- STOUT, Manual of Psychology.
- STRUMPELL, Beiträge z. Theorie des Schlafes. XV.
- SZCZYPIORSKY, Plusieurs attaques de sommeil paradoxal chez un aliene, *Annales Med. Psych.* November, 1891.
- TARCHANOFF, Obs. sur le sommeil normal, *Arch. ital de Biol.* 1894.
- TCHISH, Eksperim, isledovanie vnimania vo vremia sna, *Obosr. Psych.* 1896.
- TSCHISCH. Exp. sur l'attention pendant le sommeil, *Rev. de Ps.* 1896. Tissié, Les rêves.
- VAN GIESON AND SIDIS, Neuron Energy, *Arch. of N. and Psychop.* 1898.
- VASCHIDE et Piéron, Psychologie du rêve. 1902.
L'évaluation du temps pendant le sommeil, *Int. des Biol.* I.
- VERWORN, Beitrage z. phys. des Centralnervensystem. 1898.
General Physiology.
- VOGT, Zur Kenntniss des Wesens des Hypnotismus, *Z. f. Hyp.* 1895.
- VULPIAN, Leçons sur l'appareil vas moteur.
- WERNICKE, Lehrbuch der Gehirnkrankheiten
- WEYGANDT, Beitrage z. Psych. des Traumes, *Phil. Stud.* XX.
Beitr. z. Psych. des Schlafes. Ber. ueber d. I. Kon. f. exp. Psych. Giessen. 1904.
- WUNDT, Hypnotismus und Suggestion.
Outlines of Psychology.
Grundzüge der phys. Psychologie.
- WURTZ, The Chemistry of Sleep, *Popular Sc. Mo.* 1894.
- ZBINDEN, La crainte de l'insomnie, *Arch. de Psych.* II.