This article was downloaded by: [University of Cambridge]
On: 14 January 2015, At: 01:10
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK

Annals and Magazine of Natural History: Series 2

Publication details, including instructions for
authors and subscription information:

<http://www.tandfonline.com/loi/tnah08>

Teeth of Testacellus and Glandina

John Edward Gray

Published online: 16 Dec 2009.

To cite this article: John Edward Gray (1853) Teeth of Testacellus and Glandina,
Annals and Magazine of Natural History: Series 2, 12:72, 478-478, DOI:

[10.1080/03745485709495079](https://doi.org/10.1080/03745485709495079)

To link to this article: <http://dx.doi.org/10.1080/03745485709495079>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

alea, *Cleodora*, *Tiedemannia*, and *Pneumodermon*. In all, with the exception of *Pneumodermon*, the development of which has been described by Müller (Monatsbericht der Kön. Akad. der Wiss. zu Berlin, October 1852), and by Kölliker and Gegenbaur (Zeitschrift für Zoologie, Bd. iv.), an oval embryo is formed, furnished with membranous ciliated lobes (*velum*) and a shell (even in *Firola*). In the Pteropoda this *velum* is persistent, and becomes transformed into the finlike lateral appendages of these animals. In the Heteropoda, on the contrary, it gradually disappears as the animal acquires its characteristic form. The velum of the Heteropoda and Pteropoda corresponds exactly with that of the Gasteropoda, from which it follows that the lateral lobes or fins of the Pteropoda, which are only an ulterior metamorphosis of the velum, cannot be compared with the foot of the Gasteropoda, as was Cuvier's opinion.

2. Lastly, the author has ascertained that in many Mollusca the generative organs contain both eggs and spermatozoa. The excretory canal of these organs is not double, or furnished with two semi-canals, as was supposed by Meckel, but contains at once eggs and spermatozoa: this was shown by H. Müller of Wurzburg to be the case in *Phyllirhoë*.—*Comptes Rendus*, Sept. 26, 1853, p. 493.

TEETH OF TESTACELLUS AND GLANDINA.

M. Moquin-Tandon, in the 'Journal de Conchyliologie' (ii. 125), describes the teeth of *Testacellus*, and among other particulars states that the animal has no horny jaws, a retractile proboscis, and is carnivorous.

M. Morelet (in vol. iii. pp. 27 & 257) and M. Raymond (in vol. iv. p. 14 of the same Journal) describe the animal of two species of *Glandina* from America and Africa as having nearly similar teeth, a retractile proboscis without a horny jaw, and the same carnivorous appetite. The latter author considers *Testacellus* as "a *Glandina* with a rudimentary shell." Dr. Wyman described and figured the teeth of *Glandina* in the 'Boston Journal of Natural History,' showing them to be of a conical form.

I intended, in my paper on the Teeth of Pulmonata in the last Number, to have observed, that the illustrations of that paper were kindly drawn by Mr. S. P. Woodward from the well-mounted specimens of Messrs. Cocken and Wilton. The examination of the large series of mounted specimens belonging to these gentlemen and other microscopists, has been very useful to me in these researches, as showing the uniformity and permanence of the characters afforded by the teeth, and sometimes of drawing my attention to peculiarities of form, and inducing me to examine the teeth of the animal they were said to be taken from.—JOHN EDWARD GRAY.

On the Structure of the Retina in Man. By Profs. KÖLLIKER and H. MÜLLER.

The retina is composed of different layers—viz. 1. the layer of cylinders and cones; 2. that of nucleiform bodies; 3. the layer of