SYPHILIS OF THE STOMACH

By J. W. McNEE

(From the Medical Unit, University College Hospital)

With Plates 8 and 9

Introduction.

SYPHILITIC lesions of the stomach have always been regarded as extremely uncommon, and in British medical literature the subject has attracted comparatively little attention. Even Jonathan Hutchinson, with his enormous experience, states in his text-book on syphilis, published in 1887, 'I do not know of any museum specimens, nor of any published observations (of syphilis of the stomach), but I have witnessed, in a single case, most distressing stomach pain after food in association with extensive syphilitic induration of the tongue'. A short search, however, through American, French, and German records of visceral syphilis has shown the existence of many more authentic examples of the condition than is perhaps commonly known.

In France the relation of syphilis to certain cases of gastric disease has been fully recognized, and the French view of the matter has recently been brought to notice in this country in a lecture by Monod (1921). In the discussion following this lecture, Turnbull (1921) (13) stated that in a personal experience of nearly 13,000 autopsies at the London Hospital since 1907 he had met with only one case which might possibly be regarded as syphilis of the stomach, although even in this case the proof was not completely established. This seems to agree with the experience of other large pathological laboratories in this country.

Recently the opportunity has occurred of investigating fully a very typical case of an important variety of gastric syphilis, unique, or almost unique, from the fact that abundant spirochaetes were found in the lesion. The confusing clinical signs, and the very complete pathological examination made in this case, seem to warrant its publication in full.

For comparison of this case with others of a similar kind, a brief account is also given of some of the chief records of syphilis of the stomach in this and other countries.

Historical.

It is generally recognized that the diagnosis of gastric syphilis during life has hitherto been difficult. Many more cases have been seen and described after autopsy than have been recognized and reported during life. In the earlier literature an uncertainty in the diagnosis, which was perforce based on histological evidence only, exists in many instances; and, as would be expected, the discovery of the causal organism by Schaudinn and Hoffmann (1905) and the introduction of the Wassermann reaction (1906) place all the later reports on a more definite footing.

In the older literature the pathological diagnosis of one variety of gastric syphilis is fairly certain, from the definite characters of the lesion both macroscopically and histologically. This is the condition of submucous gumma, breaking down as a rule to cause ulceration. In the case of other varieties of the disease the diagnosis, even in the presence of obvious signs of syphilis elsewhere, must always be open to doubt, since many of the histological changes found in syphilis (round-celled infiltration, endarteritis obliterans, and cellular necrosis) are found in other granulomata, and in many chronic inflammatory lesions. This point is emphasized in a case reported recently by Fowler (1921) of a woman known to have signs of gastric ulcer of long standing, but suffering from secondary syphilis when examined. A course of antisyphilitic treatment did not lead to improvement, while gastro-enterostomy afforded short relief. Finally the pyloric half of the stomach was resected, and a large callous ulcer, adherent to the pancreas, was found. In this case obviously a very chronic round ulcer of the stomach was present, having no relation to the presence of syphilis. The syphilitic infection, however, was only at the secondary stage, whereas the great bulk of the syphilitic lesions of the stomach on record have been *late* manifestations of the disease. We as yet know little of the effects on the stomach of early syphilis. Chiari (1891), in a series of autopsies on 243 cases of known syphilis, found lesions of the stomach in two cases only. In one of these the disease was inherited, and in the other acquired. Quite recently Stoeckenius (1921) has described in great detail the results of very careful and extensive histological examination of the organs from four cases of recent syphilis in young adults. In all of them the amount of granulomatous tissue found scattered through the various organs was very great, so that in three of the cases tuberculosis was at first suspected. Damage to blood-vessels was also prominent, and although the stomach is not dealt with apart from the intestinal tract generally, these observations do at least suggest the possibility of early syphilitic lesions of the stomach.

Before the discovery of the spirochaetal cause, and the use of the Wassermann test, cases of gumma of the stomach form the bulk of the literature on the subject. The most quoted early paper is that of Chiari (1891), who reported the two cases already referred to. He accepts as authentic the cases described by Klebs (1869), Cornil and Ranvier (1884), and Birch-Hirschfeld (four cases—1887). Monod

(1921) traces the French writings on the subject as far back as 1834, and refers to the thesis of Pater (1907), who collected records of 122 cases.

S. Flexner (1898) described a very typical case of syphilis of the stomach, of gummatous nature, which bears very close resemblances to the example now discussed. The patient was a man of 52 years, who died from perforation of the stomach with resulting peritonitis. A large gumma of the liver was found at autopsy, and the stomach showed a large ulcer (5 × 5 cm.), situated close to the cardia. Very careful histological examination of the ulcer was made, and changes of two distinct types determined. The earlier lesion consisted of a cellular infiltration of the submucosa with cells of granulation tissue type. Foci of necrosis had occurred within large collections of these cells. The second and later type of lesion was more common, and was found in all parts of the ulcer, including the tissues forming its elevated borders. It consisted of dense fibrous tissue formation, containing scattered granulation tissue cells. In this tissue also the blood-vessels showed extensive damage, endarteritis and endophlebitis obliterans, and hyaline thrombosis with organization, being very prominent.

E. Fraenkel (1899) described a remarkable case, seen in 1895. A man of 47 years was admitted suffering from diarrhoea and pain in the abdomen, and died from peritonitis following acute perforation. In the stomach thirteen ulcers were present, along with thirty-one others scattered along the small intestine. The possibilities of enteric fever and tubercle were fully considered, but a diagnosis of syphilis was finally decided on after very careful histological examination of the ulcers.

Soltan Fenwick (1901) recognized the occurrence of gastric syphilis, and claimed that about 5 per cent. of chronic gastric ulcers are of syphilitic origin. Other writers have attributed an even larger number of chronic ulcers of the stomach to syphilitic infection, Ewald (1905), for example, placing 10 per cent. of gastric ulcers in that category.

Turning to the more recent publications on gastric syphilis, the comprehensive paper by Monod (1921) has already been referred to. The most complete recent review of the subject is probably that of Wile (1920), in America, who gives a list of over fifty references to other writers. He states that he himself has seen three cases of gumma of the stomach in a period of four years. Smithies (1915) has reported an analysis of 7,545 patients suffering from all types of dyspepsia, among whom 26 gave a positive Wassermann reaction. Of the patients (1,603 in number) who showed evidence of definite organic disease in the stomach or duodenum, a proportion of 1.6 per cent. gave a positive Wassermann reaction. It appears to be quite unfair, however, to suggest that in the latter patients the gastric symptoms were necessarily due to syphilis, and not merely coincident with it.

Among other recent writers, Haas (1920) describes a case of 'syphilitic tumour' of the stomach, and Florand and Girault (1921) give a description of a large tumour in the epigastrium which at operation was discovered to be

a gummatous mass involving both the liver and stomach. Other examples are also referred to later in the text.

Varieties of Gastric Syphilis.

All writers are agreed that the commonest form of gastric syphilis arises as a submucous gumma, which tends to undergo the usual degenerative changes, and finally leads to ulceration. This variety has added importance from the fact that many more of such cases have been diagnosed as carcinoma of the stomach and treated as such, than as syphilis. A palpable tumour of varying size is always found on examination. Just as in carcinoma, loss of weight and marked cachexia are the rule, and examination of the gastric contents adds a further source of error in diagnosis, which is dealt with later.

Apart from gumma of the stomach, which is admittedly uncommon, other varieties of gastric syphilis which have been described must be extreme rarities. Monod (1921) has attracted attention to diffuse syphilitic infiltration as one cause of 'leather-bottle stomach', although there is no doubt that scirrhus carcinoma is a much more common reason for the occurrence of this somewhat rare deformity.

Wile (1920) adopts a very wide classification, which certainly seems to include all the possibilities. The rarity of all other forms when compared with gumma of the stomach is, however, fully acknowledged. His classification is as follows:

I. Gastric syphilis.

- (a) Syphilitic catarrh.
- (b) Syphilitic round ulcer.
- (c) Diffuse syphilitic infiltration of the gastric mucosa, with distortion of the stomach.
- (d) Pyloric syphilis—including ulcer, cicatrix, gumma, and diffuse infiltration.

II. Perigastric syphilis (with secondary implication of the stomach).

The Relation of Syphilis to the Common Round Ulcer of the Stomach.

It is quite obvious from clinical experience that syphilis has nothing to do with the aetiology of the majority of cases of simple round ulcer of the stomach. The question, however, must be raised whether syphilis may be the cause of even a small percentage of such ulcers. Wile, Fenwick, Smithies, Ewald, and others all believe that syphilis may be a cause of round ulcer. One statement by Wile has a particular interest in this connexion. He says while dealing with round ulcer due to syphilis that 'a recent case at the University Hospital, in Cabot's service, gave a negative Wassermann in the presence of a typical syphilitic ulcer. Spirochaetes were demonstrated by Warthin in the excised material.' This case

does not appear to have been published in full, but, in passing, it may be stated that this would seem to be the only other case, apart from our own, in which spirochaetes have been demonstrated in a syphilitic lesion of the stomach.

On the other hand the case reported by Fowler, and already referred to, shows how chronic round ulcer of the stomach may coincide with, and yet have nothing to do with, a syphilitic infection.

Gastric Analysis in Syphilis of the Stomach.

The occurrence of achlorhydria in the gummatous variety of syphilitic ulceration of the stomach, in contrast to the results of analysis in round ulcer, has attracted attention. The attention is well merited, since the absence of free hydrochloric acid tends to add support to an erroneous diagnosis of carcinoma.

Glaser (1921) deals most fully with the question of gastric analysis in cases which have been diagnosed as gastric syphilis with more or less certainty. He collected for his paper ten cases (including one of his own) in which the results of examination of the gastric contents are recorded. In all of these there was achlorhydria. The same conclusion is arrived at by Hausmann (1911) and Brugsch and Schneider (1915). Glaser also refers to the experiences of Neugebauer (1914), who found that among 200 soldiers suffering from syphilis there was complete absence, or only traces, of free HCl in 18 per cent. This of course introduces another question concerning the effects of early syphilis on the stomach and the variety of gastric lesion called 'syphilitic catarrh' by Wile. Smithies also states, in the statistical study of cases of dyspepsia already referred to, that among twenty-six cases regarded as probably due to syphilis on account of the positive Wassermann reaction, free hydrochloric acid was absent in twenty-four of them.

In our own case gastric analysis showed a complete absence of hydrochloric acid and the presence of lactic acid. Taking into consideration the extent of the damage to the stomach in gummatous ulceration, these results of analysis are not surprising.

The Effects of Antisyphilitic Treatment in Gastric Syphilis.

This is a difficult part of the subject to deal with adequately since the clinical diagnosis is so rarely made with accuracy. It is certain, however, that cases have occurred associated with the presence of a large mass in the epigastrium, in which very rapid cure, with gain in weight and removal of cachexia, have occurred in a few weeks after antisyphilitic treatment.

Hausmann (1911) describes a case of gumma of the stomach, diagnosed clinically by him, and treated with injections of salvarsan. There was a large tumour, involving both the greater curvature and the pyloric region. A history of syphilis was obtained, and the Wassermann reaction was weakly positive.

The tumour had entirely disappeared six weeks after treatment was commenced. A further point of interest recorded in this case is that whereas achlorhydria was noted on admission, free hydrochloric acid began to return two months after treatment began, and a month later the chemistry of the gastric contents was found to be normal.

Pezoldt (1915) reports a typical example of ulcerating gumma of the stomach in a negro, in which complete cure followed within a few weeks after treatment with salvarsan and iodides.

Sherran (1921) has given a brief description of a case of especial interest. The patient had a large tumour of the stomach, thought to be malignant, and with all the characters of a carcinoma. On examining the patient further, however, a swelling was found in the left epididymis, which was diagnosed as of syphilitic nature. The Wassermann reaction was found to be positive. After antisyphilitic treatment, the nature of which is not stated, the tumour of the stomach disappeared completely within three weeks.

Fenwick refers to cases of ulcer of the stomach which were only cured after the use of large doses of potassium iodide or other antisyphilitic treatment, but in these cases the exact nature of the stomach lesion remains uncertain.

It is a curious fact that recent writers, especially of the French School, have supported the claims of the older methods of antisyphilitic treatment (mercury and, more especially, iodides) against the newer salvarsan remedies, in cases of gastric syphilis. This is a matter on which further experience is required.

Record of a Case of Syphilis of the Stomach, in which abundant Spirochaetes were found in the Lesion.

The patient, a clerk aged 57 years, was admitted to University College Hospital under the care of Dr. Sidney Martin, F.R.S., complaining of pain in the chest and abdomen, and of constipation alternating with bouts of diarrhoea. His previous health had been good, and no history of syphilis was elicited. He was pale and emaciated, and stated that he had lost four stone in weight during the past six months. There was considerable pain and tenderness in the left epigastrium, and a tumour could be felt in this region, protruding below the left costal margin. The mass was about the size of a hen's egg, hard, of irregular contour, and moved with respiration. A diagnosis of malignant growth was made. The liver was not enlarged, and no superficial lymph glands appeared to be involved. The gastric contents were examined on two occasions, at twelve days' interval, and on both of these free hydrochloric acid was absent, whilst lactic acid was present. About a month after admission a fairly copious haematemesis (12 oz.) took place, followed by a second very large haemorrhage about a fortnight later. After the second attack the patient collapsed, and gradually became unconscious. Signs of probable perforation of the stomach were also recognized, but in view of the diagnosis of carcinoma which had been made, and the critical state of the patient, an operation was not considered

advisable. Since there appeared so little doubt of the diagnosis of malignant growth, the Wassermann reaction was not carried out.

At the autopsy generalized peritonitis was found, resulting from perforation of the stomach. Apart from the stomach no other organs showed noteworthy pathological changes, and nothing was found suggesting the presence of an antecedent syphilitic infection. The liver was normal in size and free from disease.

On opening the stomach a remarkable condition was found, which at first sight appeared to fit in with diagnosis of gastric carcinoma of scirrhus type. On examining the organ more closely, however, it was seen that the characters of the ulceration were peculiar, while further dissection showed the lymphatic glands draining the gastric area to be small and free from all signs of cancerous invasion. The absence of secondary growth in the liver in the presence of such extensive ulceration also excited notice.

The macroscopic appearances of the inner surface of the stomach are well shown in the photograph (Plate 8, Fig. 1) and may now be described in some detail.

The lesser curvature of the stomach, from a point about one inch below the cardia to the pylorus, is completely ulcerated. The ulcer also extends over a large part of the greater curvature, and this portion has been cut in two in opening the stomach, so that part appears in the upper part of the opened-out specimen and part below. Two perforations are seen in the ulcer on the posterior wall of the stomach, fairly close together, while in two other adjacent spots penetration of the wall is almost complete (Fig. 1, D). The chief characteristic of the ulcer is its irregular shape and greatly thickened borders. The floor is bare, but the margins are covered with shaggy shreds of necrotic tissue, mixed with shreds of tenacious mucus. Only about two-fifths of the stomach wall is free from ulceration.

The stomach wall is greatly thickened in most parts of the floor of the ulcer, except at the points where perforation has ensued. The thickness is fully half an inch in most places, while in relation to the raised margins it is fully three-quarters of an inch. In section the thickened wall has everywhere a dense pearly-white appearance, such as is often seen characteristically in scirrhus carcinoma of the organ. Particular attention may be drawn to one particular area (Fig. 1, c) of the lesion where the process appeared to be more acute and recent than elsewhere. A similar small area was originally present on the extreme upper border of the specimen, but was removed for histological examination. The results of the examination of this portion of the ulcer are dealt with later.

Owing to the peculiar characters of the ulcer, portions from as many situations as possible were removed for histology, without destroying the macroscopic specimen. The results were as follows, and compare closely with those described by Flexner in his very similar case. Flexner's excellent description was of course written long before the discovery of the causal organism of syphilis in 1905.

Sections from the central portion of the ulcer well removed from the raised margins show, in all the paraffin blocks examined, the same type of structure.

Adjacent to the gastric cavity the floor of the ulcer is composed of a thin layer of necrotic fibrous tissue, infiltrated with round cells. Not a trace of the glandular acini, or of the muscularis mucosae, remains. Below this necrotic layer is a thick zone of dense adult scar tissue, with many fibrils and comparatively few cells. Here and there, however, areas of less adult connective tissue occur, with large fibroblasts, and numbers of small thin-walled capillaries. Many of these small vessels are encircled by round-celled infiltration. Embedded in the mass of dense fibrous tissue are a few arterial trunks, some of them very close to the surface of the ulcer. The walls of these arteries are thickened, and show an advanced stage of endarteritis obliterans, but no thrombosis (Plate 9, Fig. 3). The accompanying veins have also thickened walls, but there is no definite endophlebitis. Hyaline thrombosis is not seen. Thick strands of firm fibrous tissue are also seen to have penetrated from the main mass into the muscular coat of the stomach, so as to interrupt the plain muscle fibres in all directions. In the midst of some of these strands of fibrous tissue, small collections of atrophic and disintegrating muscle fibres are visible. The blood-vessels in the muscular coat do not exhibit thickening, endarteritis, or thrombosis.

In blocks of tissue taken from other parts of the ulcer, especially from the greatly thickened borders, the histological changes are of a very much more active kind. The whole of the mucosa and submucosa is entirely replaced by an actively growing granulation tissue, which is necrotic and shaggy on its surface. The granulation tissue shows abundant fibroblasts, and thin-walled capillaries running up towards the surface (Fig. 4). The whole tissue is also densely infiltrated with cells of lymphocytic type. Wide branches of the granulation tissue extend into the muscular layers of the organ, following for the most part the track of the larger blood-vessels, and separating widely the strands of muscle fibres. In other situations the muscular coat appears to be intact, and the only change noted is the presence of round-celled infiltration along the blood-vessels. The arteries and veins in relation to these actively proliferating granulomatous parts of the ulcer are entirely free from endarteritis. In a few places large masses of granulation tissue have necrosed and separated, so as to expose even the deeper layers of the muscular coat. It is easy to see how in this way perforation must have occurred, and since the main blood-vessels have not undergone endarteritis obliterans the reason for the two attacks of haematemesis is not far to seek.

Other sections from the ulcer show almost all the intermediate stages between the actively growing granulomatous lesion and the almost obsolete zone of scar tissue in the centre, so that it is quite easy histologically to follow the whole course of the pathological process.

No giant cells are seen in any of the sections examined. Except for the very localized and deep patches of necrosis which reach down to the muscular layers, the tissues below the superficial floor of the ulcer are free from degenerative changes and caseation. The subperitoneal connective tissue is in most places greatly thickened behind the area covered by the ulcer, the loose connective tissue

normally found there being converted into dense fibrous tissue, containing many fibrils but comparatively few cells.

At certain parts of the edge of the ulcer, on the greater curvature of the stomach, the margins are not notably raised and thickened, and the site of origin of the lesion is more easily traced. Here the early pathological changes are seen spreading in the submucosa only, below an intact muscularis mucosae, but much altered mucous membrane. The early stage of the lesion consists of large and dense collections of small round cells encircling the blood-vessels, with, in a few places, the development of small patches of granulation tissue. An appearance difficult to explain is the occurrence immediately below the mucosa of several oval nodules of very dense and almost hyaline fibrous tissue, so perfectly circumscribed as to resemble small fibromata.

In the parts of the stomach wall which are free from ulceration (e. g. near the cardiac end) the mucous membrane shows an interstitial gastritis of atrophic type (Fig. 5). The glands are shortened, incomplete, and run in all directions. They are embedded in interstitial tissue which contains many small round cells. Some glandular acini are distended to form small cysts filled with mucus. The blood-vessels in the submucosa are not altered, but are frequently surrounded by collections of small cells of the usual type.

The lymphatic glands draining the stomach posteriorly were also examined histologically, but exhibited nothing more than the catarrhal changes commonly found in the neighbourhood of any chronic inflammation.

Examination for Spirochaetes.

In view of the histology of the lesion as described above, the conclusion was arrived at that the lesion was of syphilitic nature, but it was felt that even more convincing proof would be given if spirochaetes could be demonstrated in the actively growing parts of the lesion. For this purpose eight small blocks of tissue, from known situations, were treated by Levaditi's silver method.

In sections from one block only, spirochaetes were discovered in great numbers penetrating deeply in the actively growing granulomatous tissue. Near the free surface the spirochaetes in the necrotic surface layer were mixed with other organisms, but in the deeper granulation tissue close to the muscular coat they were unaccompanied by any other demonstrable organism. They had the typical appearances of *Trypanema pallidum*, having from eight to fourteen small and regular curves (Fig. 2). The part of the ulcer from which the block containing the spirochaetes was obtained differed in some important respects from other areas of the ulcer. It was reddish in colour, and sharply marked off from the neighbouring parts of the floor. Another area of exactly similar character is well shown in the photograph of the specimen (Plate 8, Fig. 1, c).

Conclusions.

1. The case recorded illustrates the chief clinical and pathological facts in the commonest variety of syphilis of the stomach, which arises as a submucous gumma. A particularly interesting point was the discovery of abundant spirochaetes in an actively developing part of the lesion.

2. Cases of this kind resemble so closely carcinoma of the stomach that an erroneous diagnosis, in the absence of other overt signs or a history of syphilis, is very likely to be made.

3. The resemblances to carcinoma include :

(a) The invariable presence of a palpable tumour in the epigastrium.

(b) The occurrence of loss of weight and cachexia.

(c) The absence of free hydrochloric acid in the gastric juice.

4. Perforation of the gummatous ulcer may lead to fatal peritonitis.

5. Haematemesis may occur from necrosis of part of the gummatous mass, before endarteritis and thrombosis of the arteries have occurred.

6. The pathological lesion consists essentially of the development of a granulation tissue mass, or gumma, in the submucous layer of the stomach, which soon leads to destruction of the overlying mucosa and to ulceration.

7. The lesion may be seen at various stages, the centre of the ulcer being almost healed and transformed into scar tissue, while other parts, notably the raised margins, show the active stage of granulation tissue formation. The larger blood-vessels in the obsolete central parts of the ulcer show well-developed endarteritis obliterans, but in the zone of active growth these changes are absent.

8. From a brief historical survey of the literature it is evident that the prognosis in such gummatous lesions of the stomach is good when antisyphilitic treatment is employed. The literature also shows that this is the most frequent form of an uncommon variety of visceral syphilis. Many more descriptions of the condition have been found on record than the scanty references to the subject in British text-books might be taken to indicate.

Other varieties of gastric syphilis must at present be regarded as of extreme rarity, although it may be pointed out that little is known of the effects of early syphilis on the structure of the stomach.

The writer wishes to thank Dr. Sidney Martin for permission to publish this case.

BIBLIOGRAPHY.

1. Brugsch, T., and Schneider, E., 'Syphilis und Magensymptome', *Berl. klin. Wchnschr.*, 1915, lii. 601.
2. Chiari, H., 'Ueber Magensyphilis', *Rudolf Virchow Festschrift*, Berlin, 1891, ii. 297.
3. Ewald, 'Das Magengeschwür', *Deutsch. Klinik*, 1905, v. 457 (quoted by Wile).
4. Fenwick, W. Soltau, 'Syphilitic Affections of the Stomach', *Lancet*, Lond., 1901, ii. 835.
5. Flexner, S., 'Gastric Syphilis, with the Report of a Case of Perforating Syphilitic Ulcer of the Stomach', *Amer. Journ. Med. Sci.*, 1898, N.S., cxvi. 424.
6. Florand, A., and Girault, A. L., 'Syphilome gastro-hépatique', *Presse Médicale*, Paris, 1921, xxix. 841.
7. Fowler, W. F., 'Benign Gastric Ulcer in Syphilis', *Surg. Gynaecol. and Obstetrics*, May, 1921; extract in *Lancet*, 1921, ii. 689.
8. Fraenkel, E., 'Zur Lehre von der acquirirten Magen-Darm-Syphilis', *Virch. Arch. f. path. Anat. u. Physiol.*, Berlin, 1899, clv. 507.
- 9.¹ Glaser, F., 'Ueber Anacidität bei syphilitischem Magengeschwür', *Mediz. Klinik*, 1921, xl. 1199.
10. Haas, G., 'Ein Fall von syphilitischem Tumor des Magens', *Archiv f. Verdauungskr.*, Berlin, 1920, xxvi. 68.
11. Hausmann, T. H., 'Ein Fall von gummosen Magentumor, geheilt durch Salvarsanbehandlung', *Münch. med. Wchnschr.*, 1911, lviii. 511.
12. Hutchinson, Jonathan, *Syphilis*, Lond., 1887.
13. Monod, G., 'Syphilis of the Stomach', Occasional Lecture, reported in *Proc. Roy. Soc. Med.*, Lond., 1922, xv. 1.
14. Neugebauer, H., 'Sekundäre Syphilis und Magenveränderungen', *Wien. klin. Wchnschr.*, 1914, xxvii. 845.
15. Pezoldt, 'Kasuistische Mitteilung über einen Fall von Karzinom vortäuschendem ulcerierten Gumma der vorderen Magenwand (Pylorus)', *Archiv f. Schiffs- u. trop. Hygiene*, 1915, xix. 151.
16. Sherran, J., 'The Value of Pathological and X-ray Examinations in Abdominal Surgery', *Lancet*, Lond., 1921, ii. 689.
17. Smithies, F., 'Syphilis of the Stomach', *Journ. Amer. Med. Assoc.*, 1915, lxxv. 572.
18. Stoeckenius, W., 'Beobachtungen an Todesfällen bei frischer Syphilis', *Ziegler's Beitr. z. path. Anat. u. z. allg. Path.*, Jena, lxxviii. 185.
- 19.¹ Wile, Udo J., 'Syphilis of the Stomach', *Archiv. of Dermatol. and Syph.*, Chicago, 1920, N.S., i. 543.

¹ With literature.

DESCRIPTION OF PLATES.

PLATE 8. FIG. 1. Photograph of the specimen of syphilis of the stomach. A. Oesophagus. B. Pylorus. C. In tissue from an area similar to this abundant spirochaetes were found. D. Two perforations of the ulcer.

PLATE 9, FIG. 2. Microphotograph of a section showing the character of the spirochaetes. Stained by Levaditi's method. ($\frac{1}{15}$ " oil-immersion lens, No. 4 eyepiece.)

FIG. 3. Microphotograph of a small artery in the floor of the ulcer, showing endarteritis obliterans. Haemalum and eosin ($\times 50$).

FIG. 4. Microphotograph of a moderately active part of the ulcer, showing the granulation tissue. Part of the muscular coat of the stomach is seen below. Haemalum and eosin ($\times 50$).

FIG. 5. Microphotograph of the mucosa of the stomach in a part not involved by the ulcer. The changes are those of interstitial gastritis, and the muscularis mucosae remains intact. Haemalum and eosin ($\times 50$).

FIG. 1

FIG. 2

FIG. 3

FIG. 4

FIG. 5