number of NATURE as having been seen by Mr. Denning at Bristol on the evening of the 23rd ult., as well as a single other shooting star noted during a watch of the sky whenever it was clear by the same observer on that evening, seem, from their apparent courses, to have been both possibly directed from the now well-known radiant point of the meteors of Biela's comet, but only a small number of shooting stars appears to have been visible on that night. During a watch for ten minutes, kept at four different times between half-past 7 and half-past 9 P.M. on the following evening, near Regent's Park, in London, Mr. H. W. Jackson observed, in each watch, a shooting star as bright as a first-magnitude star directed in each case from the recently determined radiant point; while, with an equally uncloudy sky, on the following evening of the 25th, three small meteors from the same radiant point, and two brighter ones from other directions, were seen by Mr. Denning at Bristol between 8 and 11 P.M. in two intervals of a watch for half an hour. A small meteor of the same stream was recorded by Mr. Denning on the 26th, but their numbers on this, as on the previous nights, were evidently inconsiderable, a clear view of the sky on the night of the 26th, between 11h. 20m. and 12h. 40m., affording Mr. Jackson, at Tooting, no observation of a single meteor. The time of first approach and commencement of the bright star-shower seen on the evening of the 27th ult. must therefore have been later than shortly before 1 o'clock on the previous morning. A completely overcast state of the sky after midnight on the night of the shower appears to have prevented observations of its close, and probably of its complete extinction on that night, after the greatly-diminished intensity which it had then attained; but a correspondent in London informs me that, in spite of the densely overcast state of the sky, which prevented any view of the shower from being obtained in the metropolis, an exceedingly bright meteor was seen to flash like lightning through the clouds at about 4 o'clock, A.M., on the 28th. On the evening of the 28th Mr. Greg watched for shooting stars, and for any remnants of the star-shower of the previous evening which might be visible, at Buntingford, in Hert's; but although the sky was quite clear he failed to see any meteors. A strict watch for outlying meteors of the shower was also kept by two observers at Hawkhurst, in Kent, on the evening of the 28th, where the sky was quite cloudless between 9h. and 11h. 15m. P.M., but without success, only four shooting stars of ordinary character being visible during more than two hours of their simultaneous watch. An interval of about forty-four hours is thus clearly determined in which the first indications of the star-shower must have arisen, reaching the maximum of its intensity towards the middle of the period, and disappearing so completely before its termination as to leave the whole expanse of the sky almost as perfectly free from shooting stars as it was before its commencement.

In addition to the early appearance of the shower recorded in NATURE of the 5th inst. as having been seen near Dublin, I have received the following communications from observers relating to its early visibility and abundance. At Brancepeth, near Durham, Mr. Joseph Lawson first began to count the meteors at 5 o'clock, and continued to enumerate them until, towards 7 o'clock, their constantly increasing frequency obliged him to desist. During the last half hour of his estimation the rate of their appearance was about ten per minute, while a total number of 1,000 meteors was counted between 5 and 7 o'clock. The rate of their appearance in the first was thus little less than that of their appearance in the latter portion of his watch. Mr. S. J. Miller, at Wisbeach, first noticed the abundance of shooting stars at 5h. 40m., and counted in three minutes twenty meteors, or about six or seven per minute. Soon after this, at about 6 o'clock, their number was ten or twelve per minute. During their appearance before 6 o'clock, Mr. Wood, at Birmingham, considered them to be falling from 5h. 45m. when he first observed them, at a rate of about fifteen per minute, while the result of his enumeration between 6h. and 7h. was at least 600 meteors, and in occasional intervals between 6h. and 7h. 15m., when the sky became gradually overcast, the average rate of appearance was found to be increasing nearly uniformly from 8 to 10 per minute, showing that the intensity of the display before 6 o'clock was little inferior to the maximum which it appears afterwards to have gradually attained. At York the first attempt to estimate their numbers was made by Mr. T. H. Waller, at 5.30 o'clock, when the meteors were found to be appearing at the rate of 12 or 15 per minute. At 6 o'clock their numbers had increased to about

20, and at 7 o'clock to about 30 or 40 per minute for one observer. At 8 o'clock the best determination of their numbers was however obtained by two of Mr. Waller's scholars, who, standing back to back, counted 79 meteors in 1m. 15s. or 31 meteors per minute for each observer. The principal maxima of the shower at about 6h. 30m., and 8h. 15m. P.M., as mostly clearly shown by the continuous observations of Mr. Lowe at Highfield House, and by Prof. Grant, who was assisted in his observations at the Glasgow Observatory by Prof. G. Forbes, are also generally indicated by the observations which I have received, and the gradual diminution of the shower after 9 o'clock was observed at Newcastle-upon-Tyne as well as by some of the observers who combined in their watch for its appearance for the British Association. The frequency of the meteors after 9 o'clock was regarded by Mr. Wood as not more than a quarter of what it was at about 6 o'clock, when they were first seen, yet these numbers continued for some time to be considerable and quite unusual. Near Rothbury, in Northumberland, repeated estimations of their abundance were made by Mr. G. A. Lebour in different parts of the sky, and at different times between the hours of 7 and 10 o'clock, with the following results:—

From 7 ^h 20 ^m to 7 ^h 28 ^m	100 meteors falling perpendicularly in the west (the east cloudy).
7 35 7 40	100 meteors falling vertically in the east (the west cloudy).
7 45 7 48	100
9 0 9 15	Meteors too numerous to count at 9h., falling in showers at 9h. 15m., especially in the north; one bright red with beautiful red streak, which lasted more than 10 seconds.
10 0	Meteors still falling, but in smaller numbers.

In 20 minutes after 10 o'clock 35 meteors were seen by one observer in a clear part of the sky, which scarcely exceeded a quarter of the sky. In 10 minutes after 11 o'clock ten meteors only were counted in the same space, and in 20 minutes after 12 o'clock 9 meteors only could be seen. The rate of decrease of the shower from shortly before 10 o'clock until after midnight was thus apparently more rapid, and the decline of its intensity appears to have been considerably more complete than the first brightness and progress of its gradual increase may be concluded to have been during the earliest hours of its observation.

Newcastle-upon-Tyne, Dec. 7

A. S. HERSCHEL

The De Novo Production of Living Things

WILL you allow me to ask Dr. Bastian to state in your columns, in reply to this note, the specific gravity of the turnip infusion, to which a "fragment" of cheese has been added, and which, he states, has rarely failed to give him positive results in his endeavours to obtain evidence of the *de novo* production of living things from dead matter. If Dr. Bastian should be unable to give the specific gravity of the infusion, perhaps he will be so good as to state accurately the weights of water, turnip, and cheese employed. I am, of course, acquainted with the directions he has already given, but wish to avoid any objection from him or others on the score of improper preparation of the infusion to results which I may obtain and publish.

E. RAY LANKESTER

Exeter College, Oxford, Dec. 7

The Birth of Chemistry

YOUR correspondent, "A. H.," in alluding to my mention of the *Insula Cassiterides*, inquires whether the name was derived from a Sanskrit source. The word *kaṣṭhīrepa* is used both by Homer and Hesiod, and it is possible that it may have been borrowed from the Sanskrit *kaṣṭhira*, and that tin was first procured from India. The Sanskrit word for tin—*kaṣṭhira*—is clearly related to the verb *kāś*, to shine. It is strange that the Arabic word for tin is *kāṣṭir*, closely resembling the Sanskrit, although there is no family relationship between the languages. Possibly the Phœnicians first procured tin from India, and gave it a name resembling its native name *kaṣṭhira*; then the Greeks converted the Phœnician word into *kaṣṭhīrepa*, the Romans borrowed the word from the Greeks, and the fact of the scarce