2. DESCRIPTIONS OF NEW SPECIES OF EULIMA, TRIPHORIS, ETC.,
FROM THE COLLECTION OF HUGH CUMING, ESQ.

BY ARTHUR ADAMS, F.L.S. ETC.

1. *EULIMA MODICELLA*, A. Adams. *E. testá subulato-pyramidalí, albd, subpellucidá, flexuosá; anfractibus 11, planulatis, varicibus lateralibus continuis impressis, instructis; anfractu ultimo, in medio, vix angulato; aperturá ovali; columellá anticè subrectá; labro anticè producto.*

Hab. Island of Zebu, sandy mud, 7 fathoms. Mus. Cuming.

2. *EULIMA GRANDIS*, A. Adams. *E. testá subulato-pyramidalí, albd, solidá, flexuosá, opacá; anfractibus 15, planulatis, varicibus lateribus continuis instructis; anfractu ultimo angulato; aperturá obliquá, oblongo-ovalí, labio anticè calloso; labro margine flexuoso, anticè subreflexo.*

Hab. Island of Burias, coral sand, 7 fathoms. Mus. Cuming.

3. *EULIMA PORCELLANA*, A. Adams. *E. testá subulatá, albd, solidá, opacá, apice subflexuoso; anfractibus 13-14, planulatis, varicibus impressis irregularibus lateralibus; aperturá oblongo-ovalí, labio anticè calloso, vix reflexo; labro margine, in medio, dilatato.*

Hab. —? Mus. Cuming.

4. *EULIMA ACUTA*, A. Adams. *E. testá aciculato-turritá, albidá, rectá, subopacá; anfractibus duodecim, planiusculis, ultimo rotundato; aperturá oblongá, anticè subreflexá, labio subincrassato; labro margine recto.*

Hab. Sual, province of Cagisanan, island of Luzon, sandy mud, 7 fathoms. Mus. Cuming.

5. *EULIMA CUSPIDATA*, A. Adams. *E. testá subulato-pyramidalí, albidá, solidá, rectá; anfractibus 12, convexiusculis, anfractu ultimo rotundato; aperturá oblongo-ovalí, labio anticè calloso, subrecto; labro acuto.*

Hab. Sibonga, island of Zebu, in loose coral under stones, low water. Mus. Cuming.

6. *EULIMA OBESULA*, A. Adams. *E. testá pyramidalí-ovatá, albd, solidá, nitidá, opacá; anfractibus sex, convexis, ultimo rotundato; aperturá oblongo-ovalí; labro margine incrassato, nec limbato vel reflexo.*

Hab. Gindulman, isle of Bohol, in soft mud, 8 fathoms. Mus. Cuming.

7. *EULIMA TEINOSTOMA*, A. Adams. *E. testá subulato-turritá, rectá, albidá, nitidá, subpellucidá; anfractibus 12, planulatis, lined impressá infra suturas; anfractu ultimo rotundato; aperturá oblongo-ovalí, anticè producto, labio subrecto, anticè reflexo; labro margine, in medio, dilatato.*

Hab. Feejee Islands, on coral reefs, in sand, low water. Mus. Cuming.

8. *EULIMA FLEXUOSA*, A. Adams. *E. testá subulato-turritá, albd, flexuosá, solidá, subopacá; anfractibus 15, planulatis, lined impressá subpellucidá ad suturas; anfractu ultimo rotundato; aperturá oblongá; labro margine flexuoso, in medio producto.*
Hab. —? Mus. Cuming.
9. *EULIMA ACLIS*, A. Adams. *E. testá subulato-turritá, albidá, solidá, subopacá; anfractibus 11, planulatis, ultimo rotundato, anticè subproducto; aperturá oblongá, labio anticè subreflexo.*
Hab. Singapore, coarse gravel and sand, 12 fathoms. Mus. Cuming.
10. *EULIMA PYRAMIDALIS*, A. Adams. *E. testá subulato-pyramidali, albd, nitidá, subpellucidá; anfractibus decem, planulatis, lined impressá prope suturas, anfractu ultimo subangulato; aperturá oblongo-ovali; labro margine, in medio, subproducto.*
Hab. Isle of Capul, on the reefs in sand, low water. Mus. Cuming.
11. *EULIMA POLYGYRA*, A. Adams. *E. testá subulato-pyramidali, albd, subopacá, apice tortuoso; anfractibus permultis, planulatis, ultimo angulato; aperturá obliquá, subtetragonali, labio anticè reflexo; labro, in medio, valdè dilatato.*
Hab. Cagayan, province of Misamis, isle of Mindanao, sandy mud, 50 fathoms. Mus. Cuming.
12. *EULIMA VITREA*, A. Adams. *E. testá subulatá, acutá, rectá, albidá, vitreá, pellucidá; anfractibus planulatis, lined impressá prope suturas; anfractu ultimo rotundato; aperturá oblongo-ovali, labio anticè recto, in medio subtortuoso; labro anticè subreflexo.*
Hab. Feejee Islands; from the stomach of a *Holothuria* (Captain Swain). Mus. Cuming.
13. *EULIMA GUILDINGII*, A. Adams. *E. testá subulatá, rectá, nitidissimá, albd, pellucidá; anfractibus planulatis, ultimo subrotundato, elongato; aperturá oblongo-ovali, labio vix tortuoso; labro margine flexuoso.*
Hab. St. Vincents, West Indies, sandy mud, deep water. (Rev. L. Guilding.) Mus. Cuming.
14. *EULIMA CUMINGII*, A. Adams. *E. testá subulato-turritá, albd, rectá, solidá, opacá; anfractibus 13, convexiusculis, varicibus irregularibus impressis instructis; anfractu ultimo rotundato; aperturá oblongo-ovali, labio anticè calloso, incrassato; labro margine recto.*
Hab. Lord Hood's Island, South Seas, on the *Avicula margaritifera*. Mus. Cuming.
15. *TRIPHORIS VARIEGATUS*, A. Adams. *T. testá subulato-pyramidali, in medio tumidá, albd, maculis triangularibus rufo-fuscis variegatá; anfractibus planulatis, triseriatim granulatis, granis aequalibus, interstitiis punctatis, suturis impressis; canali brevi, apertá.*

Hab. St. John's. Mus. Cuming.

A large variegated species, somewhat resembling in general appearance *T. ornatus*, Desh.

16. *TRIPHORIS PULCHELLUS*, A. Adams. *T. testá subulato-pyramidalí, in medio tumidá, fuscá, serie moniliformi albo ornatá; anfractibus convexiusculis, triseriatim granuloso-carinatis, granorum serie inferiore prominulá, superiore multo minore; aperturá rotundatá, constrictá; canali brevi, recurvo.*

Hab. —? Mus. Cuming.

A handsome brown species, with a white series of bead-like granules at the lower part of each whorl.

17. *TRIPHORIS NIGRO-FUSCUS*, A. Adams. *T. testá pyramidalí, nigro-fuscá; anfractibus planis, triseriatim granulatis, granulis æqualibus, confertis, anfractuum suturis impressis, basi convexá.*

Hab. Sydney, low water, under stones (*Mr. Strange*).

A black-brown species, with three rows of regular, equal-sized granules on each whorl. Mus. Cuming.

18. *TRIPHORIS FESTIVUS*, A. Adams. *T. testá pyramidalí, basi planá fuscá, albidá, fasciis fuscis interruptis, transversis, ornatá; anfractibus planis, cingulis duabus granorum instructis; interstitiis valde punctatis.*

Hab. Port Lincoln. Mus. Cuming.

A small prettily-marked species, with two rows of granules on each whorl, and the interstices deeply punctured.

19. *TRIPHORIS SCITULUS*, A. Adams. *T. testá subulato-pyramidalí, albidá, nitidá, subpellucidá, suturis rufo-tinctis; anfractibus convexiusculis, cingulis tribus nodorum ornatis, cingulá medianá majore moniliformi, nodorum interstitiis fuscis, anfractu ultimo basi fusco; canali brevi, aperto.*

Hab. Port Lincoln. Mus. Cuming.

A semipellucid, white and brown species, with the middle row of nodules very prominent.

20. *TRIPHORIS ALBIDUS*, A. Adams. *T. testá subulato-pyramidalí, albidá, nitidá; anfractibus planulatis, subimbricatis, granoso-clathratis, granis oblongis, serie granorum inferiore prominulá, anfractu ultimo basi fulvo; canali brevi, subrecurvo.*

Hab. Honduras (*Dyson*). Mus. Cuming.

A solid, white, shining, pyramidal species, with oblong granules disposed in three series on each whorl.

21. *TRIPHORIS VESTALIS*, A. Adams. *T. testá turrito-subulatá, dextrorsá, albá, subnitidá; anfractibus 13, convexis, suturis impressis, triseriatim granulatis, interstitiis alveolatis.*

Hab. Honduras. Mus. Cuming.

A delicate and chaste right-handed species, with convex whorls, and pits between the granules.

22. *TRIPHORIS CINGULATUS*, A. Adams. *T. testâ elongato-pyramidali, cinereâ; anfractibus sexdecim ad octodecim, spiraliter tricingulatis, cingulâ medianâ minore, interstitiis carinarum longitudinaliter valde striatis.*

Hab. Red Sea (Rüppell). Mus. Cuming.

An ashy-grey species, with three smooth keels on each whorl, and the interstices strongly striated: somewhat similar to the *T. corrugatus* of Hinds.

23. *TRIPHORIS LABIATUS*, A. Adams. *T. testâ subulato-pyramidali, nigro-fusca, in medio tumida, spirâ apice obtusa; anfractibus 10, planulatis, triseriatim granuloso-carinatis, suturis concavo-impressis; labro reflexo, dilatato, albido; canali brevi, subrecurvo.*

Hab. Sydney, under stones, low water (Mr. Strange). Mus. Cuming.

A small, nearly black shell, with the outer lip dirty white or pale fuscous.

24. *MESALIA STRIATA*, A. Adams. *M. testâ subulato-turritâ, fulvâ; anfractibus 10-12, planulatis, superioribus longitudinaliter plicatis, inferioribus levibus, transversim striatis, striis impressis, subdistantibus; anfractu ultimo subangulato; aperturâ ovali, labio subplanulato, anticè subreflexo; labro acuto, integro.*

Hab. Philippines. Mus. Cuming.

25. *MESALIA DECUSSATA*, A. Adams. *M. testâ subulato-turritâ, in medio subcylindraceâ, pallidè rubro-fusca; anfractibus novem, convexiusculis, plicis longitudinalibus confertis, et sulcis impressis, transversis, decussatim ornatis; aperturâ semiovali, labio subcalloso, anticè subreflexo, integro; labro incrassato, margine integro.*

Hab. Masbate, Philippines. Mus. Cuming.

26. *RISSOINA SEMIGLABRATA*, A. Adams. *R. testâ subulato-pyramidali, albâ, solidâ, nitidâ; anfractibus convexiusculis, supremis transversim striatis, inferioribus glabratis; aperturâ semiovali, anticè subcanaliculatâ, labio incrassato; labro dilatato, crasso, intus tuberculis parvis instructo, margine subreflexo.*

Hab. Deleguete, isle of Zebu, found under stones, low water. Mus. Cuming.

A species having very much the aspect of a *Eulima*. In this species there are two tubercles on the inner surface of the outer lip.

27. *RISSOINA EULIMOIDES*, A. Adams. *R. testâ subulato-pyramidali, albâ, solidâ, nitidâ; anfractibus planiusculis, suturis impressis; aperturâ semiovali, anticè subcanaliculatâ, labio levigato, subincrassato; labro margine crasso, in medio dilatato, intus tuberculo minuto instructo.*

Hab. Island of Capul, on coral reefs in sand, at low water. Mus. Cuming.

A small polished *Eulima*-like species, with a single small tubercle on the inner surface of the outer lip.