Review: Organic Matter of the Sea Bottom

Source: *Journal of Ecology*, Vol. 3, No. 3 (Sep., 1915), p. 182

Published by: British Ecological Society

Stable URL: <http://www.jstor.org/stable/2255389>

Accessed: 26-06-2016 07:17 UTC

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at
<http://about.jstor.org/terms>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

British Ecological Society, *Wiley* are collaborating with JSTOR to digitize, preserve and extend access to
Journal of Ecology

by the metabolic products present in fir humus, including various oils, esters, tannin, formic acid, etc. These substances were added to ordinary soil and were found to retard growth, both in seedlings and in older plants, usually in a marked degree; in many cases the presence of these bodies even in moderate concentration caused death of the seedlings, though older plants were able to recover after a time. The author also tried the effect of the vapour of the ethereal oils upon the aerial parts of the plants (*Impatiens*, *Brassica*, *Lepidium*, etc.) and obtained similar results. He next investigated the influence of these substances on the soil-inhabiting lower organisms and found that they injuriously affected the growth and fermentative activity of yeasts, and either reduced greatly the numbers of bacteria present or strongly inhibited their development and activity; the bacteria tried included the ammonifying and nitrifying species. It would therefore appear from the author's results that the meagre growth of plants in coniferous woodland may be in large measure attributed to the toxic action of the oils and other metabolic substances produced by conifers and found in the humus formed by these trees.

ORGANIC MATTER OF THE SEA BOTTOM

Boysen-Jensen, P. "Studier over Havbundens organiske Stoffer." *Danske Biol. Sta.*, Copenhagen, **22**, 1914, pp. 1-36.

The author has investigated the relative contributions of the plants of the *Zostera* belt on one hand and the plankton organisms on the other to the accumulated organic matter of the sea bottom in various parts of the Danish seas. He finds that *Zostera* is much richer than the plankton organisms in pentosan compounds (a class of carbohydrates), the sea bottom organic matter being intermediate in this respect. Apparently this bottom material is almost exclusively derived from *Zostera* in the case of the more land-locked waters (bays and lagoons), while the plankton organisms play a more important part as source of this material in the more open coast waters, as in the Kattegat. Calculations of the amounts of plankton and *Zostera* produced per sq. metre in the Danish waters inside the Skaw showed that the production of phytoplankton was about 100 gm. and that of *Zostera* about 120 gm. per sp. m. Apparently only a small part of the plankton production is deposited on the sea bottom, this deposition occurring evenly, and the result of these calculations agrees well with that of the chemical analyses. The bottom organic matter contains more nitrogen than does the *Zostera*, and it was found that the latter contained more nitrogen in the later than in the earlier stages of its decomposition—a result which may be explained either by the more rapid destruction of the non-nitrogenous than the nitrogenous substances, or by the occurrence of nitrogen assimilation, or by a combination of both these processes.

ALGAL VEGETATION OF DOLOMITE ROCKS

Diels, L. "Die Algen-Vegetation der Südtiroler Dolomitenriffe." *Ber. d. deutsch. bot. Ges.*, **32**, 1914, pp. 502-526.

The author has made a minute and careful study, with interesting results, of the lithophytic algal vegetation of a dolomite rock-wall, sloping south-southeast and situated at an altitude of about 1300 m. in the Weisslahn valley, Tyrol; the rocks in other localities were also examined with similar results. The vegetation (lithophyte formation)