the relation of the heart lesions to the mental symptoms are given. It is not contended that the heart disease was the sole cause in every case, but rather that it was the exciting cause, except wherein the cardiac troubles were rather the effect, and especial emphasis is given to the relationship of mental improvement to the treatment directed to the heart trouble. During a residence of nearly two years in a large State hospital the author observed but few of these cardio-mental cases, and the inference is that they are more often treated by the internist than by the alienist, and their psychic side has thus been minimized.

DEFECTS OF WILL FROM A MEDICAL STANDPOINT. H. T. Pershing (Journal A. M. A., Oct. 28).

The author defines volition practically as the idea of motion and locates its seat in the motor centres of the brain. The transformation of a sensation of motion into a memory and into the idea of its repetition and the cerebral mechanism involved is worked out by him in detail and practical deductions are drawn. The control of acts is best obtained, not by prohibitions that arouse the idea to be avoided, but rather by displacing it by something better. In hysterical paralysis, for example, the physicians should try to raise the emotional tone, then to excite the depressed sensory centres by electricity and passive motion, and then further to awaken the lost idea of motion by encouraging the patient to aid rhythmic passive motion by voluntary effort. The fundamental thing is to reawaken the lost sensations and ideas in the kinesthetic centre. This is the main idea in his article, but he elaborates it to apply to the various neurasthenic defects of will, as well as to morbid impulses or obsessions which are to be combated, giving also in a suggestive way the general principles of treatment adapted to the several types of disordered volition.

CAMP (Philadelphia).

SURGICAL TREATMENT OF INTRASPINAL TUMORS. Richard H. Harte (Annals of Surgery, October, 1905).

Harte presents an analysis of 92 cases operated for spinal tumor. This series shows a mortality of 47 per cent, but if late deaths are deducted, this is reduced to 28 per cent. The high death rate should not deter surgeons to operate, since the results are markedly beneficial. In 88 of the cases the nature of the tumor was recorded as follows: Sarcoma, 37; adhesions, 11; echinococcus, 8; fibroma, 16; syringomyelia, 5; endothelioma, 4; psammoma, 3; cyst, 3; fibromyoma, 2; osteoma, 2; tumor of vertebra, 1; myeloma, 1; lipoma, 1; lymphangioma, 1; primary carcinoma, 1; secondary carcinoma, 1; and dermoid cyst, 1. Of the 49 patients who survived the operation 59 per cent were cured, 34 per cent were improved and 6 per cent unchanged in condition. The nature of the growth is practically impossible to determine prior to operation. Sarcomata can be diagnosed with certainty only when clearly secondary. It is usually stated that extradural tumors are more frequent and much less fatal than intradural growths. In this series 50 were extradural and 36 intradural, the mortality in the former being 50 per cent and in the latter 47.21 per cent. This may be due to the fact that sarcomata are more frequently extradural. The relative mortality in males and females is 45 and 57 per cent respectively, but again this may be influenced by the greater frequency of sarcomata in the male. Of 82 cases in which the region affected is noted the upper dorsal was involved in 33, the lower dorsal in 24, the lumbosacral in 14, and the cervical in 11. Of the cases in which the cause of death was recorded fully one-half died of shock and hemorrhage, or of infection and meningitis.

COWLES (New York).