The advantages of the instrument are that: (1) it offers a simple means of exposing the entire tonsil for diagnostic purposes; (2) it can be used to advantage in expressing pus from the tonsils for bacteriologic examination; (3) the tonsils can be evacuated without an anesthetic without pain to the patient two or three times a week, and the patients rarely object to the repetition of the treatment. In very sensitive throats a 2 per cent. solution of cocain may be used before treatment.

In many cases one or more large pus pockets are present. The most frequent seat of the pus pockets in our cases has been near the upper pole between the anterior pillar and the tonsil. This part of the tonsil is often completely buried behind the pillar and cannot be exposed by simple retraction of the pillar.

Massage and expression of the tonsil in no sense replaces tonsillectomy. It is recommended only as an emergency measure where the radical operation is contraindicated during the height of an infection.

ACCIDENTAL INJECTION OF BILE DUCTS WITH PETROLATUM AND BISMUTH PASTE

ADOLPH REICH, M.D., NEW YORK

Miss J. de M. had a pelvic operation two years before consulting me. A few days after the operation she was taken with pain over the region of the liver, and a swelling, which appeared in the midaxillary line, was incised and discharged pus, leaving a persistent fistula between the tenth and eleventh ribs. She asked to be relieved of this fistula.

In order to know the direction of this fistula, I injected about 2 ounces of petrolatum paste and bismuth with a melting

Fig. 1.—Postero-anterior view, showing bile ducts filled with paste down to the duodenum.

point of 110 F. Immediately there was considerable pain in the pit of the stomach, the following day the patient's temperature was 102 by mouth, the urine was dark brown, she was jaundiced, and she could not hold anything on her stomach. Figure 1 shows the bile ducts filled with the paste down to the duodenum.

The treatment consisted of local hot applications, morphin for the pain, calcined magnesia one-half ounce every two hours until the bowels moved freely, and injection of warm olive oil through the fistula, about 1 ounce or more every three hours. The next morning the bowel movements had a dark slate color and kept so for two weeks, and the temperature

Fig. 2.—Anteroposterior view taken recently; bile ducts free from bismuth; a few grains of bismuth near the opening of the fistula.

fell to normal within one week, and all symptoms of the obstructive jaundice passed off gradually.

Figure 2, taken recently, shows the bile ducts entirely free from bismuth and only a few scattered grains of bismuth near the opening of the fistula. Pus is discharged only occasionally, and some of the bismuth finds its way out.

245 West Twenty-Fourth Street.

STUDIES OF ROENTGENOGRAMS OF THE ACCESSORY NASAL SINUSES

WITH A DESCRIPTION OF A NEW METHOD OF TAKING SUCH PICTURES:
PRELIMINARY NOTE

MAX UNGER, M.D., NEW YORK

In common with other nose and throat specialists, I have felt the disappointment resulting from the lack of help in diagnosis furnished by roentgenograms of the accessory nasal sinuses, especially in cases of suspected ethmoid and sphenoid disease.

The method now employed in the taking of roentgenograms of the accessory nasal sinuses is to put the plate in front of the patient's face and to shoot the roentgen ray through from the back of the head. This gives us an excellent picture of the frontal and maxillary sinuses, but the recognition of the ethmoids and sphenoids is a matter of guesswork. This is due to the fact that the latter cells are small, deeply situated and overlapped by the frontal and maxillary sinuses, the turbinates and the nasal bones.

Accordingly, I have devised a new method of taking roentgenograms of the sinuses. I employ a roentgen-ray film that is cut to the shape and size shown in Figure 1. This film is wrapped in waterproof paper, similarly to dental films. A