NOTE ON ASAFÆTIDA.

BY J. M. MARTIN, B.A., M.B., AND C. G. MOOR, M.A.

(Read at the Meeting, December 6, 1899.)

A SAMPLE of gum-resin of asafætida having been found to yield a very much higher ash than the standard given by the British Pharmacopœia, a number of samples were examined to determine the general condition of the commercial article.

Twelve samples were examined as to :

1. Matter soluble in 90 per cent. alcohol.
2. Ash of the gum.
3. Portion of ash soluble in HCl.

In all the samples, on grinding, much mineral matter was found, in one case a stone weighing 3·4 grammes.

The percentage of the gum soluble in alcohol (90 per cent.) ranged from 14·8 to 39·8, instead of the 65 per cent. of the British Pharmacopœia.

The ash of the gum ranged from 63·1 to 26·4 per cent., the British Pharmacopœia figure being 10 per cent.

The solubility of this ash in 50 per cent. HCl varied from 15·7 to 57·9 per cent., but in one case reached 96·4 per cent., nearly the whole going into solution with much effervescence.

The British Pharmacopœia directions for making the tincture of *asafœtida* are to extract 200 grammes of the gum-resin with 1 litre of 70 per cent. spirit. Hence, supposing 65 per cent. of the gum were soluble, the tincture should contain 13·5 grammes per 100 c.c. of total solids. Seven tinctures were examined for total solids, and gave results ranging from 4·5 to 8·5 grammes per 100 c.c.

The results of the analyses made are as follows :

Sample.	Extract.	Ash of Gum.	Ash insoluble in HCl.
I. ...	31·9 ...	38·6 ...	82·5
II. ...	30·2 ...	45·8 ...	3·6
III. ...	14·9 ...	62·1 ...	77·8
IV. ...	20·2 ...	56·3 ...	84·3
V. ...	18·4 ...	59·3 ...	74·2
VI. ...	35·4 ...	40·1 ...	64·9
VII. ...	19·9 ...	58·4 ...	74·6
VIII. ...	23·0 ...	52·9 ...	78·2
IX. ...	14·8 ...	63·1 ...	83·9
X. ...	39·8 ...	43·4 ...	42·1
XI. ...	37·7 ...	26·4 ...	67·2
XII. ...	28·9 ...	47·2 ...	78·4

Tinctures.

I.	8·2 grammes per 100 c.c.	V.	5·3 grammes per 100 c.c.
II.	6·2 " "	VI.	4·3 " "
III.	4·5 " "	VII.	8·5 " "
IV.	7·6 " "		

It is, then, evident that commercial samples of the gum-resin of *asafœtida* are very far below the standard given by the British Pharmacopœia, and that they are probably intentionally weighted with worthless material. Consequently, preparations made from the gum-resin according to the British Pharmacopœia directions fall far below the presumed strength, and the medical profession are prescribing medicines of uncertain and variable composition.

A few other observers have noted this condition of commercial gum-resin of *asafœtida*, yet but little attention seems to have been given to their results.

MOERNER (*Journ. Pharm. Chim.*, March 1, 1888, p. 24) examined a sample, and found 86 per cent. of mineral matter, and only 14 per cent. *asafœtida*. A large proportion, on fracture, proved to be alabaster coated with a thin layer of *asafœtida* resin.

BUCKNER (*A. J. P.*, 1890) found the ash in a number of samples to range from 19 to 56 per cent.

LLOYD (*P. J.*, i., 243, 1896) said "ash is often higher than the 10 per cent. required by the British Pharmacopœia." The *National Dispensatory* states that the ash of commercial samples is sometimes as high as 40 per cent.

A thirteenth sample was examined which had been specially prepared by a firm of chemists with a view to determining how far it was practicable attempting to produce a sample approaching British Pharmacopœia strength from this raw material. They succeeded in increasing the matter soluble in 90 per cent. alcohol to 57·5 per cent. (British Pharmacopœia, 65 per cent.). The ash of this was only 5·8 per cent.

Although asafœtida in its various preparations is now prescribed somewhat less than formerly, yet we certainly ought to have reliable material. The standard of the British Pharmacopœia is too high, unless means can be taken to prevent adulterated resin reaching the market. In this connection it may be remembered that a few years ago many samples of ginger were discovered with an abnormally low soluble ash, and that when attention had been drawn to this, and a few prosecutions followed, such ginger almost entirely disappeared from the market.

DISCUSSION.

The PRESIDENT having invited discussion,

Mr. CHATTAWAY said that the British Pharmacopœia, in addition to laying down certain regulations as to the proportion of ash, solubility in alcohol, etc., described the physical characters of the drug; and it would be interesting to know if the samples referred to were in accordance with the Pharmacopœia in this respect.

Mr. E. M. HOLMES said that about a year previously, in his capacity as one of the referees in connection with the compilation of the present Pharmacopœia, he had had analyses made of several samples of asafœtida. He was aware that the commercial form of the drug was largely mixed with mineral matter, but, at the same time, the gum in the form of "tears" was almost pure, and he had experienced no difficulty in obtaining it. When the gum was being collected, a certain number of the tears were obtained pure, except for a small quantity of sand attached to their lower surface, where they fell on the ground; but there was a large quantity of more or less liquid asafœtida, which could not be collected in the ordinary way, and this was mixed by the collectors of the drug with a certain quantity of whatever soil might be obtainable in the neighbourhood in order to solidify it. Sulphate of lime, in the form of alabaster, was probably used as an adulterant, not only for asafœtida, but for benzoin, in which it could be used to represent white tears, a certain proportion of which was normally present in the lump form of the drug. In the samples which had been analysed for him the ash in no case exceeded 8 per cent., but in the crude material in lumps it often ranged from 20 to 60 per cent. He thought that, while for medicinal purposes a drug as pure as possible was desirable, a commercial article was admissible which could be used for drain-testing, or for denaturing tea in the manufacture of caffeine; and a normal standard might be laid down for the crude drug, quite apart from the medicinal point of view. The supply of asafœtida sufficiently pure for medicinal purposes was limited; but there was a fairly abundant supply of the ordinary form, in which a moderate proportion of mineral matter—say, from 20 to 30 per cent.—was for commercial purposes not of much consequence,

seeing that for such purposes the article would be bought on the basis of its percentage of active matter.

Mr. ALLEN said he felt some difficulty in agreeing with Mr. Holmes's suggestion to establish two qualities of the drug. The addition of calcium sulphate in the case of the lump asafœtida was an intentional adulteration, which it would, he thought, be dangerous to countenance. Even though the adulterated material were intended only for commercial purposes, he did not see why the purchaser should be supplied with calcium sulphate when he might imagine that he was getting pure asafœtida.

Mr. CRIBB said that he had found the ash in asafœtida to vary from 15 to 35 per cent., and the proportion soluble in alcohol was correspondingly low. The samples were purchased indiscriminately in London. As far as he could ascertain, the drug in the solid form was hardly ever asked for except in remote country districts, and it was now rarely employed in medicine in any form.

Mr. EKINS said that the drug was very seldom used in the solid form except for veterinary purposes, and the prescriptions in which the tincture appeared were very few.

Mr. MOOR said that the samples he had examined were all in the form of a hard concrete mass, which was the only form of the drug to which his experience extended. Regarding the tincture, which was the chief form in which the drug was used medicinally, it was perfectly easy to make the tincture up to the proper standard by taking a suitable quantity of asafœtida.

Dr. RIDEAL observed that in the Pharmacopœia it was directed that, in making the tincture and fetid spirit of ammonia, a certain quantity of the drug should be taken, and it seemed open to question whether the use of a less pure drug in larger quantity would be in strict conformity with the directions.

Mr. MOOR said that to sell a tincture of as nearly as possible the proper strength—even though it might be made from a larger quantity of the drug than the Pharmacopœia directed—would, he thought, be better than deliberately to sell a weak tincture, and accordingly he would suggest that the tincture itself should be standardized. It would be unreasonable, he thought, too strongly to insist on conformity with the British Pharmacopœia tests, and he would suggest that for the present asafœtida containing not more than 20 per cent. of ash, and not less than from 45 to 50 per cent. of matters soluble in alcohol, should be accepted.

Mr. CHATTAWAY said that the Pharmacopœia laid down that the drug should be in the form of tears—indeed, the actual size of the tears is specified; and he thought that the specimens referred to by the author, although they might be admissible commercially, could not be regarded as representing the British Pharmacopœia article, which frequently contained as little as 2 or 3 per cent. of ash.

Mr. HOLMES said that, while the British Pharmacopœia limit of ash was 10 per cent., a much smaller percentage was quite common in the drop or tear asafœtida. He thought that lump asafœtida would be admissible only provided that it contained less than 10 per cent. of ash. The question of purity was merely a question of the price given for the drug, which varied from about 45s. or 60s. for the lump, to 120s. per cwt. for the finest tears. Seeing that the Pharmacopœia intimated that only the pure drug should be used in medicine, he did not think it would be justifiable to use

the drug in an impure state in medicine, and in its purification some volatile active ingredients might be lost, and the British Pharmacopœia standard of strength would not be maintained.