what is perhaps to be considered remains of an embryonic arrangement in the coccygeal region, or more accurately the result of tardy or arrested development. It is in short a depression or dimple and sometimes a blind tubular canal in the median line a little above the coccyx. Dr. R. M. Hodges called attention to it in the Boston Medical and Surgical Journal, 1880. He had several times found it the centre of an inflammation due to the irritation of hair, which apparently had collected in it. It is mentioned in the Surgical Memoirs of the late Dr. J. Mason Warren, and otherwise had not received notice till a short time before Dr. Hodges' paper. It formed the subject of a communication presented by Dr. Lawson Tait to the British Association in 1878, and published in *Nature* during the same year, who we believe thought it corresponded to the depression above the origin of the tail in dogs. It was written on at considerable length by Ecker, at about this time, in volumes xi., xii., and xiii., of the *Archiv für Anthropologie*. He describes as occurring in this region a whirl of hairs, differing from most others in being convergent instead of divergent.

Supernumerary nipples and mammae have received attention, and the reader is referred to a curious paper on the subject by Bruce, in the *Journal of Anatomy and Physiology*, vol. xiii., 1879, and one by Leichtenstein in Virchow's *Archiv*, Bd. lxxiii., 1878. It appears that this peculiarity is by no means a very rare one, and that it may occur apparently with equal frequency in either sex. This last point, however, is still unsettled.

Accessory Thyroid Bodies.—Kadyi has shown in the *Archiv für Anat. und Entwickl.*, 1879, that extra thyroid bodies are formed in the submental region and even in the floor of the mouth, the result, probably, of separation of portions of the gland at an early period; a fact which we mention on account of its surgical importance. Zucherbandl has published a monograph on the same subject at about the same time.

By way of conclusion to this sketchy and imperfect retrospect we may say that while no great new fact has been brought to light, much good work has been done that has not yet borne its full fruit. Startling discoveries are not to be expected, but the collection of properly observed and classified facts may lead to the enunciation of laws. The most valuable work appears to us to have been done in mechanics, in the brain, and in the collection of anomalies.

Hospital Practice and Clinical Memoranda.

NEW ENGLAND HOSPITAL.

CASES OF LACERATION OF THE CERVIX UTERI OCCURRING IN THE NEW ENGLAND HOSPITAL DURING THE TWO YEARS BEGINNING OCTOBER 1, 1880.

REPORTED BY DR. MARY A. SMITH.¹

In this report are included only such cases as were considered of sufficient gravity to require surgical interference, and not the cases of slight laceration which needed only a few weeks of rest and tonics, combined with local uterine treatment.

During these two years there were forty cases recorded. Of these three finally so far yielded to treatment that an operation was considered unnecessary.

¹ In a paper read before the New England Hospital Medical Society, January 18, 1883.

Altogether there were thirty-nine operations performed upon thirty-seven patients, a secondary operation being necessary in two instances.

SITUATION OF THE INJURY.

On the right side.....	1
On the left side.....	8
Bilateral.....	28
Stellate.....	3
Total.....	40

CHARACTER OF LABORS IN WHICH THE LACERATIONS OCCURRED.

Rapid.....	16
Tedious.....	7
Difficult.....	3
Forceps used.....	8
Unknown.....	6
Total.....	40

In fifteen instances the laceration occurred in a primipara and twenty-five times in a multipara. In regard to these last the date of the injury was so unsatisfactory as to be of little or no value.

General condition before operation: Of the whole number only four might be said to have enjoyed fair health before the operation. As a rule, the usual symptoms in such cases were complained of. Pain in head, back, or extending down the limbs; discomfort on standing, or inability to walk without fatigue; generally some menstrual disorder prevailed, either irregular or painful menstruation or more frequently menorrhagia. Subinvolution and leucorrhœa were generally present, and sometimes the remains of a former cellulitis and subsequent displacement of the uterus were found, where the rent had extended beyond the vaginal junction. Laceration of the perinæum was also a frequent accompaniment.

In all cases some preparatory treatment was given either in the hospital or at our dispensary.

Squibb's ether was the anæsthetic invariably used.

The only antiseptic measures employed were laying the instruments in a one to forty solution of carbolic acid, and wringing the sponges used out of a similar solution. All assistants at an operation were obliged carefully to disinfect the hands.

Silver wire sutures were used in thirty-six operations, catgut in the remaining three.

The after-treatment consisted in keeping the patient in bed, though she was allowed to lie in any position; liquid diet the first week. The catheter was not employed, but the vagina and external genitals were cleansed with carbolyzed water after each micturition. The bowels were confined for the first week, then moved with oil. The stitches were removed on the tenth day, unless menstruation had come on in the mean time, when they were left twelve or fourteen days. In ordinary cases the patient began getting up after two weeks.

RESULTS OF OPERATIONS.

Perfect union.....	33
Partial union.....	4
Secondary operations.....	2
Total.....	39

In three of the cases which failed to unite menstruation recurred in a day or two after the operation, and possibly prevented the healing.

In twenty cases perineorrhaphy was performed either at the same sitting or before the patient was discharged.

In three of these double cases the catgut sutures were used. Twice the union was entire, once only partial.

In only one case was a fever curve required. This patient had, previous to the operation, suffered from pelvic cellulitis, for which she underwent a long course of treatment till the tenderness subsided. She was probably operated upon a little too soon, for a fresh attack of cellulitis supervened, which required ten weeks of after-treatment before the patient could be sent home well.

All the patients were requested to report their subsequent condition. Most of them have since been heard from, and are either steadily improving or entirely restored to health, in many cases the change being very marked.

Reports of Societies.

PATHOLOGICAL SOCIETY OF PHILADELPHIA.¹

C. B. NANCREDE, M. D., RECORDER.

A UNIQUE SPECIMEN OF OSSIFICATION AT THE AORTIC ORIFICE.

EXHIBITED BY DR. J. T. ESKRIDGE.

It was sent him from a distance and consisted of about one inch of the cardiac end of the thoracic aorta, the aortic semi-lunar valves, and the immediate portion of the heart. The specimen was removed from a man, who, aged about seventy, had suffered a number of years from severe heart disease. The walls of the large arteries were thickened, rigid, and contained numerous deposits of inorganic matter. The left ventricle was enormously enlarged. He was unable to obtain any information with regard to the condition of the cardiac valves, other than those of the aortic orifice.

Description of the Specimen.—The aorta, where it surrounds the valves for about half an inch in extent, is a hard, unyielding substance of fibrous tissue, and calcified and ossified matter. The valves in several places are about one fourth of an inch thick, and seem to have been almost entirely transformed into bone-like material. They are rigid and immovable, and have almost completely cut off all communication between the heart and aorta. One of the leaflets, about three fourths of an inch in all directions with its vegetations, stretches across the aorta, lies against, and is apparently adherent to, the other segments of the valves, the latter being curled upon themselves. The central portion of the aorta is entirely occluded, and only two small openings through which the blood could have escaped from the left ventricle are seen between the valvular leaflets near their peripheral attachments. The larger of these holes admits a flattened probe three millimetres wide by one thick; the smaller is about two thirds as large. Three other smaller orifices have existed, but these were obliterated before death by a thin, fibrous, transparent membrane, which is still seen. The valves on the cardiac side are tolerably smooth, but on the aortic side they are very rough, one of the leaflets supporting a vegetation ten millimetres long. One of the segments of the valves is adherent to the inner coat of the aorta for about half an inch in extent, the free end of the valve being folded upon itself, and pointing towards the nearly closed aortic orifice. After macerating the specimen in water

for forty-eight hours, the diseased valves still remain inflexible.

DRS. FORMAD, DUNN, GRIFFITH, and EDWARDS reported having similar or nearly similar cases.

DRS. TYSON and NANCREDE called in question the correctness of calling the disease *ossification* of the valves, as it was in reality a *calcification*.

DR. SHAKESPEARE concurred in this view, and thought that ossification, rarely, if ever, occurred in this situation.

DR. ESKRIDGE said that Hayden² referred to bony deposits in the aorta and its valves as follows: Sir Dominic Corrigan exhibited before the Pathological Society of Dublin³ the heart of a young woman, in which the root of the aorta had undergone complete osteoid transformation; it was likewise greatly dilated, and the aortic valves had been rendered thereby inadequate. During the patient's last illness a systolic murmur of metallic quality, appropriately designated a "trumpet bruit," was audible at the base, and in the ascending aorta and carotid arteries; there was likewise a soft diastolic murmur. He regards a "trumpet bruit" as absolutely diagnostic of bony deposit in the aorta, either in the form of a "rim of bone" or a "projection or tongue of bone." In the same paragraph Corrigan refers to Dr. Banks's specimen of a "tongue of bone" projecting into the aortic orifice.

REPORTS OF THE COMMITTEE ON MORBID GROWTHS.

"A section of the growth presented by DR. NANCREDE for Dr. Seltzer on microscopical examination is shown to consist of an external covering formed of the histological elements and arrangement as found in the skin; the papillæ were in places much atrophied and flattened, also in some parts elongated. There were no hair follicles, sweat or sebaceous glands in the section. Below the papillæ was seen fibrous tissue, at some points in an active state of proliferation, and numerous blood-vessels. The growth is a fibrous polypus."

"A microscopic examination of a section made from the growth removed from the uterus, and presented by DR. PARISH, December 28, 1883, shows it to be adenomatous in structure, consisting of small cavities or spaces lined with cells, which cavities are separated from one another by fibrillar connective tissue."

"A section of the lymphatic gland presented by DR. PARISH, December 28, 1882, shows that its normal structure is metamorphosed into that of a carcinomatous nature, consisting of an alveolar fibrous stroma having the spaces filled with cells of an epitheliomatous type."

Report on Dr. Eskridge's specimen of brain, lung, liver, spleen, kidney, and bladder troubles, presented November 9, 1882: "A section made from one of the nodules of the liver presented by Dr. ESKRIDGE, examined microscopically, shows it to be a new formation consisting of epithelial cells placed in alveolar spaces; the tissue forming the alveolar walls being fibrillar connective tissue. The cells in the spaces at the periphery of the groups, and lying next to the alveolar walls, have a columnar shape, and are quite regularly arranged, while those of the centre are flat or squamous epithelial cells. The neoplasms are cylindrical-celled epithelioma. The post-mortem changes undergone by the brain were such as to prevent any satisfactory histological examination being made."

² Diseases of the Heart and Aorta, vol. ii., p. 839.

³ See Proceedings, vol. ii., New Series February, 1864.

¹ Concluded from page 302.