The Boston Medical and Surgical Journal

TABLE OF CONTENTS

January 22, 1920

ORIGINAL ARTICLES.		EDITORIALS	
PROSTATISM. By Edward L. Keyes, Jr., M.D., New York.	79	COMPULSORY VACCINATION.	96
DIFFICULTIES IN THE DIAGNOSIS OF MENINGITIS. By Harold W. Dana, M.D., Boston.	84	METHYL ALCOHOL POISONING.	97
BLOOD PLASMA CHLORIDES VERSUS RENAL FUNCTION. By W. C. Rappleye, M.D., San Francisco.	89	THE INCREASE OF VENEREAL DISEASE IN GREAT BRITAIN.	97
INSTITUTIONAL CONTROL OF DIPHTHERIA. By Francis A. Finnegan, M.D., Boston.	93	BOSTON MEDICAL LIBRARY.	98
CLINICAL DEPARTMENT		MEDICAL NOTES.	98
TWO CASES OF FOETAL ASPHYXIA WITH UNCOMMON HISTORY. By Charles J. Kickham, M.D., Boston.	94	MISCELLANY	
BOOK REVIEWS		A DISTINGUISHED FRENCH PHYSICIAN.	102
Industrial Nursing. By Florence Swift Wright, R.N.	94	HENRY VIII TH AS A PATRON OF MEDICINE.	103
Military Surgery of the Ear, Nose, and Throat. By Hanau W. Loeb, M.D.	95	EXAMINATIONS FOR APPOINTMENT IN THE MEDICAL CORPS OF THE REGULAR ARMY.	105
A Textbook of Urology in Men, Women, and Children, Including Urinary and Sexual Infections, Urethroscopy and Cystoscopy. By Victor Cox Pedersen, M.D.	95	NOTICES, RECENT DEATHS, ETC.	106

Original Articles.

PROSTATISM.*

BY EDWARD L. KEYES, JR., M.D., NEW YORK.

PROSTATISM is the best word I know to express that condition which results from obstruction to urination at the bladder neck due to adenomatous or sclerotic changes in the glands of the prostate itself, or of the posterior urethra. This description may seem unduly complex to you, but there is even today no absolute and general agreement as to the precise nature and origin of the so-called prostatic hypertrophy or sclerosis which are the two causes of prostatism. Prostatic adenoma, commonly known as hypertrophy, is an adenomatous change affecting various groups of glands about the bladder neck, some of them part of the prostate proper, and some of them small glands lying along the posterior urethra. As a result the prostate enlarges, protruding somewhat toward the rectum, but chiefly toward the bladder into which it projects, generally speaking, in the form of lateral or so-called median lobes or all together as a sort of horse-collar enlargement. The urethra is thus

lengthened and distorted, so that the effort to pass water closes the urethra instead of opening it. Hence the retention, acute or chronic, complete or incomplete, which is the basis of the disease.

The same phenomena of retention result from prostatic sclerosis. The sclerotic prostate may or may not also show adenomatous changes. It is usually but slightly enlarged, it is characteristically a rigid scarred prostate. This condition is appreciated when the surgeon performs a suprapubic section, and introduces his finger into the posterior urethra for the purpose of enucleating the adenomatous portions of the gland. The finger may not enter the posterior urethra at all without tearing this which has been changed into a sclerotic ring. Once in the urethra the mucous membrane tears with considerable difficulty, and the finger finds great difficulty in identifying any plane for the enucleation of lateral or median lobes.

This sclerotic prostatitis may occur at any age after adult life has been reached. I have seen it in men in the twenties, though it is commonest between the ages of 50 and 65. This sclerotic process may result from antecedent gonorrhea of considerable severity, or from bladder stone, or its cause may not appear. The effort to associate adenomatous prostatism

* Read at a meeting of the Surgical Section of the Suffolk District Medical Society, Nov. 21, 1919.

with gonorrhea has completely failed. Cabot and I have independently made observations which show that patients with stricture of the urethra are singularly free in their later lives from the symptoms and signs of adenoma of the prostate. Young has noted the rarity of this condition in Catholic priests, and its frequency in married men. I have been struck by these same facts, but beyond this the etiology of prostatism is open for discussion. Adenoma of the prostate, be it said, shows itself to the pathologist at about the fifth decade of life, but usually does not cause symptoms which persuade the victim to consult a physician until some twenty years later, usually when he is about sixty-five years of age.

Let us assume a working knowledge of the symptoms of prostatism and of the pathological processes which it implies, processes of dilatation and secondary infection of the bladder, the ureters and the kidneys. And let us turn our attention at once to some high lights in the diagnosis of this condition.

Retention of urine in the absence of grave ureteral stricture or of such obvious conditions as abscess in the prostate, stone in the urethra or periurethritis may be due to a number of causes. Let us enumerate them in order of frequency:

1. Adenoma of the prostate.
2. Carcinoma of the prostate.
3. Tabes.
4. Sclerosis of the prostate.
5. Diverticulum of the bladder.
6. Tumor or stone of the bladder.
7. Prostatic stone.

The differential diagnosis between these various conditions is by no means always easy, though sometimes it is obvious enough. Yet it is always confused by the fact that several among them may exist together. Any specialist will recall cases in which adenoma of the prostate has been associated with carcinoma, with tabes, with prostatic stone, with bladder stone, diverticulum or tumor. Thus the differential diagnosis implies, not only the proof that one of these conditions does exist, but that all of the others do or do not exist. So complete a diagnosis is perhaps not always essential to the general practitioner; it certainly is to the specialist.

In the first place one must define retention of urine. This is usually described as a con-

dition in which at least 50 c.c. of urine may be drawn from the bladder by catheter immediately after the patient urinates. Such a definition will suffice in general, but there are a number of functional conditions, notably the diurnal polyuria familiarly but incorrectly labelled hysterical, and the nocturnal polyuria which is often an indication of deficient vitality in a person suffering from chronic nephritis or arterio-sclerosis. There are a number of such polyuric conditions which are not infrequently met with in middle life, and which give symptoms resembling those of prostatism. But their cause is an increased production of urine, not a retention. Let such a patient urinate, then talk with him a few minutes while he is removing his coat, then catheterize him and you may get as much as 200 c.c. from the bladder though the lower urinary organs are actually quite normal and this great quantity of urine has been manufactured within a short time. Let it be understood, therefore, that a single observation of a small amount of residual urine does not establish the diagnosis of prostatism unless supported by many other data.

To enumerate all of the differentiating points between the several conditions that cause or simulate prostatism would but confuse you. Let us rather touch upon the more striking features of each.

Prostatic Adenoma. Prostatic adenoma causes the symptoms of prostatism which bring a patient to consult a physician when he is over fifty-five and not more than seventy years of age. Examination reveals retention of urine and rectal touch usually discloses a bilateral symmetrical enlargement of the prostate in all of its dimensions. Exceptionally the adenoma is at the bladder neck and the prostate feels normal by rectal touch. If a sound or cystoscope is introduced into the urethra this will be felt plainly through the apex of the prostate, while the base of the prostate forms more or less of a mass between the finger in the rectum and the instrument. Inspection of the prostate through the cystoscope reveals a bulging within the bladder quite comparable to the bulging felt by rectum. The demarkation between the lobes is identified by deep clefts in the bladder neck for which Young has devised an ingenious chart. The extent of projection into the bladder is estimated by the degree to which the trigone is obscured.

Carcinoma of the Prostate. Carcinoma of the prostate presents a wholly different picture, most confusing to the inexperienced practitioner. It occurs both earlier and later in life than adenoma, and is the commonest cause of retention of urine beginning in men over seventy. The prostate feels only very slightly irregular in some cases, suggesting ancient gonorrheal infection. But almost always there are areas of hardness in the gland, extending upward around the seminal vesicles, and if felt upon a metal instrument in the urethra this hardness is found to cross the median line at the apex of the gland in sharp contrast to the condition found in adenoma. The cystoscope shows a bladder neck that is irregularly thickened and edematous and possibly ulcerated or showing papillary carcinoma. The most confusing element in the diagnosis is the fact that carcinoma and adenoma may exist together though beginning in different parts of the gland, for it is to be remembered that Geraghty has shown that carcinoma does not begin in portions of the gland which have become adenomatous.

On the other hand, a very large proportion of prostatic carcinomata give as their first symptom something other than urinary disturbance, such as an enlarged gland in the left side of the neck, sciatica (bilateral sciatica is said to be pathognomonic to the prostatic carcinoma) or secondary tumors in bone or lumbar glands. Prostatic carcinoma is often overlooked through failure to make a rectal examination. It may be confused with prostatic stone (see below).

Tabes. Bladder paralysis is not infrequently an early sign of tabes. I have several times seen it in patients whose only other physical sign of the disease was a loss of ankle jerk, and of deep muscle sensibility. These tabetics can often hold their water very well. They show much less vesical irritability than do the prostatites as a class, and by the same token they usually do well on catheter life. The maximum incidence is between the ages of forty and sixty. The prostate is normal, though it may be pronounced enlarged if palpated while the bladder is still full of urine. The trabeculation of bladder wall and laxity of sphincter said to be characteristic of the tabetic bladder are not to be depended upon as diagnostic signs.

Sclerosis of the Prostate. Sclerosis of the

prostate is to be distinguished from paralysis of the bladder with considerable difficulty in some instances. The sclerotic prostate is often not enlarged, it does not project in any characteristic manner into the bladder, indeed the prostate of a tabetic may be sclerotic. Therefore before operating upon such a case blood and spinal fluids should be carefully investigated for the various syphilitic reactions.

Diverticulum of the Bladder. Diverticulum of the bladder is usually congenital, often associated with some sclerosis of the prostate, usually gives symptoms of retention of urine before the forty-fifth year, may be associated with any of the above conditions and is diagnosed by those who look for it with the cystoscope, or by cystography (radiography of the bladder filled with a fluid opaque to the rays).

Bladder Tumor. Bladder tumor may cause retention due to clots, or to the tumor itself blocking the urethral orifice. It is more likely to be mistaken for prostatic carcinoma than for adenoma.

Stone in the Bladder. Stone in the bladder is not an uncommon cause of prostatic sclerosis and certain cases of recurrent bladder stone are curable only by relief of the prostatic retention even though the amount of urine retained be less than 100 c.c.

Prostatic Stone. Prostatic stone is one of the most elusive of the prostatic conditions causing retention of urine. Most cases of prostatic stone show no symptoms whatever. A fair number have a certain bladder irritability. Prostatic sclerosis may result and cause retention. Three times I have seen prostatic stone that by rectal touch gave an irregular hardness of the prostate absolutely characteristic of carcinoma. But the stones are formed chiefly in the line of cleavage about the lateral lobes and the distinguishing feature between carcinoma and stone in the prostate is that stone causes no thickening of the apex. Fortunately the x-ray always discloses prostatic calculi since these are made up largely of calcium salts.

TREATMENT OF PROSTATISM.

The two real problems in the treatment of prostatism are when to operate, and how to operate.

When to Operate. The prolonged preparation of the patient for operation upon the prostate and the prolonged and careful

after-treatment constitute an orderly systematized medical procedure. But the operation itself remains a blind, bloody, brutal, blundering performance,—an emergency operation, if ever there was one. Such being the case, it is small wonder that a very large proportion of physicians and public still look askance upon the operation, and refuse to submit to its dangers. A patient so minded will seek medical aid only when his painful and frequent urination becomes an unbearable agony, or when his urethra shuts up entirely. He then asks for the catheter, and not for operation and is delighted to carry away this novel toilet article rather than to submit himself to the vicissitudes and dangers of operation.

We should not blame him. He is nearing three score and ten, an age at which the preservation of life is not worth much suffering, and if he can manage to slip comfortably downhill in a gradually increasing renal deficiency, or to be snuffed out by an acute focal renal infection he is certainly better off than some of his friends who have been left with suprapubic fistulae or incontinence of urine, though they may outlive him many a year.

Naturally the urologist dissents, and perhaps he exhibits a very moving series of cases to support his contention that prostatectomy is the sole guaranty of eternal life, both physical and sexual. The cunningest way to reach this conclusion is to divide your operative procedure into two parts, kill off all the feeble patients with the first operation, and then brag about the results of the second one.

As a matter of fact, it is probably still the case, as has often been alleged, that in a cross section of the community the general surgeon or practitioner will promptly dispose of about forty or fifty per cent. of the operable prostaties who seek his assistance, while the specialist will reduce this immediate mortality to about ten per cent.

This is not to say that the specialist does the operation any better than the general practitioner. I think I could teach a qualified general surgeon everything I know about the operation of prostatectomy in one brief lesson. A properly prepared prostatic should never die of prostatectomy (I make the statement rather roughly). But some will die in the course of even the best of preparation. Others will die

because of improper preparation, and many will die because of the fantastic improvements that the surgeon introduces into his operation.

But the pleasure of hitting out at the general surgeon is distracting my attention from the subject I wish to discuss. When should the prostatic be operated upon? The answer, of course, is that he should be operated upon just as soon as he shows symptoms of prostatism. When we get patients at that stage of the disease in sufficient numbers we shall be able to develop an operative technic which shall be a little more decent and cleanly than that we now employ. And we shall be able to save these patients' lives in more than ninety per cent. of cases, and not leave them with irreparably damaged kidneys.

What type of operation should we employ? I can best answer this question by distinguishing three types of cases:

1. The frankly adenomatous type with involvement of all the lobe.
2. The adenomatous type with partial involvement, one of the lateral lobes being apparently not enlarged, or with complicating sclerosis, so that one may question whether the obstruction is due to the sclerosis or to the enlargement of the lateral lobes.
3. The frankly sclerotic prostate and the adenomatous bar due to enlargement of Albarán's glands.

The frankly adenomatous prostate is, I believe curable by the ordinary suprapubic prostatectomy on condition that the mucous membrane of the bladder neck is spared at the superior commissure, and that the fistula is left at a point as distant as possible from the bladder neck. We spare the mucosa so as to prevent postoperative stricture of the bladder neck. We place the fistula high so as to encourage its early closure.*

I did my first prostatectomy on a case of this type in 1898. I have yet to see a recurrence, though this operation does not cure pyonephrosis, renal or bladder calculus, or bladder diverticulum or tumor.

The intermediate class of cases that we may call partial hypertrophy or adenomato-sclerotic cases must be operated upon according to the same principle. If, for example, one of the lateral lobes seems not to be enlarged the finger must nevertheless be introduced into the

* I have no intention of interrupting our evening with discussion of the best methods of operating or of preparing the patient for operation.

urethra, the mucosa torn open over this lobe and such fragments of it as can be enucleated should be torn away. The procedure may seem very futile at the time, but if it is not carried out the patient is doomed to a relapse. Under the best circumstances he will be relieved of his symptoms from four to ten years and then they will recur, due to the adenomatous enlargement of the lobe which was not enucleated.

I have apparently permanently relieved the retention of a number of cases of the adenomato-sclerotic type, by removing absurdly small fragments of the lateral lobes—lobes which did not project into the bladder and whose adenomatous condition could only be appreciated either by the urethroscope or by the finger introduced into the prostatic urethra through the suprapubic wound. The presence of sclerosis, however, makes one fear that these cases will relapse after a certain number of years have gone by.

The problem of the frankly sclerotic prostate is with us still. Bottini, Chetwood, Young, Goldschmidt, Bugbee and Buerger have devised various types of operation, any of which will temporarily relieve a certain proportion of cases.

The most brilliant example of the failure of these procedures that I have encountered is the following case. The patient came to me having had a Buerger operation performed for an obstruction at the bladder neck, which I understand was a small median lobe. This operation did not relieve the chronic complete destruction to urination from which he suffered. He entered Bellevue Hospital last spring, and I performed a suprapubic operation upon him and discovered the bladder neck to be very tightly strictured. I tore open this stricture, and enucleated two very slightly enlarged and sclerotic lateral lobes. This left him with a retention of about 200 c.c. and an open suprapubic fistula. I used the Young punch upon him twice, obtaining no results the first time, and a reduction of his residual urine to 100 c.c. the second time, but within six weeks it had returned to 200 c.c. I then did a Bottini operation upon him (the only one, by the way, that I have performed), and his fistula closed, and his residual urine dropped to about 30 c.c. when he urinates sitting down; but if he attempts to urinate standing up he still has 200 c.c. residual urine. In this condition he has re-

mained for five months, and the amount of his residual urine will doubtless increase.

Fortunately, not every sclerotic prostate behaves quite as badly as this. Slight enlargement of Albarran's glands and sclerotic prostates may be successfully operated upon by the Young punch, and the author of this operation has reported cases that remained well for a number of years. I have reported equally satisfactory results from the Chetwood operation, but only two of my cases have remained well for ten years. One, a purely sclerotic prostate, is apparently well twelve years after operation; the other has now a very large adenomatous prostate, but empties his bladder completely although his operation was performed in 1905. Such accidents may happen.

What, then, is the best operation for the sclerotic prostate, the contracted bladder neck, the slight enlargement of Albarran's glands? To this question I do not know a universal answer. One of my patients remains well twelve years after a tearing open of his bladder neck, thanks to an occasional treatment with the Kollmann dilator. Another died recently of renal calculus, an attempt at suprapubic prostatectomy having been made upon him by myself and by another surgeon. Neither of these operations did him the least good. The residual urine of about 150 c.c., which he had when he came to me six years ago, and which was the occasion of his bladder and kidney stones, remained unchanged by the two operations. At postmortem I was shocked to find that his bladder showed no diverticula and the prostatic orifice was so wide open that the tip of the little finger could be introduced into it scarcely touching its sides, yet his residuum had been undiminished.

Dr. Squier has recently suggested that for such cases the proper operation may be the total removal of the prostate, dividing the urethra at the bladder neck and at the membranous urethra and sewing the two together. In two or three cases his immediate results following such an operation appear to have been excellent.

It seems a pity to close one's remarks in such a pessimistic strain. We can but cheer ourselves with the hope that the next generation of surgeons will succeed where we have failed.