X.—*On a Hydride of Boron (Preliminary Notice).*

By FRANCIS JONES, F.R.S.E.

I AM not aware that any attempt has been made since 1809 to obtain boron hyhdride. In that year Davy stated that a gas, which he supposed to be a compound of boron and hydrogen, was evolved by the action of water on potassium boride; and he obtained a similar gas by the action of acids on a boride of iron prepared by heating a mixture of boracic acid and iron filings to a high temperature. These results have been attributed to impurities in the substances employed, and it is generally stated that boron is the only non-metallic element which forms no compound with hydrogen.

Thinking it probable that a hydride might be obtained by other

42 L. T. WRIGHT ON THE OCCURRENCE OF CERTAIN NITROGEN

reactions, I attempted in the first place, to obtain magnesium boride by the action of sodium on a mixture of magnesium chloride and potassium borofluoride, thus imitating the process employed for the preparation of silicon hydride. A very violent reaction occurred on simply mixing the substances, but no successful result was obtained on adding hydrochloric acid to the product.

It then occurred to me that magnesium boride might be obtained more simply, by strongly heating a mixture of magnesium dust and boron trioxide. On trying the experiment in the proportions suggested by the equation—

I found the action succeeded perfectly, and a grey friable mass was obtained, containing magnesium boride. When hydrochloric or nitric acid is dropped on this substance, great heat is given out, and an inflammable gas is evolved, which is without doubt a hydride of boron. On repeating the experiment on a larger scale, I obtained the gas in sufficient quantity to make the following observations. It is a colourless gas, burns with a bright green flame, and has a very disagreeable odour, resembling both that of phosphine and that of stibine. It is soluble to a slight extent in water; the solution gives with silver nitrate a brownish-black precipitate, containing both boron and silver. The gas is decomposed by passing it through a red-hot tube; but I have not yet obtained it unmixed with hydrogen.

The reaction of magnesium on boron trioxide also serves very conveniently for the preparation of boron itself. Half the quantity of magnesium represented in the above equation is used, and the product when extracted with hydrochloric acid, leaves amorphous boron as a brown powder.