there is a large amount of needless repetition, for the estimation of the heat of combustion is a simple calculation, which might have been made once for all with reference to each proximate principle, especially since the bare facts, as they are put, convey but little idea to the general reader. The chemistry of foods is very superficially and imperfectly treated, not nearly so full as it deserves; and the botany would have been better if a more thorough study of *materia medica* had been undertaken. There is one sentence we have in vain attempted to understand. When speaking of the sweet chestnut, the author, after remarking that at present it may be regarded as a luxury, says, "The first step to a great extension of its use would be to make the ordinary horse-chestnut a safe and agreeable food, since it grows in our climate, and could be obtained in large quantities." How this can be, seems extremely difficult to understand; as is well known, the two fruits having nothing whatever to do with one another.

The descriptions of the various methods that have been proposed for the preservation of meats which have to travel long distances and through hot climates is very complete and clear. The preference is given to the method of heating, and that adopted by Mr. Jones, in which the meat is heated in vacuo, to 280° F., in the cans, is fully described. It is shown, however, that by this process the meat is stewed, and over-stewed, not roasted nor boiled. In this, and all similar processes, it is found impossible to expel all the air without over-cooking the meat.

Another subject of particular interest which is discussed is the preservation of milk. Two methods, it appears, are adopted in America, one in which the milk is simply evaporated to one-fourth its original volume, when it will often keep for a month, and another in which sugar is added; by the latter process it remains good for an indefinite time, and contains about one-third of its weight of sugar. The author agrees sufficiently with Dr. Daly in his condemnation of the employment of this preserved milk for infants, to quote an article by him which appeared last year in the *Lancet*.

Extract of meat, especially Liebig's, occupies the greater part of one chapter, and we think the author has done good service in setting in a clear and unmistakable light the true value of that expensive luxury. He shows that its chief value depends on the meaty flavour it is capable of imparting, and that its nutritive value is *nil*. He remarks—"Its proper position in dietetics is somewhat more than that of a meat-flavourer, but all that is required for nutrition should be added to it. . . . Used alone for beef-tea it is a delusion." That this is correct is evident from a consideration of the method by which it is prepared, for "during the process, all the fat and as much of the gelatin and albumen as can be extracted are removed from the solution of flesh, whilst the fibrin, being insoluble, is necessarily left behind. Hence there remain water, salts, osmazone, and the extractives of flesh, or, in general terms, the flavouring matters and the salts of meat—thus leaving out all that is popularly (and correctly) regarded as nutritious."

Many tables are given to show the effects of different substances on the respiration, pulse, exhalation of carbonic anhydride and aqueous vapour. There seems to be a

want of association between the great mass of facts, which must have been the result of long and continuous labour; and they are undoubtedly put forward in a way which is not best suited to convince the scientific student. For example, the effects on the pulse, &c., of tea dissolved in water is given in full, but under the head of water no mention is made of its physiological action, though decidedly, by itself it changes the pulse rate, if nothing else.

Several recipes of the fourteenth century are quoted from "Cury," a copy of ancient manuscript recipes of the master cook of Richard the Second. There are also many scriptural references, and a very inappropriate abstract of an incident which occurred at the Worship Street Police Court.

OUR BOOK SHELF

A Manual of Metallurgy. By George Hogarth Makins, M.R.C.S., F.C.S., &c. (Ellis and White, 1873.)

THE present edition of this work presents a marked improvement over those which have preceded it, but it is still far from being all that even a small manual might be. In the preface the author expresses a hope that the volume, "in which the leading points connected with the principal metals are set forth, may be found useful," and as there are singularly few metallurgical works in the English language, we have but little doubt that this hope will be realised. Mr. Makins has long enjoyed the reputation of being a most accurate assayer, and the descriptions of the processes of assaying gold and silver are careful and valuable. The portion of the work which is the least satisfactory is that devoted to iron.

LETTERS TO THE EDITOR

[The Editor does not hold himself responsible for opinions expressed by his correspondents. No notice is taken of anonymous communications.]

The Huemul

THE Huemul of Chili and Patagonia, referred to in *NATURE*, pp. 253 and 263, was first recognised in modern scientific literature by MM. Gay and Gervais, who in the *Annales des Sciences Naturelles* for 1846 (v., p. 91), showed that the so called *Equus bisulcus* of Molina, was a species of Deer (*Cervus*), which they proposed to call *Cervus chilensis*.

In Gay's "Fauna Chilena" (plates 10 and 11), the female and skull are figured. Concerning the nomenclature of species, I have published some remarks in the last volume of the "Annals of Natural History" (ser. 4, vol. vi. p. 213), to which I beg leave to refer such of your readers as are desirous of further information on this subject. P. L. SCLATER

11, Hanover Square, W., Aug. 6

Perception and Instinct in the Lower Animals

IN answer to Mr. George J. Romanes (*NATURE*, August 7) I beg to say that I particularly inquired of my friend whether he had been to or near his old house on the day the dog returned, or shortly before, and he assured me that "he had never been near it since he left." I ought to have stated this in my account of the circumstance.

I shall make no further remarks on the subject, because I believe that nothing satisfactory can be arrived at till experiments of the nature indicated in my last letter have been systematically carried out. ALFRED R. WALLACE

Collective Instinct

THE writer of one of the books on Indian sport relates how he saw a herd of antelopes, driven backwards and forwards by four wolves, which surrounded the herd, each guarding a diffe-

rent side, until at length the antelopes passed over a ditch in which a fifth wolf lay concealed. This wolf, jumping up as the antelopes crossed, secured one of them, upon which his four companions joined him, and assisted in making a meal of the captured animal.

A civilian of the N.W.P.* told me that he witnessed a very similar occurrence in Oudh. He saw two wolves standing together, and shortly after noticing them was surprised to see one of them lie down in a ditch, and the other walk away over the open plain. He watched the latter, which deliberately went to the far side of a herd of antelopes standing in the plain, and drove them, as a sheep dog would a flock of sheep, to the very spot where his companion lay in ambush. As the antelopes crossed the ditch, the concealed wolf jumped up, as in the former case, seized a doe, and was joined by his colleague.

Here are two well-authenticated instances of an action or series of actions requiring the exercise of combined sagacity of a high degree on the part of two or more individual animals, being performed in exactly the same way by different members of the same species. Was the method employed by the wolves to secure their food, which they could not have caught single-handed, the result of separate experience or of inherited habit? The identical character of the stratagem employed in the two cases points to the latter.

I have noticed some similar instances of collective action on the part of other animals which I believe to be as much inherited as the habitual actions of individual animals. I have constantly seen a flock of pelicans when on the feed form a line across a lake, and drive the fish before them up its whole length, just as fishermen would with a net. The capture of the fish is rendered doubly easy by this method. I have witnessed exactly a similar plan pursued by a large number of Ganges crocodiles which had been lying or swimming about all day in front of my tent, at the mouth of a small stream which led from some large inland lakes to the Ganges. Towards dusk, at the same moment, every one of them left the bank on which they were lying, or the deep water in which they were swimming, and formed line across the stream, which was about twenty yards wide. They had to form a double line, as there was not room for all in a single line. They then swam slowly up the shallow stream, driving the fish before them, and I saw two or three fish caught before they disappeared.

Where a large number of individuals constantly repeat in continuation the same action, it is possible that the younger members may merely copy the older members of the species, and so carry on the habit generation after generation. This is less likely where few are concerned, as in the case of the wolves. A pair of wolves are probably of the same age. It is a marked habit of some species of birds to hunt in pairs, and assist each other in the capture of their prey. The *wokhab*, or common eagles of the Indian plains, hunt in this way. When one of the pair misses in its swoop, the other descends on the victim before it has time to make a fresh attempt to escape. The circumstance that some species of birds of prey are in the habit of combining for the capture of their food, while others hunt singly, would tend to prove that the combined habit is as much inherited as the habits of individuals are known to be.

Gregarious actions, which require combination of purpose on the part of two or more individuals, entail the exercise, if not of a higher degree of intelligence, at any rate of a greater number of intelligent qualities than the isolated actions of single individuals. This class of actions possesses, therefore, a special interest. Those instances in which different individuals perform totally different acts for the attainment of the same end, as in the case of the wolves, are the most interesting, as requiring the most intelligent qualities. I should be glad to learn if any of your readers have ever witnessed or heard of the stratagem described above being employed by wolves for the capture of their prey.

Allahabad, June 29

E. C. BUCK

Ants and "the Taint of the Hand"

IN NATURE, July 24, Mr. James D. Hague, writing on the habits of ants, attributes their dislike to the place across which a finger has been drawn to "the taint of the hand."

Now, Sir, I have frequently drawn a line with a piece of chalk across the track of ants, and observed in them the same symptoms of dislike as Mr. Hague's ants showed to the finger-mark.

* Mr. Elliott, B.C.S., now Secretary to Government, N.W.P.

I have also drawn a small circle with chalk round one or more ants, who will seek a spot untouched by the chalk through which to make their escape; but should there be no such opening, they will presently cross the circle. If, however, this enclosure be made upon a perpendicular wall, &c., they will frequently drop to the ground rather than walk across the line.

Now, as I have never observed this same dislike—exhibited by dropping—of the "taint" when ants have been running over my hands, and as the chalk-line has the same effect as the finger-mark, may it not be something else than the "taint of the hand" to which the ants object when their usual track is interfered with?

Stamford, Aug. 8

G. E. C.

Venomous Caterpillars

WITH reference to a paper published by Mr. Murray in NATURE, vol. viii. p. 7, on Venomous Caterpillars, I wish, in corroboration, to add my testimony from personal experience, that a species of caterpillar has the power of inflicting a very painful sensation (I will not say wound, as such was not visible) by its sting.

In 1868, when travelling in company with Capt. Street in the Burmese forests on a botanical trip, and whilst in the act of detaching a specimen plant of *Deudrobium farnerii*, from the naked branch of a tree, I felt a severe and painful sting on my thumb. On examination I noticed I had seized hold of a large caterpillar lodged amongst the roots of this orchid. It was about two inches long, clothed with erect hairs; its colour was a reddish brown, the lower part of the abdomen being darker, with well-developed legs.

My thumb continued painful for three days; it was considerably swollen, the skin having a drawn glazed appearance.

The Burmese told me that this kind of caterpillar was exceedingly venomous, and one fellow was particularly consoling by informing me that unless the pain subsided in three days the sting might prove fatal. I am inclined to think that the caterpillar for self-protection has the power of detaching these hairs; whether any propelling force is present at the time of detachment it would be difficult to prove.

I found steeping my thumb in Eau de Cologne gave me the greatest relief.

Whether these hairy caterpillars have a special venom or otherwise I do not feel qualified to express my opinion either one way or the other; but I lean towards the conclusion that the irritation is set up by the mechanical action of the spine during its penetration of the skin, and my reason for inclining towards this opinion is because we have a somewhat parallel case in the irritation caused by the hairs of the prickly pear.

I was present when an officer was thrown off his horse into a prickly pear hedge; he suffered the greatest pain, and could not bear the parts, where these minute spines had penetrated the skin, to be touched. On his being placed in a warm bath the relief was almost immediate, especially to those parts capable of total immersion, and this I attribute to the prickles or hairs floating and becoming removed from the skin by the oscillatory motion of the water.

Madras, July

R. BENSON

Abnormal Ox-eye Daisy

IN 1868 I gathered among the ruins of Pompeii a very curious monstrosity of the common ox-eye daisy. The flower and flower-stalk were confounded into a strap-shaped mass which was fringed with the florets. I showed it to Prof. Wyville Thomson, who told me it was an instance abnormal in this species, of the form of inflorescence which is normal in the coxcomb.

JOSEPH JOHN MURPHY

Old Forge, Dunmurry, Aug. 1

Canarese Snakes

FAM. *Erycida*, Gen. *Gonygophis*? Sp.?—Captured in Mangalore, December 2. Gape wide; fangs in sup. and inf. maxillaries.

Body moderate, tail short, obtuse scales, smooth, 48;—ventrals narrow, 197, terminating with three rows of scales between last ventral and anal; latter entire. Subcaudals single, 24, last forming conical point.