Thrombosis, carrying with it the possibility of embolism, may develop after any type of operation, but the element of risk seems greater in the case of operations in the upper abdomen than in those involving the pelvic viscera. Post-operative pulmonary embolism usually occurs between the eighth and twentieth days of convalescence. In the majority of cases, fortunately, the embolism is not fatal. When it involves a small vessel or when the embolism is lodged in an organ where vascular anastomosis is free, embolism is often unrecognized. In an analysis of 16 fatal cases, Busch found that death was instantaneous in one-fourth. In the others, ten minutes or more elapsed before death took place. Novak discusses the various factors to which an etiological role has been attributed.

A Statistical Study in Serology. By Robt. S. Preston, Richmond, Va. *The Old Dominion Journal of Medicine and Surgery*, April, 1915, pp. 202-205.

The author in his study states that he has found it essential to the correct interpretation of incomplete complement fixation tests that the clinical and laboratory finding be correlated. Incomplete Wassermann reactions and indefinite luetin tests are as apt to be overestimated as are negative Wassermanns. The latter does not exclude the possibility of a luetic infection. To do this to the best of our ability several tests should be performed at intervals, and if necessary after a provocative injection of salvarsan has been administered. Salvarsan is more efficient in producing the Herxheimer phenomenon than mercury, though mercury may be utilized previously for this purpose on account of the facility of its administration.

Of 11 cases that had received antisppecific treatment 41 per cent gave a positive Wassermann reaction. Those cases (368) diagnosed and grouped as being syphilitic in nature gave 79 per cent positive reactions. Two hundred and forty-four cases treated or untreated diagnosed as primary, secondary and tertiary lues gave 90 per cent positive tests.

In the routine cases for diagnosis one or two cases of paroxysmal hemo-globinuria gave a positive Wassermann, as also did a case of hematomphrynia, one case that had previously had a splenectomy performed, one case clinically a gastric carcinoma, and one with gastric crisis operated on for perforating ulcer. Of seventeen cases of aortic diseases, six gave positive tests. Of three cases of Menier's diseases partial fixation was obtained in two, but no response could be obtained by treatment. Of eight cases of syphilis of lungs and pleurae, two were diagnosed pleurisy and four were known to have been considered phthisis previously.

The Significance of Bacillus Coli in Pasteurized Milk. By L. P. Shippen, Baltimore, Md. *Journal of the American Medical Association*, April 17th, 1915, pp. 1289-1291.

The presence of *B. coli* in pasteurized milk is commonly taken to indicate either improper pasteurization or subsequent contamination of the

milk. This organism was supposed to have a thermal death point below the temperature used in pasteurization, and in consequence it was believed that it could not survive this process. However, De Jong and De Graef claim to have isolated strains which are capable of withstanding a temperature of 149 to 152.6 degrees F. for thirty minutes. These strains would not be killed by the degree of heat usually applied in the process of pasteurization, and their presence would not mean that the milk had been improperly heated.

The writer succeeded in isolating from pasteurized milk seven strains of *B. coli* which survived temperatures ranging from 143.6 degrees F. to 147.2 degrees F., applied for thirty minutes. This confirms the work of De Jong and De Graef, and indicates that the presence of *B. coli* does not mean necessarily improper pasteurization or subsequent contamination of milk.

Diuretics in Renal Diseases. By Leslie Byron Wiggs, Richmond, Va. *The Virginia Medical Semi-Monthly*, April 9th, 1915, pp. 7-10.

The author protests against the promiscuous use of diuretics in the presence of renal damage. He supports his views by the results of experiments on rabbits in which he had produced fatal acute nephritis. The use of diuretics in these animals in each instance shortened their lives. He believes that if this is true in the acute conditions that it is equally true in the chronic types, for in both instances the cause of the nephritis is irritation, and he argues that it is illogical to further irritate them by the use of diuretics, for "cells that fail to functionate as a result of stimulation or irritation will probably not be permanently benefited by further stimulation or irritation." What they need under such circumstances is rest, and his clinical experience justifies such an hypothesis.

He summarizes as follows:

1. In chronic cases of nephritis with rising blood pressure and a normal chloride and watery content in urine associated with a low urinary urea, diuretics of the vascular type, may be used with caution.

2. In severe acute cases diuretics are contraindicated.

3. In cases definitely known to be mild or subacute in type, less irritant diuretics may be used guardedly. . . . Since, clinically, these types are difficult to differentiate from the early manifestations of severe forms, it seems a good rule to interdict the use of diuretics in doubtful cases.

4. Theoretically, there seem to be few logical reasons for the use of renal stimulants or irritants; therefore their indications in any case are clothed in extreme doubt.

Bell's Palsy and Neuritis of the Eighth and Twelfth Nerves Following Salvarsan Injection. By Virgilus Dabney, Washington, D. C. *Interstate Medical Journal*, May, 1915, pp. 478-480.

Foreign observers have reported deaths and cases of neuritis but such reports are not common with us, whatever be the motives. This has served to lead many into the use of salvarsan through lack of warning as to the possibility of