ounces of serum. The arachnoid itself was quite healthy. The brain, which was lying collapsed at the base of the skull, was much more vascular than natural, but its consistence was normal. The marks of the convolutions were quite distinct. The lateral ventricles were empty, and enormously dilated; their lining membrane was thickened, but smooth. On one side a small pellucid cyst, the size of a grape, was attached to the inner wall of the lateral ventricle by a small vascular pedicle. The cerebellum, the cerebral nerves, and other structures at the base of the brain, were in a natural condition. No fluid was discovered in the spinal canal. The opening made by the trocar through the right lateral ventricle had evidently become wider since the operation; the edges of the aperture, nevertheless, presented a smooth and healthy appearance.

Remarks.—Although the above case was only partially successful, it has been the means of eliciting some important facts, which may serve to guide us in the future treatment of a melancholy and intractable disease. It has shown that iodine may be injected into the brain without causing any poisonous effects, and that its strength may be so regulated as to produce an amount of stimulation sufficient to restore the balance between the secreting and absorbing functions, without causing destructive inflammation of the cerebral tissues.

In the first case, in which I merely tapped the brain, without employing an injection, the water began to re-accumulate in the head on the third day; whereas in the latter instance, where iodine was injected, there was no perceptible increase of fluid on the tenth day after the operation. The small quantity of fluid found in the head after death was most probably part of the original serum contained in the brain before the operation, as I had purposely refrained from evacuating every portion of the serum.

Congenital hydrops cerebri is essentially a non-inflammatory disease, and the fear of exciting inflammation in these cases has led to the neglect of efficient measures. The brain is in a debilitated condition, and requires careful stimulation.

The details of the case which I have narrated prove that the shock of a severe operation on the brain is not necessarily fatal. The little patient recovered from the shock, and the principal functions of the body were restored. Had the child resided in a healthy locality, and possessed more stamina, the results of the case would probably have been more encouraging.

Finsbury-square, October, 1855.

ON SOME OF THE PHENOMENA OF VISION.

By WILLIAM HENDRY, Esq., M.R.C.S., Hull.

DURING the month of July, 1851, an article appeared in the pages of THE LANCET, from the pen of Mr. Rainey, of St. Thomas's Hospital, ascribing to the ciliary processes of the eye the function of adaptation to distinct vision of near as well as remote objects; to which, in a subsequent communication, I took occasion of asserting my own long-cherished convictions, that the delicacy of the structures in question (ciliary processes) precluded all possibility of their adequacy even to antagonise, much less to exert a preponderating influence over the ceaseless and powerful activity of the larger muscles implanted upon the exterior of the globe,—considerations which induce me to withhold assent to the popular theories, ascribing to any of the interior structures, thus so feebly developed, so important and essential a function.

But although endeavouring to refute such hypotheses, relative to the capacity of the combined efforts of the numerous ciliary processes, in the longitudinal displacement of the crystalline lens, yet, considering the lamellated character of the tissues involved, and the motions of such lamellæ being transverse to the optical axis, I am willing to allow what I conceive to be a proper function of such processes, as constituting one amidst the several agencies by which the disturbing impulses of exterior muscular action are rectified, the rectilinear adjustment of such axes being further aided by the elasticity of other structures (the sclerotic), and also by the restored equilibrium of muscular tension; the degree of tension alone determining every alteration of focus.

Views thus entertained might be deemed mere assumption, and consequently fallacious and justly demanding still further evidences.

I would hence refer to a close attention and study of the varied developments and phenomena of what have long been understood by *muscæ volitantes*, for a representation and description of which I have great satisfaction in recommending

the perusal of page 341 of Wharton Jones's Manual on Diseases of the Eye. *Muscæ* will be found to exhibit isolated vesicles, or others in linear arrangements, like strings of beads; others enclosed in tubules in single file, approximating, or in a greater or less degree apart; or tubules straight or otherwise spiral, and without vesicles contained within: vesicles also with and without nuclei; other vesicles enclosed in one or even two concentric margins; vesicles clustered or aggregated, or under other arrangement, forming beauteous figures or constellæ; while, under a more rigid examination, the entire field of vision presents a basement of non-nucleated vesicles, to such an extent as to render it probable that the whole solid structures seen are thus resolvable.

Travers long since published that "these bodies present themselves to most persons at some period or other during life;" and so far as my own investigations extend, they may be seen by every one using necessary endeavours, under a proper modification of light, whence it may be reasonably inferred that they constitute the normal condition of the structures in which they are seen, and if proving at any time a source of annoyance, do so through a morbid sensibility, induced through gastric, cerebral, or some other cause of disturbance.

I do not, however, regard any of the vesicles or clusters as truly floating, as considered by some authorities, or even so detached as to move with freedom and confusion in a chambered fluid of diminished density to themselves; but that all, of whatever description, whether apparently isolated or entering into complex groupings, are attached to or otherwise enter into the structure of the solid tissues wherein seen; so that under circumstances of observed movements, whatever oscillations or intersections of different groups, or approximations or glidings may occur, such are not indicative of the confused motions of stray fragments, varying in the degree and direction of their course according to impulse received, and the relative gravity and magnitude of the particles, but movements corresponding to the entire plane to which each visible series properly belongs, so that in many instances they are seen through several planes deep, the axilinear aggregations amidst which, in some cases, accounting for occasional temporary opacity.

Such has been the constant result of my own observations for many years past, and which may account for my enforcing these views of a lateral deflexion of the axis of vision, requiring a mechanism of adjustment, which I am disposed chiefly to ascribe to the ciliary processes.

As a passing remark, I believe these bodies we ordinarily term *muscæ* constitute that ultimate organization which fulfils the vital functions of the more simple and diffused structures in secretion and absorption beyond the confines of ordinarily demonstrated vascular distribution, with other interspersions of nerve vesicles and tubules, not necessarily belonging to these of common sensation.

When an individual looks with considerable attention, as for instance, at some one point of small print, (supposing him to have full command over the observation of *muscæ*.) then, on a sudden after a moment's elapse, turns attention to some other distant point on the same page, he will observe an interval of time before such latter point becomes distinctly defined. Such observations enable one to ascribe the delay to oscillation of the structures of the eye itself, occasioned by muscular directive impulse; and I apprehend that in advanced life, with declining integrity of muscular contractility, so also may the impaired vision of old age be in part imputed to the vacillatory movements of the muscular fibres of the globe, both external and internal, irrespective of the usually recognised alterations in the curve of the lenses.

The groupings or configurations of *muscæ*, are dissimilar in the two eyes of the same individual; so that besides the laminated movements observable in either, the more complicated motions discernible in both at the same moment, and their ever-altering relations to each other under muscular disturbance, meanwhile in a state of quiescence falling back to their normal position, only confirms, in greater degree, what has hitherto been advanced under separation observation (*id est*) of each eye alone; and I am still further prepared to state, that not only, as just observed, are the groupings dissimilar in the two eyes of the same individual, but the foci and visual powers, even in a normal condition, oftentimes, if not for the most part, as widely differ in each also.

One may pause with astonishment at the precision of an article contained in the *Athenæum* of September 21st, 1839. When referring to a paper read at the British Association of that year relating to "personal equations" as needful to be allowed to reconcile existing differences in time, noted by two or more observers of the instant of chronometric indications,

or of the appearance of signal rockets employed in the measurement of intervening distance between two or more stations in geographical survey; but such precision falls into comparative insignificance when experience testifies to the fact, that even in the same individual *an equation* also is necessary to reconcile ocular indications of each eye separately; hence, when the medical attendant may be consulted relative to the using lenses, as in double eye-glasses, he should be aware of the not unfrequent occurrence, wherein a different focus may be required for each eye, and that glasses (spectacles), as a constant rule, should not be exactly paired in numerical and optical accordance, but that each eye be suited to its own actual requirement. Thus, also, most serious errors are committed in the construction of our double opera-glasses possessing connected sliding tubes, instead of a detachment, enabling the most perfect adjustment of focus to each eye separately.

At the present crisis, when so pressing are the demands for medical gentlemen in the navy, and when our medical services have become an essential accompaniment to almost every scientific expedition, wherein the telescope and various astronomical instruments are extensively employed, questions of a perplexing character may be occasionally submitted, but little contemplated, and such as have occurred to myself not unfrequently, thus: Why, in observed lunar distances, does the index of a repeating circle or sextant obtain more, by four or five seconds, with the left eye than with the right, and *vice versa*? The reply suggests itself, that the two halves of the human body are seldom precisely equal in development and function; the right hand mostly exceeds that of the left, as in the fitting of gloves; the left foot mostly that of the right, as in the fitting of boots; the meatus and osseous canals of the ears are oftentimes of different calibre, the listener unconsciously turning that auditory organ in the direction of sound which is the most exquisitely sensitive; and so with the eye there exists a predominating sensibility or power of one over that of the other, which although of trifling import perhaps in the ordinary engagements of life, yet, under more particular pursuits, may prove of more serious consideration—important, doubtless, when, under neglected adjustments in the use of glasses or otherwise, one eye may be found to be performing an undue amount of labour, while the other is merely occupied in the dim and profitless gaze: important, also, under other considerations, as when a justly needed equation, as that of several seconds of an arc, in angular measurement, being neglected in its application to other elements of a geometrical problem, occasioning discrepancy of a nature and magnitude impossible otherwise to reconcile, neither ascribable to tabular sources of error, nor to accidental omission of instrumental adjustment, but exclusively to individual peculiarity in the development and function of a special organization, as in the present argument—viz., that pertaining to vision.

October, 1855.

A Mirror OF THE PRACTICE OF MEDICINE AND SURGERY IN THE HOSPITALS OF LONDON.

Nulla est alia pro certo noscendi via, nisi quam plurimas et morborum et dissectionum historias, tam aliorum proprias, collectas habere et inter se comparare.—MORGAGNI. *De Sed. et Caus. Morb.* lib. 14. Proœmium.

ROYAL OPHTHALMIC HOSPITAL.

SOME useful hints are occasionally gleaned by comparing the results of practice throughout the year at the various Ophthalmic Hospitals. Having met M. Worlomot, of Brussels, frequently at Moorfields, we can confirm his impression, "that if the London operators do not perform better than their Belgian brethren, they certainly operate differently," and we would almost add, no two alike. We have been, also, very much struck with the differences of opinion, even amongst our best ophthalmic surgeons, on such familiar, if not commonplace, points as the treatment of purulent ophthalmia, and that of strumous ophthalmia, the use of the ophthalmoscope, &c. At St. Thomas's Hospital, Mr. Mackmurdo uses very generally, in ophthalmia cases, a solution of one grain of nitrate

of silver to an ounce of water, as a lotion. Mr. Wharton Jones and Mr. Guthrie, we have observed, on the other hand, in similar cases, find a strong ten-grain solution preferable. A favourite ointment with the former surgeon is, a drachm of levigated red precipitate of mercury to an ounce of lard, in place of the nitrate-of-silver ointment, which we have seen used elsewhere; at University College Hospital the granulations on the lids are freely scarified with a sharp, round-pointed lancet, and the red precipitate ointment freely applied, and we must say, with most excellent effect. Dr. Taylor and Mr. Haynes Walton, at the Central London Ophthalmic, are more in favour of solution of sulphate of alum, as at once more economical, more effectual, and not injuring or discolouring the eye. The change effected by nitrate of silver, especially in soldiers dismissed from the army, where it is quite a panacea in ophthalmia cases, being most marked in the production of troublesome pannus, dyeing or staining of the conjunctiva, &c., (the latter is usually lessened, if not removed, by a lotion of six or eight grains of iodide of potassium to an ounce of water.) Mr. Critchett and Mr. Bowman, at the Royal Ophthalmic, Moorfields, we find, in chronic ophthalmia or conjunctivitis, usually employ mild stimuli, solutions of acetate of lead, and sulphate of zinc, though the former may be objectionable if ulcer exists or threatens. If the symptoms are attended with much irritation, the old-fashioned opium wine is perhaps the best remedy. A solution of hydrocyanic acid, about a fluid drachm to the ounce of water, is also very serviceable. If the case have progressed, so as to give rise to troublesome mucous discharge, Mr. Critchett feeds the patient well, and uses the sulphate of copper or nitrate of silver. The excellent directions as to diet and treatment, recently given by Mr. Critchett, (*THE LANCET*, vol. i. 1855, p. 602,) leave us little to add on this subject. In *acute* conjunctivitis, if attended with lachrymation, the practice of Moorfields consists of leeching, with the subsequent use of iced water. Mr. Critchett believes that stimulation does harm. If, however, there be a puro-mucous discharge, he applies a weak solution of the nitrate of silver, with ordinary internal remedies to suit each case.

THE USE OF THE OPHTHALMOSCOPE.

Believing as we do that the student who neglects his attendance on the ophthalmic portion of his studies, neglects an opportunity of the most valuable kind, and one that may never again present itself, we would direct his attention, in a very especial manner, to the use of the ophthalmoscope. It may appear at first to be a mere plaything, as did the stethoscope and microscope; but it is destined to play a very important part in the interpretation of eye-cases before long. This very useful instrument continues to be more and more used and valued every year. Some ophthalmic surgeons, however, deny its value altogether; it requires—as we have seen it used by Mr. Wharton Jones, Mr. Bowman, and others—some careful attention to details, as regards the arrangement of the artificial light, the particular kind of disease, the relative position of the patient to the operator, &c. No little care and some perseverance are also necessary in acquiring the requisite aptitude for using it, even in the most marked case of disease, or the student will see nothing, as he too often will hear nothing with the stethoscope at first; but he afterwards comes to diagnose endocardial from pericardial sounds, and other subtleties of a like kind. The student, as observed by Mr. Paget, must train and educate his senses if he wish to keep on a level with the advance of scientific surgery at present. The most minute speck in the lens, or vitreous humour, for instance, is readily detected by the ophthalmoscope, the deep-seated and more obscure changes of the retina are also intuitively caught up in the mind before they reach the stage of incurable disease. It shows a very curious group of changes in the delicate arrangement of the vitreous humour where such exist, as pointed out to us in a case recently by Mr. Bowman; it also demonstrates the condition of the circulation in the small ramifications of the arteria centralis of the retina and its accompanying vein, and the general state of that very important nervous expansion, the little spot of Scemmering, with its yellow folds, connected, it is thought, with the lymphatic system, all so infinitely delicate and minute. We have seen of late several new forms of this instrument—that of Helmholtz, Coccuss, and Mierstein, of these the last-named form is the best.

STRABISMUS.

This operation is now so frequently performed that it may be regarded as one of the established facts of surgical science.