In one breast operation I had to scoop out a lot of enlarged glands from the axilla, and I employed two skeins of catgut for the drainage. As the healing went on, although the skin presented its natural appearance, there was a considerable serous discharge, and the separate strands of the catgut evidently underwent a process of solution. This free serous discharge is characteristic of Listerian treatment, and I attribute its occurrence in this last case to the action of the strong carbolic acid (5 per cent.) solution with which I washed the cut surfaces. Here then, it appears to me, we have a satisfactory explanation of the point of variance between Neuber and myself. In my practice the discharge is trifling, after the immediate oozing has ceased, and the tube remains incorporated in the wound. With Neuber it is melted down and washed away. The strong solution is used, the cut surfaces acted upon are cauterised and have a whitened or greyish appearance. That is to say, they are covered with a film of coagulated albumen, which is nothing more than dead matter, and, impregnated as it is with a powerful irritant, must come away in great measure in the form of discharge. This is the peculiarity of Listerian wounds, and hence the necessity of Listerian drainage. It may be true that without the drainage the germs are destroyed, but the mischief done mechanically by tension from pent-up fluid would be as disastrous as anything occasioned by living organisms.

Why this inconsistent cauterisation should be put forward as essential to perfect Listerian practice I have always been at a loss to understand, seeing that a solution of half the strength is considered sufficient to cleanse the instruments, and for the spray. Surely, if the germs on the hands, instruments, and sponges succumb to a solution of $2\frac{1}{2}$ per cent., it cannot be necessary to kill a part of the raw surface with a 5 per cent. solution to involve the few germs that may be there in the catastrophe. It is not maintained that the external germs—the spongeal, instrumental, manual, imported, vagabond germs—are in any way different from the germs in the wound, the colonised germs; so that if $2\frac{1}{2}$ per cent. is enough for the one, 5 per cent. must be too much for the other; or if 5 per cent. be the true germicidal proportion, the germs in the $2\frac{1}{2}$ per cent. spray may still dance and be merry as motes in the sunbeam. Five per cent. either is or is not necessary: if necessary, the spray is a mockery; if not necessary, the wound treatment is bad surgery. It looks as if there were no escape from the horns of this dilemma.

The conclusions, then, that I have to draw with regard to the bone drainage-tubes are—1st, that they may be used without any immediate mischief in cases where they are wanted only for a short time; that is, in cases where there is a discharge only for the first few hours after operation. But in my experience there is even then this disadvantage, that with rapid union they remain encased and unabsorbed too long, and in that condition they generally cannot be said to do any good, and they may do harm. Time will show this. 2. When, however, from the nature of the case or the germophobic strength of the dressings the discharge is expected to be copious, and to continue for several days, the bone tubes are wholly inapplicable; hence it need excite no surprise that in Neuber's ovariectomies their use was "disastrous." For the drainage after this operation the glass tube must hold its own against all comers.

Granville-place, Portman-square.

TWO CASES OF MYXCEDEMA.

BY THOMAS INGLIS, F.R.C.P. ED.

So few cases of myxœdema have been recorded in the medical journals that it is clinically important to note a short account of them as they occur, as this form of disease is somewhat uncommon, is insidious in its symptoms, and its treatment hitherto difficult and unsatisfactory. The morbid condition was first pointed out by Sir William Gull in an article on "A Cretinoid State supervening in Adult Women,"¹ and was subsequently more fully described, and its pathology partly elucidated by Dr. Ord to the Medical and Chirurgical Society of London;² while cases have been observed by Drs. Duckworth, Sanders, and Savage.³ Its

most prominent feature consists of a jelly-like formation in the subcutaneous areolar tissue, which gives the patients a stolid, puffy appearance, approaching that of the cretin. Their appearance is suggestive of the dropsy of recent renal disease and the passive anasarca in cardiac complaints, but we find the urine usually normal till the termination of the illness, and the heart is also healthy. All the patients hitherto observed have been females, except a doubtful male case under the care of Dr. Savage. I will give a general outline of two cases at present under observation.

T. W—, a plumber, of temperate habits and good general health, became depressed in mind owing to his wife's bad conduct, and was admitted to an Asylum, labouring under melancholia, with strong suicidal tendencies, when the following particulars were ascertained. There was no hereditary predisposition to insanity or other neurosis, and he had two healthy children. He had taken a large dose of laudanum with suicidal intent, and required the application of remedies and artificial respiration when admitted. He answered questions in a confused way; stated he wished to take away his life.

In three months he made a recovery from the melancholia, but was observed to be slightly enfeebled in mind.

When the first symptoms made their appearance, a general swelling, affecting principally at first the eyelids and hands, and afterwards extending over the extremities and body, gradually and insensibly came on. This swelling was first noticeable about the face, where the skin of the eyelids became puffy, wrinkled, and waxlike, widening the nose, and causing a general blurring or obliteration of the facial expression, and giving it a flattened, cretinoid aspect. The nose looked levelled down, the lips were thick and heavy, the lower one protruding. The tongue looked too large for the mouth, but was not œdematous, while the interference with the function of speech was very evident, articulation being slow, indistinct, and paretic. The mode of speech resembles that of the hemiplegic, and the thickened articulation of the epileptic, and was most affected in the morning. The hands got much enlarged and "spade-like," and he complained of their being stiff, cold, and numb. On pressure over the front of the tibiae, where the œdema was considerable, there was no pitting. The urine contained no albumen; sp. gr. 1015. The heart's action was slow and laboured, but the sounds were healthy. The surface of the body was always cold to the touch. Temperature in axilla, 95° . He never felt warm even in the hottest weather. All over the skin is dry and brawny, and its sensibility is slightly impaired, somewhat resembling scleroderma. Although the skin is dry it is often subject to short-lived perspirations. There are no reflex contractions on tickling the soles of the feet. There is no atrophy of the muscles, but the grasp of the hands is feeble, chiefly from inability to flex them. There is an absence of excitability of the muscles of the legs and arms. On faradising the muscles of his face only a slight response (Weiss, fifty cells) was obtained in them. On applying the electrode of the intermitted battery current to his scalp, where there is little œdema, there is strong reaction, and he then says he sees flashes before his eyes.

The fact that the faradaic contraction of the muscles where the œdema was marked is much impaired would point to a degeneration of the nerve-tissue at the periphery, as it is known that the normal excitability of the muscles is retained after gross lesions of the central ganglia, as in hæmorrhage or softening; while, after an injury to a nerve, such as the sciatic, there is no electrical response. On the other hand, that there is no atrophy of the muscles militates against the theory of peripheral nervous degeneration, and it is probable that the myxœdema presents a mechanical obstruction to the stimulation of the nerve endings. The helplessness and slow mobility appear to be due more to the mechanical obstruction of this mucoid degeneration than to defective innervation. He presents a blown-up, water-logged appearance, and waddles like a duck when walking. The appetite is usually very bad, and he can go without food for days and not feel hungry. When he does eat, the food tastes badly to him. Bowels are constipated. He has become rather deaf, while sight and smell remain good. He has become very lethargic, and sleeps a great deal. The mental characteristics are now slight enfeeblement; memory slightly impaired as to recent events. In the mind, as with the body, the period of activity is over, and his small routine duties are performed in an automatic, listless manner.

¹ Clin. Soc. Trans., vol. vii.

² THE LANCET, Oct. 18th, 1879.

³ Journal of Mental Science, Jan. 1880.

The second case is that of a woman, aged thirty-nine, of good general health, the mother of eight healthy children. She comes of neurotic stock, with an hereditary predisposition to insanity and other neuroses. Her mother died of apoplexy, and a sister was treated in an asylum for puerperal mania. Her illness came on five years ago, at the same time as an attack of melancholia, and was attended with perversions of the special senses. She said everything smelt of gunpowder. She presents the same bodily symptoms as T. W—, but in a lesser degree. Her mental condition is now permanently impaired. Three years ago an attack of mania followed the melancholia, and she had to be sent to an asylum. There she expressed many changing delusions, as that she was a queen, and sent to Edinburgh Castle for an escort to take her to Leith Fort. Sight is good, but the senses of smell and taste are impaired. She alleges that she smells snuff in her food, and that it tastes like poison. Numerous warty growths are scattered over her arms. Like the man she feels worst in the morning, when, she says, she can hardly see. The urine is natural, sp. gr. 1010. The heart's action is regular, the second sound being sharp and accentuated.

Her natural temperament was amiable, but she is now irritable, passionate, and sometimes violent without cause. Is becoming demented, is silly and childish in conversation, and frequently talks incoherently to herself. There is loss of that important faculty, the attention, and the judgment is warped, but the memory is still tolerable, and her affection for her children remains.

That the central nervous system is largely involved in myxœdema seems unquestionable, as in all the cases on record mental disturbance in some degree has been observed, ranging from perversion of the affection, irritability, slight depression, and loss of enjoyment in life, to aggressive mania and fatuity.

It appears to be a general progressive disease, which distinguishes it from simple scleroderma.

Post-mortem examination by Dr. Ord revealed a general œdema of the subcutaneous areolar tissue, which was composed largely of mucin and not of serum. It was also found in the internal viscera. The arteries of the brain were much degenerated and atheromatous, but the microscopic cerebral structure does not appear to have been fully examined. Where the extremities of the cutaneous nerves were observed, it was seen that they were enveloped in a soft transparent substance, so as to be padded and removed from the ready operation of incident impulses, tactile, thermal, or chemical.

When fluid accumulates in the subcutaneous areolar tissue we have a tense shining skin, pitting on pressure, and making its appearance in the lower extremities. When from impoverishment of the blood or pressure on veins we have dropsy, the œdema at first is not stationary, but is altered by position. In myxœdema we have the reverse. Scleroderma differs from it in being a sclerosis of the areolar tissues, and not an elastic mucoid change, but a limited indurated change unaltered by compression.

Although mental disturbance or aberration in some phase exists in all cases, the disease does not necessarily lead to insanity when it supervenes. This is possibly due more to the individual temperament or constitution of the patient than to the *materies morbi*, and in this particular it resembles many lesions and injuries of the brain, which frequently but not inevitably cause mental disorder. Its characters are so varying that it does not follow any typical course, but generally the lethargy and heaviness of the body find a reflex in the inactive and torpid mind. While the delusions and perversions of the special senses in the case of Mrs. H— are undoubtedly attributable to disordered brain function, the occasional profuse perspirations are also the indication of a failure of nerve power, and, like plithisical sweating, are due to the defective innervation of the nerves controlling the sweat-glands.

The irritability and mental excitement noticed in the first stages are probably the resultant of the action of poison on the cerebral convolutions, as when poisons act on the intelligence, such as alcohol or opium. There is first a high stimulation and sur-excitation, and ultimately derangement of the intellectual functions.

In some cases the departure from the normal mental state at the onset of the disease is marked more by a change in tone than by an actual state of insanity. The man feels ill, loses his sense of well-being, spontaneity; he feels uncomfortable, irritable, and listless—a phase of mild melan-

cholia. The changes come on slowly and insidiously; the alteration is noted first in temperament and disposition.

Dr. Ord is uncertain whether this mucoid change is a new product, the result of a disordered nutrition, a degeneration altering the function of the supreme cerebral centres, or a retrograde metamorphosis to a lower type, such as cretinism. Certainly the mental and bodily symptoms show a marked filiation to this disease.

I look upon myxœdema as primarily a nervous disease, and that the grosser changes in the skin and viscera are caused by deficient absorptive action of the lymphatics, the result of trophic changes in the cells of the higher ganglia.

The treatment of the disease has hitherto been very unsatisfactory, and the administration of nervine tonics, galvanism, and baths appear to have no beneficial effect, either in favouring the absorption of the deposit, or checking the progress of the disease, and at the best exercise only a palliative influence.

Deeping St. James.

ANTISEPTIC MIDWIFERY.

By WALTER VERDON, F.R.C.S.

THE following case is, I think, of interest in showing the value of Mr. Lister's treatment applied to midwifery.

Lately I have used antiseptics in ordinary midwifery cases, and with marked benefit. The discharge ceases sooner and does not become offensive. In some cases this disappears entirely during the first week. It is necessary, however, in order to obtain a good result, to guard against the admission of septic air both during and after confinement—to view the uterine and vaginal lesions in the same light you would the wound made in a subcutaneous operation, and as carefully protect them from septic influence.

In the case I quote below, the patient was exposed to the poison of erysipelas both during and after labour; and though she had a slight attack of cutaneous inflammation accompanied with fever, the uterine wound remained intact; the poison having no more influence upon it than it would have upon a wound caused by subcutaneous rupture of a muscle.

Mrs. T—, weak and anæmic, at full time, had suffered from frequent pains for two days. She had not sent for assistance because she did not think the labour far enough advanced, but now felt so exhausted she wished for help. The os was now fully dilated, membranes ruptured, head above the brim, pelvis roomy, pains frequent and short; the head descended with and receded after each, making no progress; pulse 98, weak and compressible; the patient was much exhausted. I then determined to use the forceps, and also to take every antiseptic precaution.

I syringed out the vagina with warm carbolic water; also had a small basin of carbolic oil at hand for the instruments, hands, &c., and to anoint the labia and adjacent parts. All vaginal manipulation was performed beneath the spray, and with very little traction the child was born. The after-applications were diapers soaked in carbolic oil and carbolic tow, and an injection of carbolic lotion three times daily.

On the second day after confinement I found the nurse was suffering from an acute attack of erysipelas of the face; she told me she was feeling ill and her head felt tender the day before the confinement. She had slept with the patient on the night of the labour and on the following night.

On the third day the patient complained of sore-throat. Pulse 120; temperature 102° F. Vaginal discharge free from putrefactive smell; urine clear and copious.

On the fourth and fifth days the patient felt very ill and complained of pain in the right leg. Her tongue was coated; pulse 120. Still the vaginal discharge was going on satisfactorily, but there was a patch of cutaneous inflammation on the right thigh from the buttock extending almost to the knee.

On the sixth day the discharge ceased and did not return. The pulse remained above 100 until the eleventh day; the redness disappeared on the eighth day.

The child also was the subject of an unhealthy ulceration round the umbilicus, preceded by a superficial inflammation extending to the groin.

Both the child and mother made a good recovery.

Brixton-road.