MAP OF LOCALITIES DESCRIBED

WEATHERING OF GRANITIC ROCKS OF GEORGIA *

BY THOMAS L. WATSON

(Read before the Society December 29, 1900)

CONTENTS

	Page
Introduction.....	93
Granite.....	94
Character and composition.....	94
Discussion of the chemical analyses.....	96
Porphyritic granite.....	97
Character and composition.....	97
Discussion of the chemical analyses.....	97
Granite-gneiss.....	98
Résumé.....	99
Cause of red color of the soil.....	104
Results based on assumptively insoluble Al_2O_3 and $Al_2O_3 + Fe_2O_3$	106

INTRODUCTION †

For the past three years the writer has been engaged in a careful detailed study of the granitic rocks of Georgia, giving special attention to the phenomena of their weathering. The natural outcrops have been carefully studied in the field, and specimens of the fresh and weathered rock, representing various stages in the decay, have been collected from the most typical and widely separated localities in the state and analyzed in the chemical laboratory of the State survey by the writer. The material is sufficiently representative and the work extensive and detailed enough to arrive at definite conclusions regarding the changes involved in the transition from fresh to decayed rock in the granitic group of rocks.

Based primarily on texture and structure, three types of the granitic rocks are distinguished: (1) The massive even-granular granites; (2) the

* Published by permission of the State Geologist of Georgia.

† For a detailed discussion of the weathering of the Georgia granites, including complete chemical analyses of the fresh and accompanying decayed rock from numerous localities in the state, the reader is referred to a report on the granites and gneisses of Georgia, to be issued shortly by the State Geological Survey, by the present author.

porphyritic granites, and (3) the banded or foliated granites—gneisses. Laboratory study shows the three types to be nearly identical in mineral and chemical composition. In the field, the porphyritic facies of the granite masses are found to grade peripherally into an even-granular, medium, coarse textured granite of the same mineral and chemical composition. Likewise the gneisses are elsewhere shown to be the metamorphosed equivalents of the irruptive massive granites from which they differ only in a pronounced banded or foliated structure, secondarily induced by metamorphism. For this reason the gneisses are referred to as granite-gneisses. That comparisons may be effected and emphasis given, whatever differences arising from the influence of texture on weathering, as illustrated by the degeneration of the Georgia granitic rocks, the rock-decay is separately treated under (1) granite; (2) porphyritic granite; and (3) granite-gneiss. The several places described in the following pages are shown on the accompanying map, plate 6.

GRANITE

CHARACTER AND COMPOSITION

The fresh unaltered granites are massive, fine to medium grained rocks in texture, light gray to dark-blue gray in color, and show an admixture of fine, granular quartz, feldspars, and biotite to the unaided eye. The principal minerals are quartz, orthoclase with micropertthitic structures, microcline, plagioclase near oligoclase, brown biotite, some muscovite, and a little apatite, zircon, and magnetite.

The quartz is frequently intergrown with the feldspar in the form of rounded ovals or disks of micropegmatitic structures, clearly indicating the contemporaneous crystallization of the two minerals. It is sometimes inclosed as drop-like inclusions in many of the larger feldspar crystals. The larger individuals further show numerous lines of fracture and undulatory extinction.

The potash feldspars usually show good cleavages; are intergrown with a second feldspar, albite, as micropertthitic intergrowths, and are commonly twinned according to the Carlsbad law. The presence of considerable soda in the analyses corroborates the inference that the intergrown feldspar with the potash varieties is albite. Microcline varies in quantity, but may equal or even exceed in a few sections the orthoclase. Plagioclase is inferior in amount to the potash feldspars, and, as a rule, affords low extinction angles in basal sections, indicating an acid feldspar near oligoclase. This inference is corroborated by the percentage of lime shown in the analyses. The biotite is deep brown in color and strongly pleochroic. It is present in irregular shreds and elongated

FIGURE 1.—GRANITE QUARRY NEAR GREENVILLE, GEORGIA

Showing gradation from residual clay to fresh granite and illustrating method of granite weathering

FIGURE 2.—GULLIED SLOPE IN DEEP RED PLASTIC CLAY DERIVED FROM FRESH GRANITE

GRANITE QUARRY AND ERODED SURFACE IN RESIDUAL CLAY

plates, with good crystallographic boundaries, and intimately associated with muscovite when present. A part of the biotite is invariably partially altered to chlorite. Muscovite varies in quantity, and is always inferior in amount to the biotite. Epidote is somewhat abundant as a secondary constituent in some of the granites. The remaining microscopic accessories present show the usual characteristic features.

The first decided change in the weathering of the massive granites, in which an almost complete change in appearance from the perfectly fresh granite is indicated, is represented by a hard and firm, dull grayish granitic mass, tinged from a faint to a highly ferruginous rusty brown color. It varies in tenacity from the firm, hard rock to specimens which readily crumble under the gentlest pressure. Aside from change of color and lack of luster in the component minerals, the rock has lost its compact, close grained texture, and the partially changed granite presents instead a somewhat spongy or loose texture in appearance, in which the individual mineral grains are slightly forced apart from each other. The feldspars are of a decided white color, perfectly opaque to the naked eye, and have commenced to split along the cleavage planes. Under the lens, the feldspars show comparatively fresh luster in places; the biotite appears somewhat leached from hydration and oxidation, and the adjacent areas are slightly discolored from the hydrous iron oxide derived therefrom. In the center of the biotite plates the luster on the cleavage faces is as brilliant and the color as strong as that in the fresh rock. Thin-sections made from hand specimens representing this stage in the weathered rock show a marked yellow color to the naked eye, indicative of the iron oxide staining derived from the decomposition of the biotite.

A microscopic examination of the thin-sections confirms the above macroscopic description. Under the microscope the feldspars appear somewhat cloudy and opaque, but otherwise are comparatively fresh looking. The plagioclase feldspars are more altered, as a rule, than the potash varieties. The alteration takes place mostly along the twinning planes in the plagioclase, and mainly along the lines of cleavage in the potash feldspars. The only visible change in the quartz is a splitting up along the innumerable fracture lines subsequently induced through dynamic action. The quartz and feldspar areas, particularly along the lines of cleavage and breakage, are stained yellow by the hydrous iron oxide derived from the partial decomposition of the biotite.

The biotite appears in part perfectly fresh, but a large proportion of this mineral has lost its cleavage and optical properties, and is replaced by an amorphous yellow to brown colored mass, with the surrounding areas partially discolored from the ferruginous constituent. In still other shreds of the biotite the edges and borders are frayed out and leached,

while the centers remain entirely fresh. As observed in the potash feldspars, alteration takes place in the biotite to a marked degree along the cleavage planes.

No calcite has been observed in any of the thin-sections of the partially decomposed rock under the microscope; and the further absence of carbonates in the residual granite-decay was proved by testing portions of the finely powdered rock in dilute acid.

Digestions of separate portions of the residual products, in various stages of decomposition, in very dilute HCl at the temperature of boiling water, gave solutions varying from slightly colored to a deep wine-red in color, according to the extent of decay and the amount of iron oxide liberated from the iron-bearing silicate minerals and retained in the residual product. The residues from these separately treated portions of decayed rock indicated, according to the degree of decay, varying amounts of the fresh and partially fresh minerals, quartz, feldspar, and biotite. Numerous magnetite grains, just as fresh apparently as in the unaltered granite, were found to some extent in all the residues. In the advanced stages of decay, represented by highly ferruginous red-colored plastic clays distinctly gritty in feeling, owing to the presence of free quartz and partially decayed silicate minerals, the biotite shreds are changed from black, highly lustrous plates in the fresh granite to brittle brass-colored foliæ in the ferruginous clay, and are slowly soluble in both cold and hot dilute acid.

DISCUSSION OF THE CHEMICAL ANALYSES

Five widely separated sections of the massive even-granular granites were selected for making chemical analyses of the fresh and corresponding decayed rock for purposes of illustrating the changes accompanying the transition in weathering. The residual product represented stages of decay, varying from distinctly discolored, hard and firm rock to highly ferruginous red plastic clays. A careful study of the tables of analyses, representing the rocks in various stages of decay, shows the principal features in the change from fresh to decayed granite. On the assumption of constant Fe_2O_3 , the following calculated maximum and minimum percentage amounts of each constituent have been retained or saved, according to the corresponding stage in decay:

	Maximum	Minimum
SiO_2	92.20	22.80
Al_2O_3	99.37	56.18
CaO	57.28	1.18
MgO	54.21	12.06
Na_2O	85.86	7.85
K_2O	96.60	8.26

FIGURE 1.—PARTIALLY WORKED UP BOULDER OF PORPHYRITIC GRANITE
Near Palmetto, Georgia. Vertical section, 10 feet

FIGURE 2.—GULLIED SURFACE IN RESIDUAL CLAY DERIVED FROM PORPHYRITIC GRANITE
Along the Atlanta and West Point railroad, $4\frac{1}{4}$ miles south of Palmetto

PORPHYRITIC GRANITE NEAR PALMETTO, GEORGIA

FIGURE 1.—TWENTY-FOOT SECTION OF RESIDUAL CLAY DERIVED FROM PORPHYRITIC GRANITE
Along public road, near depot, Chatham, Virginia

FIGURE 2.—GULLED SURFACE IN RESIDUAL CLAY OF PORPHYRITIC GRANITE
Near Depot, Chatham, Virginia

RESIDUAL CLAYS DERIVED FROM A FOLIATED BIOTITE PORPHYRITIC GRANITE

The transition from fresh to decayed granite is accompanied in every case by a gain in H_2O , which increases in amount proportional to the stage in weathering. The percentage loss for the entire rock is also proportional to the degree of decay reached in the residual product. The amount lost for the entire rock varies from 8.14 to 71.82 per cents respectively. The figures further indicate that the early stages in weathering are mostly in the nature of disintegration, while the chemical forces—decomposition—become the principal agents in the transformation in the later stages. The residual soils are accordingly gray in color and principally sandy, or deep red in color and principally clayey.

PORPHYRITIC GRANITE

CHARACTER AND COMPOSITION

Three of the largest and most typical porphyritic granite areas were selected for indicating the processes involved in the weathering of this type of rock in Georgia. Two of the areas represent the massive porphyritic granite, while the third is a pronounced foliated type of the same rock.

This type of granite only differs from the equivalent even-granular facies in porphyritic structure, and, as a rule, somewhat coarser grained in texture. The porphyritic texture grades, in most cases, into the even-granular granite facies peripherally, with no difference shown in mineral and chemical composition. They consist of a coarse grained granite matrix of quartz, feldspar, and biotite, in which are imbedded potash feldspar phenocrysts 10 to 50 millimeters long and 5 to 10 millimeters across; idiomorphic to allotriomorphic in crystal outline, with pronounced cleavages parallel to (001) and (010), and are twinned according to the simple type of Carlsbad twins.

The weathering of the natural rock exposures in the field is in every way analogous to that of the massive granites, with usually no visible difference apparent between the two, on textural grounds. The chemical changes accompanying the transition from fresh to decayed rock are closely similar in the two types of granite.

DISCUSSION OF THE CHEMICAL ANALYSES

The principal features to which attention need here be called are (1) a gain in H_2O , which increases as the change becomes more advanced; (2) a total percentage loss for the entire rock, varying, according to the

completeness of the change, from 15.84 to 35.07 per cents respectively ; (3) the transition from fresh to decayed granite has been accompanied by the retention of the following maximum and minimum percentage amounts of each constituent saved :

	Maximum	Minimum
SiO ₂	81.72	60.64
Al ₂ O ₃	92.51	89.11
CaO	86.64	3.71
MgO	69.04	58.23
Na ₂ O	91.29	26.26
K ₂ O	93.16	52.18

GRANITE-GNEISS

The Lithonia area of contorted gneiss in DeKalb county best illustrates the character of and changes incidental to the weathering common to this type of rock in Georgia. It has been shown elsewhere by the writer that the gneisses of Piedmont, Georgia, are closely similar to the granites in chemical and mineralogical composition, and are accordingly metamorphosed granites. The gneisses are only distinguished from the massive granites by a pronounced banded or schistose structure subsequently induced by metamorphism. They are therefore designated granite-gneisses in contradistinction to gneisses of known sedimentary origin.

The sap (partially decayed surface rock) of the doming masses of Lithonia granite-gneiss and the adjacent famous Stone Mountain granite boss is very thin, rarely averaging, as a rule, more than two inches in thickness. It is hard and firm surface rock, discolored a slight red from the partial hydration and oxidation of the biotite. The feldspars are dull and opaque from kaolinization, and in most cases the rock indicates alteration to a greater depth than the discolored portion, manifested by general dullness in appearance of the stone through more or less feldspathic alteration. Numerous exposed sections of apparently completely disintegrated and decomposed granite-gneiss are exposed, varying in depth from 10 to 15 feet, with the schistosity of the original fresh rock almost perfectly preserved in the residual clays forming the sections.

Microscopically the gneisses are composed of quartz, orthoclase and microcline, intergrown with a second feldspar, albite in the form of micropertthitic intergrowths, an acid plagioclase near oligoclase, and biotite. A little magnetite, garnet, some apatite, and zircon are present.

FIGURE 1.—COMPLEXLY CONTORTED GNEISSIC BANDS OR LAYERS

FIGURE 2.—ANOTHER TYPE OF VARIATION IN STRUCTURE OF THE GNEISS

GRANITE-GNEISS, LITHONIA, GEORGIA

FIGURE 1.—THIRTY-FOOT SECTION OF VARI-COLORED RESIDUAL CLAY
Derived from decay of gneiss

FIGURE 2.—FORTY-FOOT SECTION OF RESIDUAL CLAY
Derived from decay of gneiss. Original schistosity planes of fresh rock perfectly preserved

RESIDUAL CLAYS DERIVED FROM DECAY OF GNEISS, NEAR ATLANTA, GEORGIA

More or less chlorite, muscovite, and epidote occur as secondary minerals derived from the alteration of the feldspars and biotite. The potash feldspars predominate and the species vary in amount from place to place. Microcline is somewhat more abundant than in the even-granular granites and their equivalent porphyritic facies. The plagioclase also varies in amount. The chemical analyses corroborate the inference that the plagioclase is oligoclase in the percentage of lime present. They further indicate a larger percentage of free quartz than the massive granites and their porphyritic facies, which is confirmed by the microscope. Slight peripheral shattering of the larger quartz and feldspar crystals, numerous lines of fracture, and wavy extinction common to these two minerals afford evidence of the effects of dynamo-metamorphism.

The physical and chemical changes involved in the processes of decay of the gneisses are strikingly similar to those of the granite types.

RÉSUMÉ

The following conclusions are based on the percentage amounts saved and lost per each constituent, and the total loss for the entire rock, calculated from the analyses of the fresh and corresponding decayed rock, on an assumed Fe_2O_3 constant factor.

A careful study of the tables of chemical analyses of the fresh rocks and their corresponding decayed products shows the amount of water to rapidly increase as the decomposition advances, which, as Merrill* has shown, becomes the most important factor in the earlier stages of rock weathering. The change has in every case been accompanied by a loss in the silica, a proportional greater loss in the alkalies, lime, and magnesia, with a proportional increase in the iron oxide, and in some cases in the alumina.

The decayed product of the Georgia granites invariably show an abundance of quartz grains derived from the free quartz in the fresh granite. In most cases the quartz granules have been only slightly corroded and are prevailingly angular in outline. Frequent examination of the quartz grains in the residual clays was made under the microscope for evidence of etching and solution, but positive evidence was lacking in every case. The loss, therefore, in this constituent in the Georgia granites has been derived in large part, if not entirely, from the silicate-bearing minerals, feldspar and biotite, which was doubtless removed with the alkaline solu-

* Rocks, Rock Weathering, and Soils, New York, 1897, p. 234.

tion of the more soluble constituents, as liberated in the free-nascent-state. It has been shown,* however, that rocks composed of silicate minerals, almost or entirely lacking in alkalies, lost a portion of their silica with equal readiness. This has been explained on the supposition that as the silica was liberated in the nascent state it was soluble either in pure or carbonated water.

The iron is mostly present in these rocks in the form of ferrous iron. The stability of this constituent characterizing all the analyses of the fresh and decayed rocks may be readily accounted for on the supposition that the decomposition was promoted in a sufficient supply of oxygen, whereby all of the iron was converted into the form of insoluble hydrated sesquioxide and retained with the residue. The analyses indicate only a slight loss in most cases in the alumina, while in others it has all been retained in the transition from the fresh to the decayed rock.

The lime, magnesia, and alkalies obey the usual laws in the proportional amounts lost and saved for such rocks. The lime has invariably disappeared in larger percentage amounts than the magnesia, and likewise the sodium salts have been removed, as a rule, in larger quantities than those of potassium. The feldspars are, of course, the principal source of the soda and potash in the rocks, and the removal of soda in larger amounts than the potash indicates that the potash feldspar varieties are more refractory toward normal atmospheric agencies than the soda-lime or plagioclase varieties.†

The accessories—tourmaline, magnetite, zircon, and garnet—all occur to some extent in the residual granite sands scarcely affected at all by the atmospheric agents.

In the most advanced stages of decay the residual product is a highly plastic ferruginous clay, rendered gritty from the presence of free quartz granules mainly, along with a small percentage amount of the undecomposed silicate minerals. In such cases the transition from fresh to decomposed rock has been accompanied by an unusually large proportional loss in the constituents, greatly exceeding in several instances 50 per cent.‡ In these cases the change is not far from complete, as shown, not only in the chemical analyses, but in the relatively small amounts of undecomposed minerals present in the clay as well. Where the change

* Ebelmen: *Ann. des Mines*, 1845, vol. vii.

Merrill, Geo. P.: *Op. cit.*, pp. 226, 227.

† Roth's *Allgemeine u. Chemische Geologie*, 3d edition, 2d heft. Geldmacher M., *Beiträge zur Verwitterung der Porphyre*, Leipzig, 1889. *Lemberg, Zeit. der Deut. Geol. Gesellschaft*, 1876, p. 28.

‡ Merrill, Geo. P.: *Rocks, Rock Weathering, and Soils*, New York, 1897, p. 234.

is accompanied by so large a total loss in the constituents for the entire rock, the process is an advanced one and entirely in the nature of chemical decomposition. In the earlier stages of weathering the rock begins to crumble and is ultimately reduced to a light gray-colored granitic sand, through hydration and temperature changes—disintegration—accompanied by only slight chemical change. The following summary of results, showing the total percentage loss of constituents for the entire rock in the various stages of decay, as calculated from the analyses, will indicate to some extent whether the process involved has been mainly one of decomposition or disintegration :

Locality.	Total loss for entire rock.
Average of analyses from the Crossley and Southern Granite Com- pany's quarries.....	7.68
Swift and Wilcox quarry, near Elberton.....	7.92
Coggin's Granite Company's quarry, near Ogleby.....	8.14
The Lexington Blue Granite quarry, near Lexington.....	14.56
Heggie Rock, Columbia county.....	15.84
Crossley quarry, near Lithonia.....	26.49
Brinkley place, near Camak.....	34.04
McCollum quarry, Coweta county.....	35.07
Coggin's Granite Company's quarry, near Ogleby.....	36.38
Cole quarry, near Newnan.....	38.48
Greenville Granite Company's quarry, near Greenville.....	61.85
Greenville Granite Company's quarry, near Greenville.....	71.82

In the Georgia granitic rocks, it is observed that in some localities the biotite begins to give way first, while in others the feldspar appears to be the most affected ; and in others still the rate of decomposition is seemingly about equal for the two minerals. The problem is somewhat complicated, however, in the case of the Georgia rocks, owing to the large proportion of plagioclase, soda-lime, feldspar present, and the abundance of micropertthitic structures common to the potash feldspar varieties.

During the early stages of weathering in these rocks chemical decomposition is subordinated to physical disintegration, and the rocks have suffered mainly through temperature changes, such as produce granulation and weakening in the adhesive power between the individual minerals—feldspar and biotite—resulting in crumbling along the planes of cleavage. Freezing and solution have aided in the process, but owing to the low absorptive power of the Georgia rocks, they can not be con-

sidered as essential factors. The following tabular statement shows the percentage or ratio of absorption for the Georgia granites, determined by the writer in the laboratory of the State survey:

Locality.	Gain in absorbed water.	Percentage of absorption.
The Charley Rocker quarry, Hancock county.....	0.04	0.037
The Sparta quarry, Hancock county	0.04	0.049
The Lexington Blue Granite quarry, Oglethorpe county.	0.08	0.092
The Diamond Blue Granite Company's quarry, Oglethorpe county.....	0.11	0.088
The Swift and Wilcox quarry, Elbert county.....	0.08	0.090
The Wright place, Elbert county	0.09	0.092
Tate and Oliver quarry, Elbert county.....	0.08	0.093
Coggin's Granite Company's quarry, Elbert county...	0.15	0.090
The Childs's quarry, Elbert county.....	0.17	0.092
The Snell quarry, Gwinnett county.....	0.10	0.075
The Linch quarry, Putnam county.....	0.04	0.060
Greenville Granite Company's quarry, Greenville....	0.11	0.086
The Odessa quarry, Meriwether county	0.05	0.056
Stone Mountain Granite quarries	0.09	0.067
Arabia Mountain Granite quarries.....	0.04	0.050
Pine Mountain Granite quarries.....	0.06	0.073

The fact that granitic rocks may undergo extensive disintegration with but slight decomposition has been emphasized by several writers.* Numerous cases where the rocks have been resolved into residual products from disintegration with little decomposition, and also where decomposition follows so closely on disintegration, the former greatly predominating, are well illustrated among the various types of the Georgia granites. The data are yet insufficient, however, to draw any definite conclusions concerning the difference in weathering, if there be any, between the fine and coarse grained granites of the state; that is, to state definitely whether the coarse grained varieties suffer mainly from disintegration and the fine grained ones by disintegration closely followed by decomposition.

The early stages in the processes of weathering, as elsewhere pointed out in this paper, for each type, fine and coarse grained, of the Georgia

* Voyage of the Vega, 1881, vol. ii, p. 420. Quoted by Merrill, Geo. P., op. cit., p. 242. Bull. Geol. Soc. Am., 1895, vol. 6, pp. 321-332; also, Rocks, Rock Weathering, and Soils, 1897, pp. 206-213; Rocks, Rock Weathering, and Soils, 1897, p. 243.

rocks is mainly one of disintegration accompanied by some chemical change, while the later stages are as prevailingly characterized by decomposition, resulting in the production of a stiff, plastic red clay. The earlier stages in the weathering of the fine grained granite are apparently the same as for the similar stages in the coarse grained porphyritic granites. The same statement holds for the two types in the advanced stages of weathering—decomposition. The striking similarity in the granite, porphyritic granite, and gneiss, as relating to solubility in acids, is well illustrated in the percentage amounts of soluble matter extracted from weighed portions of the finely powdered fresh rock, by digestion in 100 cc. of half normal HCl for three hours, at the temperature of boiling water.

<i>Granites</i>	Per cent of soluble matter.
The Oglesby blue granite, Elbert county	10.56
The Hutchins blue granite, Oglethorpe county.....	9.77
Swift and Wilcox light gray granite, Elbert county.	9.20
Echol's mill light gray granite, Oglethorpe county.....	8.37
Average	9.475

<i>Porphyritic Granite</i>	
Heggie Rock, Columbia county	10.98
Heggie Rock, Columbia county (decay).....	12.69
Brinkley place, near Camak	16.29
Brinkley place, near Camak (decay).....	23.69
Porphyritic granite near Line creek, Fayette county.....	9.47
Average of fresh rock.....	12.246

<i>Granite-gneiss</i>	
Southern Granite Company's quarry, near Lithonia.....	7.58
Arabia Mountain quarries, De Kalb county.....	6.06
Tilley quarry, Rookdale county	5.94
Average	6.526

The lower percentage of soluble matter in the granite-gneisses over the granites and their porphyritic facies is readily accounted for on the basis of this rock type containing a larger amount of free quartz, as shown in the chemical analyses. The inference is also corroborated by microscopic study. In each case where the total percentage of extract from

the decayed rock was determined an increase over the corresponding extract from the fresh rock was indicated.*

CAUSE OF RED COLOR OF THE SOIL

The cause of color variation in soils has been discussed in recent years by Crosby,† Dana,‡ Russell,§ and Merrill.|| Thus far the writer's observations are in accord with those of Crosby and Merrill in supposing this to be a superficial phenomenon. The most brilliant hues are observed to be confined to the immediate surface portions, varying from an inch to many feet in depth, which gradually give way in turn to more modest hues of yellow and brown, and finally pass into the grays and other colors of the fresh parent rock beneath, as the depth increases. Such a transition may be observed in almost any recent excavation in the crystalline belt of the southern Appalachians. There are, however, two exceptions to this which without proper differentiation will prove very misleading:

(1) In those cases where the sections have been exposed for some time the transition in vertical section becomes quickly obscured by the washing down of the brilliant ochreous coloring matter from above, frequently coating the section face to the entire depth of exposure with a varying thickness of the ferruginous wash of apparently the same degree of color. This invariably happens, and may be observed to advantage after heavy rains in almost any part of the above belt.

(2) In still other cases exceptions are to be found in the weathering along the joint planes and other approximately vertical as well as horizontal planes of weakness found cutting the rock masses. In these the color is most intense along the immediate surface of the planes, and extends oftentimes, in case of the vertical planes, to a considerable depth.

Crosby is inclined to regard the increase in color of the superficial portions to be due wholly to dehydration of the ferric salts, while Merrill recognizes not merely the *quality* of the coloring matter, but emphasizes the importance of *quantity* as well. Merrill¶ says: "This is well

* Geo. P. Merrill : Op. cit., 1897, p. 219 ; Bull. Geol. Soc. Am., 1896, vol. 7, pp. 349-362.

† W. O. Crosby : Proceedings Boston Soc. Nat. Hist., 1884-1888 (1888), vol. xxiii, pp. 219-222 ; Technological Quarterly, 1891, vol. iv, p. 36.

‡ J. D. Dana : Amer. Jour. Sci., 1890, xxxix, pp. 317-319.

§ I. C. Russell : U. S. Geol. Survey, Bulletin no. 52, 1889, 65 pp.

|| Geo. P. Merrill : Bull. Geol. Soc. Am., 1897, vol. 8, pp. 161-162 ; Rocks, Rock Weathering and Soils, 1897, pp. 384-386.

¶ Bull. Geol. Soc. Am., 1897, vol. 8, p. 162.

shown in the fractional separations made by washing these residual clays and sands, whereby they are separated into proportional parts of varying degrees of fineness." It has been recently shown in climates where rock weathering is accompanied by a leaching process, in which the more soluble constituents are more or less gradually removed in solution, the ferric and alumina salts are found to be the most refractory of the essential constituents in igneous rocks. The resulting final product of decomposition consists therefore of a highly ferruginous clay, composed mainly of the hydrated aluminous silicates, free iron sesquioxides, and, in the case of acid rocks, additional free silica.

Merrill* remarks that, all things being equal, the depth or degree of color indicates advanced decomposition and also geological age. This statement accords well with the writer's work on the Georgia granites. The gray soils derived from the granitoid rocks of the state represent not an advanced stage of decomposition, but mainly a process of disintegration manifested in the partial kaolinization of the feldspars and a slight rusting, oxidation of the biotite; while the red soils derived from the same rocks are invariably deep, red ferruginous clays in which the biotite is thoroughly leached and oxidized and the feldspars nearly, if not entirely, kaolinized. Stated somewhat differently, the gray soils contain an abundance of all the component minerals present in the fresh rock, but slightly altered chemically; are highly porous, sandy soils, from which clay is essentially absent, and have been derived from the fresh rock chiefly by disintegration. The red soils, on the other hand, contain comparatively only a very small percentage of the undecomposed silicate minerals; are deep red, ferruginous, stiff clays, rendered more or less gritty from the presence of free quartz, and have resulted from decomposition. The gray soils represent, therefore, the earlier stages, disintegration, in the weathering of the rocks, while the red soils are conversely the products of decomposition and represent the advanced stages in the change.

While quantity of pigment—ferric oxide—seemingly controls or rather accounts for the brilliant coloring of the soils in general, conditions afford strong indications, at least, of the suggestion of presence of some other factor or factors of marked importance. In support of this statement the author has critically examined all available authentic analyses of the decayed rock products and carefully noted each writer's description as to color. One or two illustrations will suffice:

Dr Merrill† has described the weathering of a micaceous gneiss from Albemarle county, Virginia, the decayed product—residual soil—of

* Rocks, Rock Weathering, and Soils, 1897, p. 386.
Bull. Geol. Soc. Am., 1897, vol. 8, pp. 157-168.

which he characterizes as "highly plastic" and "deep, red-brown" in color, yielding on chemical analysis 12.18 per cent of Fe_2O_3 .

The writer (Watson) has described elsewhere the changes incident to the weathering of an even grained granite near Greenville, Meriwether county, Georgia, the accompanying residual product from which is a highly ferruginous deep red, dark colored, stiff, plastic clay, slightly gritty, chiefly from free quartz, and yielding on chemical analysis only 6.33 per cent of Fe_2O_3 .

Still another case in question is the weathering of a similar fine and even grained granite near Newnan, Georgia. The residual product from this granite is a clayey mass, less plastic and not so dark in color as the Greenville material, but is a bright cherry-red in color, and yields only 1.91 per cent of Fe_2O_3 on analysis. Practically the same results are shown on comparing the recorded results of basic igneous rocks. According to these data, we apparently have residual products of essentially the same character and depth of color, but varying evidently in the amount of coloring matter—ferric oxide—present. The two extremes are represented by 12.18 and 1.91 per cents respectively, the former containing more than six times the amount of ferric oxide shown in the latter, and yet the same results as regards color are seemingly produced.

RESULTS BASED ON ASSUMPTIVELY INSOLUBLE Al_2O_3 AND $\text{Al}_2\text{O}_3 + \text{Fe}_2\text{O}_3$

It has been shown* that of the essential constituents in silicious crystalline rocks the iron and aluminum oxides are the most refractory, and accordingly the least likely to suffer loss during the processes of rock degeneration. The work of Dunnington,† Clarke,‡ and Smyth§ strongly indicates concentration, instead of loss, in titanium, in the residual product derived from the titanium-bearing minerals, upon weathering. This element occurs so sparingly distributed through the acid crystalline rocks, averaging, as a rule, less than 0.5 of 1 per cent, that it can not be regarded as a safe basis of assumption for calculating the percentage amounts of the more essential constituents saved and lost present in the rock. The same statement, as regards concentration instead of loss, also applies in some cases to several other ones of the accessory minerals sometimes present.

* Merrill, Geo. P., op. cit.

† Amer. Jour. Sci. (3d ser.), xli, pp. 491-495.

‡ U. S. Geol. Survey, Bulletin no. 78, pp. 34-42.

§ Bull. Geol. Soc. Am., 1898, vol. 9, pp. 257-269.

In the Georgia granitic rocks the iron and aluminum oxides have proved the most resistant of the constituents present toward the weathering agents, and while the ratio of increase for the two is nearly the same, that of the iron is, with several exceptions, slightly greater than alumina, and has therefore been assumed to remain constant for convenience* of comparison of the analyses of the fresh and decayed rock in tables I, II, V, VI, VIII, and IX. The figures in the tables of analyses disclose the fact that the percentage of iron oxide in the fresh and decayed rock is strikingly smaller than that of the alumina. While the ratio of increase in the iron oxide, as pointed out above, is slightly greater than that of the alumina for the decayed and fresh rock, the writer has recalculated in a similar manner all of the results: first, on the assumption of an Al_2O_3 constant basis; and, second, on the assumption of an $\text{Al}_2\text{O}_3 + \text{Fe}_2\text{O}_3$ constant basis in order to make comparisons and point out the possible differences resulting. The percentage amount of alumina is greater in every case of the Georgia granites for the decayed product than for the corresponding fresh rock, and is present in larger amounts than any other single constituent where increase in the decayed over the fresh rock is indicated. Briefly stated, a comparative study of the tables based on the assumption of, first, an Al_2O_3 , and next, an $\text{Al}_2\text{O}_3 + \text{Fe}_2\text{O}_3$ constant, shows:

1. That on the assumption of constant Al_2O_3 , the Fe_2O_3 like the H_2O indicates a slight gain, owing to the ratio of increase being greater for the iron oxide than for the alumina.

2. The calculated percentage amounts of the constituents saved and lost are in similar ratios for the three cases, but differ in actual amounts in each one. The constituent showing the greatest loss under one assumed constant likewise indicates similar loss for the other two assumptions. The percentage amount lost or saved for any one constituent is not the same in the three cases.

3. The total percentage loss for the entire rock is invariably greater on the assumption of an Fe_2O_3 constant than for the other two assumptionally insoluble constituents. In each case the lime salts have been removed in larger amounts than the magnesia, and similarly the soda has suffered a greater loss than the potash. The fact particularly emphasized is the greater loss for the entire rock on the Fe_2O_3 assumed constant over that of the Al_2O_3 and $\text{Al}_2\text{O}_3 + \text{Fe}_2\text{O}_3$ assumptions.

The tables show that the conclusions stated under 2 and 3 are governed directly by the numerical value of the ratio of increase—factor—of the constituent assumed to remain constant in the change from fresh to residual rock.

The losses for the entire rock, based on the three assumptively insoluble constituents, are tabulated below for convenience of comparison :

Table.	Fe ₂ O ₃ constant.	Al ₂ O ₃ constant.	Al ₂ O ₃ +Fe ₂ O ₃ constant.
I.....	{ 8.14 44.52 }	{ 7.40 33.63 }	{ 6.73 34.29 }
II.....	14.56	7.78	8.43
III.....		7.92	7.92
IV.....		38.48	33.72
V.....	{ 61.85 71.82 }	{ 43.60 51.64 }	{ 44.41 52.90 }
VI.....	15.84	5.59	6.65
VII.....		34.04	34.04
VIII.....	35.07	29.88	30.33
IX.....	26.49	21.10	21.61
X.....		7.68	7.08