There is a patient at St. Bartholomew's now under my care, aged twenty-five, who was born with an irregularly-shaped, dark-coloured papillary mole, covering about two-thirds of one cheek. A year ago this began to be irritable, and, yielding to the sensation, the patient has from time to time freely scratched the surface. At the present date the part originally occupied by the mole is covered by a prominent fleshy mass, about two inches high in the thickest part; it is very tender to the touch, and is still growing. There are no signs of malignancy about the disease.

The only plan of treatment for moles is by complete removal either with the knife or by some kind of cautery. As a caustic, I should recommend the strong nitric acid as the most manageable and efficient means of cure; indeed it is the only one that I have employed, and it has given me the best results. I need scarcely say that in infants the only moles that require treatment are such as by their size or position are manifest blemishes.

A SHORT GENERAL ACCOUNT OF THE EPIDEMIC OF FEVER AT MAURITIUS.

By WILMOT H. F. POWER, B.A. LOND.,
ASSISTANT-SURGEON TO THE 2ND BATT. 13TH LIGHT INFANTRY.

THE fever that has prevailed since the beginning of this year, and which still continues to some extent, has apparently attracted considerable attention in England, chiefly on account of the great mortality attending it. I have thought a few general remarks on its nature and the causes of the mortality might be acceptable to the profession at home, and having been now for seven months in constant attendance on the sick of my regiment, I may lay claim to some right to offer an opinion on the subject.

Nature of the fever.—The profession in Mauritius is agreed on this point—viz., that the fever was *malarial*; but there was, when I left the colony at the beginning of June last, a difference of opinion as to the type of the disease, some holding the greater number of cases to be "*remittent*," and others holding it to be "*intermittent*." For my own part, amongst the military and some few civilians I attended, by far the greater number were of intermittent fever, and I believe I am warranted in stating that my friend, Dr. S. Stone (the resident surgeon of the Government Civil Hospital), and consequently practising amongst a large number of coloured people, holds the same view as to the relative proportion of cases of remittent and intermittent fevers. Perhaps this discrepancy of opinion arises from the enormous amount of work thrown on the hands of the medical men, who were consequently often able to spend only a few seconds, and that not every day, on each patient; and this, with the condition of numbers of the patients, who, from natural ignorance or from the effects of the fever, were unable to give any account of their illness, rendered the diagnosis extremely difficult. And, again, I observed some tendency to put down at once as remittent fever any case which showed great prostration, vomiting, coldness of body, &c., whereas such a condition was brought about by both intermittent and remittent fevers if allowed to pass unchecked beyond a certain time. In such cases almost the only method to be sure of the type was to have yourself seen the patient previously. Perhaps, on the whole, it would be advisable to call this fever simply "*malarial fever*," especially as in some cases the patient was struck down at once, and never showed a symptom of intermission or remission.

Area over which the fever occurred.—Speaking very roughly, a line drawn from the northern to the southern end of the island, and dividing it into two portions—viz., a western and an eastern portion, this fever may be said to have been confined to the western portion alone. If we cross the island (a distance of some thirty-two miles) from Port Louis, which is near the centre of the western coast, to Mahebourg, which occupies a similar position on the eastern coast, after ascending some distance we lose all traces of the fever in the high central lands of the island, and then, descending to Mahebourg, we still find a district free from fever. This is a most striking instance of the extremely local character of the fever—two towns within a short distance of each other, and both on the sea-coast, the one teeming with the fever-poison, the other absolutely free. Till more certain intelligence reaches us of the

exact range of the fever, I refrain from endeavouring to explain this remarkable difference, beyond noticing the fact, that on the western coast there is a band of flat and damp, and in some places, marshy land, all of it undrained and much uncultivated, while on the eastern coast the mountains run directly, or by long slopes, almost to the sea-shore.

Causation.—Taking the usually accepted factors in the generation of malaria, we find the hot and rainy season of 1866–67 in Mauritius to have been characterised by unusually high and prolonged temperature, with an unusually small rainfall, acting on undrained soil full of decomposing vegetable matter, or on streams which were in reality ponds. (I may add that the fever was as fatal, if not more so, in some country districts as in the dirty, ill-drained town of Port Louis.) This amount of decomposing vegetable matter has been always present, but this year, apparently, it was first exposed to the influence of sufficient heat &c. to form the malaria. It may be a question, too, whether, owing to the large amount of forest that has been cut, a greater surface of undrained soil was exposed to the high temperature than was the case only some fifteen or twenty years ago. This is, I believe, the first time that malarial fever has broken out (to any extent, at least) in Mauritius; and I suppose the case is almost unique—namely, the appearance of a non-contagious fever (endemic in a large portion of the globe), for the first time, in a portion only of a small island near no extent of land, and, moreover, an island exposed for seven or eight months in the year to the cool south-east trade-wind, the only other winds being, as a rule, the usual local land and sea breezes.

Causes of its fatal character.—I have no doubt these were, (1) the want of quinine; and (2) the want of regular medical attendance, of proper nursing, of food, &c. (I might add a third reason—namely, the opinion published in one of the local papers by, I regret to say, a medical man, that he had found *very* large doses of quinine caused dysentery and other internal disorders, and that he never used it; being content with a mercurial purge, followed by ammonia mixture!) Comparing the mortality of my regiment with that of the civil population for the three months January, February, and March: Among a strength of 673 men with some 540 admissions from malarial fever, none died; and of 44 women and 79 children, one infant only died. (These figures refer only to the three months above specified, and while the regiment remained in Port Louis.) The exact number of the civil population who died during this time cannot be told, as it was only from the beginning of February that exact mortality returns were made; but it will suffice to say that many thousands died. These figures, I think, go far to prove the causes I have given above for the fatal character of this disease; and that the slight mortality amongst the men, women, and children of this regiment was due to a greater amount of medical attendance (the surgeon and myself), with nursing, food, and stimulants. These the mass of the poor civil population did not receive, in spite of the unwearied labour of the medical men. Further, we know well that in such incurable diseases as cholera and yellow fever no such difference is found between the military and civil population, when equally exposed to the disease. As this is an important element in estimating the actual amount of danger to life due directly to the character or type of the disease itself, I may give a few instances which occurred in my own practice. One night, while making my hospital rounds, I found three men apparently asleep, but who were comatose; they had been free from fever and tolerably well during the day, and the orderlies believed they had gone to sleep. A quantity of quinine was immediately put down their throats, followed by wine, &c. These three men are alive and well now; had they been poor blacks, or even of better rank, they would probably have been found dead next day or subsequently, in huts or in the cane fields or in the roads. These were cases of intermittent fever which had apparently recovered from a first mild fit.

There is not the slightest necessity to try to explain, as has been already done, the mortality of this epidemic by calling it a malignant fever, or one which early assumed a grave typhoid character; but given an atmosphere, and perhaps water, charged with a large amount of malaria, given also insufficient medical attendance, with want of nurses, or sick attendants with want of quinine (and consequently many repetitions of the fever), want of suitable food and stimulants—given all these factors, and the mortality of the fever is perfectly explained, without resorting to any loose jargon of the malignancy or the like of this particular epidemic.

At the end of March I saw a gentleman who had been ill with fever some days. His regular medical attendant had

been unable to see him for two days. He had been taking a small quantity of ordinary claret, with a few grains of quinine, for that time, and had taken no food. When I saw him he had fever, with low wandering delirium, &c. *A large dose of quinine I gave at once*, and ordered the usual stimulants and plenty of sherry. I saw this gentleman some weeks afterwards, well: he had recovered from the day of my visit. Now, had he not got medical advice &c. that night, he would probably have died.

To give one more instance. I was called to see a strong, hearty Englishman, in good circumstances, who had been only about a year in the island. I found him dead in his room, while in the adjoining room was another, who, hearing his friend groaning during the night, got out of bed, but fell down on the floor (he had the fever also), and remained there till put to bed again next morning by some friends who came in. These two men had no servants; they were dead or had run away. I am perfectly certain that, with a proper attendant to give him medicine &c., that man would not have died. He had been up and about the day before, and went to bed saying he was much better, and felt no more fever. He was not in the slightest degree emaciated or cachectic.

Conclusion.—I trust I have now given a fair view of the nature of the fever, and of the causes which have led to a mortality far greater than any of the epidemics of cholera which have at various times attacked that colony. I may add that there was nothing like yellow fever during my stay there—i. e., till the beginning of June, 1867. Till the various returns which, I believe, have been called for by the civil government of Mauritius have appeared, together with the various meteorological and other returns, it would be premature to attempt a full account of the fever, including its exact limits, and the relative mortality in the different districts.

Treatment.—My treatment, which I learnt from my esteemed teacher, Deputy Inspector-General Maclean, was simply quina in full doses (as a rule, eighteen to twenty-four grains were sufficient) during the intermission or remission, but giving it at once if dangerous prostration with a tendency to coma were present. I differed from, I believe, most English authorities in giving it in the form of pill, as I found it was retained better in that form in cases where there was gastric irritability, also because it had to be given as carefully as possible for fear of running short of it, and in the form of pills it was far more convenient to carry round with one to administer to 200 or more patients than in mixture. Ordinary diuretics were given, but not as a matter of routine; a good purge in some cases, with nitrate of bismuth to allay the vomiting. I seldom gave an emetic unless in the case of a man who had been drinking hard shortly before my seeing him.

Portland Barracks, Aug. 5th, 1867.

A Mirror

OF THE PRACTICE OF MEDICINE AND SURGERY IN THE HOSPITALS OF LONDON.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum, tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

GUY'S HOSPITAL.

A CASE IN WHICH THE BONE EXTREMITY OF AN ELASTIC CATHETER FORMED THE NUCLEUS OF A PHOSPHATIC CALCULUS; LITHOTOMY.

(Under the care of Mr. COOPER FORSTER.)

We had the opportunity, on July 30th, of witnessing the operation in the following case, which was one of more than ordinary interest.

A middle-aged man was placed under chloroform on the table. Mr. Forster, in some remarks to his audience, called attention to the peculiar circumstances connected with the presence of the stone. The man, he said, had been in the

habit of introducing a catheter into his bladder for the last seventeen or eighteen years; for what reason it was hard to say, for he had no stricture or paralysis. Just a twelvemonth since, on withdrawing the instrument, he found that its extremity, which was made of bone, had been left behind. Three days after this he applied to Mr. Forster, who, on introducing a sound, struck some foreign body in the bladder. On the next occasion, however, after the lapse of a few days, he failed, on careful examination, to discover it. As the alleged object was of bone, and he (Mr. Forster) did not know how far the urine might have affected it, he instituted an experiment on the subject. He kept a piece of bone in urine of a temperature of 100° for twenty or thirty hours. No effect whatever was produced upon the bone; he concluded, therefore, that no explanation of this kind would account for the disappearance of the object. He saw the man in a few months, and he was not suffering any inconvenience; but a week since he applied, when he had marked symptoms of stone. The sound now readily detected a calculus.

Mr. Forster proceeded to operate by the lateral method, using Aston Key's straight staff. The patient was not bound, but was held in position by two assistants, and the operation was performed without any difficulty, the patient remaining as still as if ligatures had been employed. With the gush of urine which escaped when the bladder was opened Mr. Forster succeeded in grasping the calculus with the forceps. The stone, however, was so very soft that, despite great care, it broke down. It was removed, however, with great ease. Examination showed that a cylinder of bone about an inch and a half long, and of the calibre of a No. 10 catheter, formed a nucleus, which had been invested with a phosphatic deposit. The patient has not had a bad symptom since the operation.

KING'S COLLEGE HOSPITAL.

PARTIAL ANCHYLOSIS OF THE JAWS, THE RESULT OF AN INJURY; TREATMENT BY FORCIBLE DILATATION.

(Under the care of Mr. SPENCER WATSON.)

In the early part of March, 1867, a girl was brought to the hospital by her mother with a partial ankylosis of the jaws, which allowed a separation of the front teeth of only half an inch. This limitation of movement was attributed to an injury to the jaw in a fall from a first-floor window six years previously. Ever since that time there had been difficulty in masticating, and the child was now almost continually munching soft food, such as bread-and-butter, in order to get sufficient nourishment.

The lower jaw, when opened, moved somewhat obliquely to the left, and this obliquity was probably due to a shortening of the ramus on that side. Some amount of thickening could be felt near or at the temporo-maxillary joint on that side, and this may very possibly have resulted from a fracture in the neighbourhood of the joint, with subsequent inflammation of the joint itself.

On March 9th, acting on the supposition that there were some fibrous bands around the joint which checked its movements, Mr. Watson had the patient put under the influence of chloroform, and then, thrusting a screw-dilator as far back as possible between the molar teeth on each side, commenced forcible dilatation. The screws were worked alternately on each side, and with very gradual and slow movements; and the jaw very slowly yielded, but without any audible crack or other evidence of rupture of fibrous bands.

The operation was not followed by any bad symptoms, and the only thing complained of was tenderness and pain in the region of the left joint for the first two or three days, with a very trifling amount of swelling. After this swelling had subsided, wedges of cork were placed between the molars on each side, and allowed to remain there constantly (excepting at the child's meal-times), with the view of preventing a return of contraction. The result was, that the jaw could be opened freely, so as to leave an interval of an inch and a quarter between the incisor teeth, and to give much more freedom of mastication.

The same operation was repeated on March 30th, but without giving any further mobility. The result, however, was considered by the patient and her friends as very satisfactory, as the child was now able to take her meals with very little difficulty.