combination of 40-inch focal length was used by reversing the lens and putting the flint and crown about 1 mm. apart. A number of views were shown to illustrate the advantages of such lenses.

Several pictures were taken with an ordinary spectacle lens, 34-inch focus, stopped to about $\frac{3}{8}$ -inch diameter, which were very good indeed and scarcely distinguishable from those taken with the telescopic lens or a telephotic combination. Such a lens can be bought for ten cents.

J. F. Kemp spoke of the comparative uselessness of ordinary photographs in the study of mountain geology, and believed that such a simple camera would be of great value in field work. The paper was discussed by others.

WM. HALLOCK,
Secretary of Section.

NEW YORK ACADEMY OF SCIENCES—SECTION OF
GEOLOGY, FEBRUARY 15, 1897.

THE first paper of the evening was by Mr. F. C. Nicholas, and was entitled 'Explorations in the Gold Fields of Western Colombia.' Mr. Nicholas described the curious placers in western Colombia, which, while extremely rich in limited portions, are of very low grade when considered as extended propositions. The gold gravels occur along the western base of the Andes Mountains, and extend from the Gulf of Darien southward, up the Atrato River, to Quibdo. They are also found to the southward of the San Juan River and are in the form of terraces similar to the terraces of the Atlantic States. After the formation of the auriferous gravels the speaker supported the view that igneous intrusions and upheavals had cut them off from their parent hills in the interior and had recognized the drainage, so that the streams do not now head in auriferous rocks. The surface geology indicated that the Gulf of Darien formerly extended a long distance up the valley of the Atrato. Quite detailed descriptions of the gravels and of the character of the terraces were given in the paper. Mr. Nicholas described a route by which a man could sail in a canoe from the Atlantic to the Pacific in the wet season by going up the Atrato River to the Quito River, thence

to the divide, which is in a series of swamps, thence into the San Pablo River and on down the San Juan to the Pacific.

The second paper of the evening was by Professor R. E. Dodge, entitled 'Recent Work in Physiography.'

Professor Dodge gave an outline of De Laparent's 'Leçons en Géographie Physique,' of Sir John Lubbock's 'Scenery of Switzerland,' and of two recent papers, one by M. R. Campbell, entitled 'Drainage Modifications and their Interpretation,' and the other by C. F. Marbutt, 'On the Physical Features of Missouri.'

The last paper of the evening was by A. A. Julien, on the 'Sculpture and Sorting of Sands.' The speaker, by means of lantern slides, illustrated various varieties of sand and their chief methods of origin and their composition. After citing the schemes for the classification of sands advanced by Zirkel and Daubrée he gave one of his own which was more elaborate and was partly based on the method of origin and partly on the physical characters.

J. F. KEMP,
Secretary.

AMERICAN CHEMICAL SOCIETY—NEW YORK
SECTION.

THE meeting was held at the College of the City of New York on Friday, February 5th, at 8:30 p. m., Dr. Wm. McMurtrie presiding, and about fifty members present.

The first hour was occupied with the 'Discussion of the Relations of the Section with the Scientific Alliance.'

Professor Breneman opened the discussion. Dr. Wiley described the work done by the Affiliated Societies of Washington, the advantages resulting from cooperation and more which might result from a little additional effort. He stated that, with possibly one exception, the Washington Societies were all strictly professional.

Professors Sabin, Doremus and others spoke strongly in favor of the Alliance; others thought the promised advantages had not materialized and that the returns were not proportionate to the annual subscription.

Dr. H. W. Wiley read a paper on the 'Value of Foods and the Methods of Ascertaining it,'

pointing out the disparity, at least until very recently, in the relative amounts of investigation and interest bestowed upon 'man foods' and animal foods. He described the classes of foods rated according to their fuel values, digestibility, etc., and noted the divergence between price and actual food value of many articles, some of them not luxuries.

The papers announced on 'Volumetric Estimation of Lead,' by J. H. Wainwright, and 'Electrolytic Production of Alkali Nitrites,' by Wm. M. Grosvenor, were held over until next meeting.

DURAND WOODMAN,
Secretary.

GEOLOGICAL SOCIETY OF WASHINGTON, 58TH
MEETING, JANUARY 9, 1897.

MR. WHITMAN CROSS read a paper on 'The igneous rocks of the Leucite Hills and Pilot Butte, Wyo.,' in which he stated that the rocks of the Leucite Hills occur as surface flows and in volcanic necks or plugs. The leucite rock described by Zirkel, 20 years ago, is the least abundant type in the region, the other varieties containing more or less potash feldspar, as pointed out by Kemp in his recent communication to the Geological Society of America. Chemical analyses of various rock types were submitted, together with analyses of the pyroxene and mica.

The rock of Pilot Butte, an isolated point near the Leucite Hills, was also described and its chemical composition shown by an analysis. This rock is closely related to the leucite rocks, although containing much glass.

In a large cavity of the leucite rock was found a quantity of potash nitre, and on a protected face of the Boar's Tusk, a volcanic plug, was observed a white coating of soda nitre. The origin of these nitrates, whose mode of occurrence is so unusual, is not explained by any observations made.

This communication will soon be published in some scientific serial.

Mr. W. Lindgren read a paper on 'The Granitic Rocks of the Sierra Nevada,' in which he called attention to the large areas of intrusive granitic rocks occurring along the Pacific coast and to the fact that these intrusives are of com-

paratively recent date, probably early Cretaceous. A map of the distribution of the various kinds of granitic rocks in the northern part of the Sierra Nevada was exhibited. It was shown that, while some true granite exists, the largest mass is made up of grano-diorite, a rock intermediate between granite and diorite or, more accurately, intermediate between a quartz-mica-diorite and a quartz-monzonite, recently defined by Brögger.

W. F. MORSELL.

U. S. GEOLOGICAL SURVEY.

BOSTON SOCIETY OF NATURAL HISTORY.

A GENERAL meeting was held January 6th, eighty persons present.

Mr. A. W. Grabau read a paper on the sand-plains of Truro, Wellfleet and Eastham. (For an abstract see above p. 344.)

Professor N. S. Shaler, in commenting upon Mr. Grabau's paper, said that his observations agreed with those of Mr. Grabau as to the origin of the sand plains. The slopes are due to a complexity of causes and frequently cannot be discriminated. The hypothesis of fresh-water lakes and the ponding of streams was rejected as inadequate. Professor Shaler said that the agency of ants in the formation of these sand plains was very great, and should be considered in relation to any theory accounting for them.

Mr. J. B. Woodworth spoke of the difficulty of making out the internal structure of sand plains; he had observed that the gravels near the head were coarser than those from other parts. Mr. Woodworth compared the sand plains of the Cape region as described by Mr. Grabau with those he had studied in the Narragansett Bay district. The hypothesis of fresh-water lakes applies equally to the Narragansett Bay district. The kettle holes indicate masses of ice after the melting of the ice sheet.

Prof. W. M. Davis claimed that the slopes could be discriminated, but that they should not be solely relied upon.

Professor Shaler contended that the slope depended upon the material, and that the original angle is unreliable until the material is known. The difference of height of sea level should be