our former paper * on the great *Paradoxides* such pleuræ were described of great length. In the present species they are shorter, but still appear to be distinctly separated by simple grooves from the circular appendix as in *P. Davidis*. In *P. Loveni* they are perfectly soldered, and are indicated only by the increased number of lateral grooves and marginal spines of that fossil; and in *Anopolenus* also the limb is many-ribbed and the margin spinose.—J. W. S.

EXPLANATION OF PLATES II. & III,
(Illustrative of Fossils from the Menevian Group.)

PLATE II.

Figs. 1 & 2, 4 & 5. *Conocoryphe applanata*, Salter. Natural size.

3. — *perdita*, spec. nov. Natural size.

6. Embryo of *Conocoryphe applanata*. Magnified.

7. *Conocoryphe humerosa*, Salter. Natural size.

8. — *bufo*, Hicks. Natural size.

9–12. *Paradoxides Aurora*, Salter.

PLATE III.

Paradoxides Hicksii, Salter.

4. ON QUATERNARY GRAVELS. By A. TYLOR, F.G.S.

(Read May 6, 1868†)

[PLATES IV.–IX.]

IN continuation of a paper on this subject in the Quart. Journ. Geol. Soc. vol. xxiv. p. 103, I now submit a number of measured sections selected as instances of the deposition of gravels of different character under different circumstances both of height above the sea-level and distance from the sea.

The illustrations and remarks in this paper relate to the unfossiliferous gravels of the rivers Taff, Rhondda, and Cynon, in Glamorganshire, South Wales, to those of the Aire, in Yorkshire, and to the fossiliferous gravels and brick-earths of the lower part of the Thames and Lea rivers, in the counties of Kent and Essex.

One section of the fossiliferous gravel of the river Avon, at Salisbury, is given for the purpose of comparison with those of the Thames; and a section across the valley of the Aire, in Yorkshire, is compared with those in the valleys of the Taff and Rhondda.

A map of part of Glamorganshire, and a longitudinal section of the bed of the river Taff, with the heights along its course measured above the height of spring tides at Cardiff, have been prepared to define the exact locality of the Welsh gravels alluded to. A similar map and sections of the course of the river Aire, in Yorkshire, with heights measured from the Ordnance datum, are given for comparison with the Welsh illustrations.

A sketch-map of part of the river Thames, and of the adjoining

* Quart. Journ. Geol. Soc. vol. xx. p. 235, pl. 13. f. 2.

† For the Discussion on this Communication see Quart. Journ. Geol. Soc. vol. xxiv. p. 456.

counties of Kent and Essex, and separate maps, on a large scale, of the Chalk-pits and Brickearth-pits of Erith, Crayford, and Grays define the exact position of the sections across the valleys of the Thames and in the pits; and the heights explain their relation to the present river Thames. Sections of the raised beach and mammaliferous gravel at Brighton, and of the fossiliferous gravel and raised beach at Sangatte, showing the contour and the modelling of the upper surface of the gravel-series, have been also drawn. A comparative Table of fossil shells &c. from the above localities is also added.

The evidence of numerous sections teaches us that, prior to the deposition of the gravel, there was a land-surface smoothly denuded by rain and streams so as to form a perfect system of principal and minor valleys, the ground sloping from higher to lower points, so as to admit the rainfall to flow with the minimum of obstruction into the side-valleys and thence into the ancient Thames. The subsequent deposition of the gravel-series did not in any way alter old lines of drainage; but where concavities existed the new deposit had a tendency to fill them up with a thicker stratum of material than was spread over the general surface of the chalk or clay. Thus the Quaternary beds reach a thickness of 80 feet at the maximum, while the average is perhaps only 25 feet in the whole district. Some boulders on the upper part of the gravel-deposits reach to many tons in weight; and they are as large or larger than those in any part of the series, so that there is evidence of as great intensity of pluvial action at the end as at the commencement of the Quaternary series.

Except where the old river-channels and concavities are filled up, the contour and the modelling of the upper surface of the gravel-series resembles that of the clay or chalk on which the lower gravel rests, and is perfectly adjusted to carry off the rainfall occurring in the later part of the gravel-period into the ancient Thames without any impediments. The contour of the land is such that it could only have reached its present form by pluvial and fluvial action, and not by marine denudation. Many of the minor valleys in which gravel and brick-earth were deposited with each flood in the Quaternary period are now dry. After the heaviest rainfall in recent times there is not sufficient force of water to remove vegetation, so as to make any change in the present surface.

We are therefore justified in stating that the character of the denuded surface of the London Clay and Chalk above the level of the Thames is evidence of the occurrence of an enormous rainfall in the commencement of the gravel-period, and that the character of the surface-deposits of gravel is evidence of nearly as much rainfall at the close of that period.

With a rainfall such as we now have, it would be impossible that such widely extended gravel-beds could be spread over an extensive area, and reach to a height more than 150 feet above the level of the Thames. It is equally impossible for the present volume of the Thames to have produced fluviatile beds at all equivalent in size to those of the ancient Thames.

The condition of the beds which rise above the 50-feet level points

rather to pluvial than to fluvial action. Of marine remains in the Thames-valley gravels there are no traces.

In the south and east of England the hills are rarely higher than 1000 feet, and the valleys are deep and wide, and are adapted for rivers 20 times as large (or more) as those occupied by them now.

The banks of the old rapid and large rivers are now separated from the banks of the present small streams by a great extent of *remanié* alluvium; and modern gravel has accumulated in the old river-channels, owing to the decreased velocity of the present streams.

This is well shown in the large flat plains of gravel through which the Lea river flows, where the river, even in floods, only reaches the bottom of the 30-feet bank at Clapton, while the brick-earths and gravels of the ancient Lea reach to 80 feet above the level of the present stream.

Some modern streams have still force enough to cut horse-shoe bends and new courses out of the flat plains which were probably in flood-times constantly overflowed by the old rivers running in the gravel-period; but the covering of recent alluvium in the river-channel prevents any observation of the character of the gravel itself.

Large and deep side-valleys open into the Thames valley, containing small streams or rivers; but I do not know of any instance of a stream in this district having risen high enough, or had force enough, to cut through any of the Quaternary deposits, so as to expose a good natural section.

The absence of such natural sections is of important geological interest; for it is evidence that no marked changes of level occurred during the gravel-period. If elevations of the land had occurred during the latter part of the gravel-period, as has been supposed, and rivers had cut down these beds 30 or 40 feet, then all the side-streams would have had to adjust their beds to the new circumstances of level, and we should have had natural sections in every side-valley, as well as in the Thames valley itself. There are no waterfalls in the district; and every stream reaches the main river by a very gradual course in obedience to a definite law.

The fact that no good natural sections of this kind exist in the Thames valley, or in the valleys of its tributaries, is a proof that neither the principal river nor its confluent streams had force enough to break through the upper gravel and to expose the brick-earth and sands below. It is also evident from the same facts that the change in rainfall, and consequent decrease in volume and speed of all rivers and streams in the district, occurred immediately after the close of the gravel-period.

We find natural sections of gravels along the water-courses of the streams and rivers of South Wales, where the rainfall is still nearly 40 inches, and the side-streams fall with great rapidity to the main rivers from the high land. The materials of these gravels are essentially different from those of the south and east of England; and although the two sets of gravels may be identical in age, there would be as much relative difference between the deposits in high and low

districts in the gravel-period as there is now. I have introduced illustrations of these Welsh gravels, because it would be impossible to form any clear view of the general features of the gravel-period without comparing the sections of the deposits found on the upper part of rivers flowing off high land with those formed along the lower parts of rivers near the sea in the Quaternary period. The general law of deposition will be the same; and it is this law I wish to investigate by the aid of the measured sections. Those sections are chosen specially for description where the rock or clay and the incumbent gravel can be observed together. This gives an opportunity of observing the points or lines on which denudation ceases and deposition commences.

In the London basin, the series of Quaternary strata generally commences with a coarse gravel, a few feet thick, eating into the chalk or London clay in which it lies, and mixing up the clay or chalk with the gravel.

The coarse gravel often passes into coarse sand, and is evidence then that the movement of the water at that point was not sufficiently rapid to transport gravel, or that there was no gravel present to be deposited at that moment.

I have not observed any finely bedded clay or loam in contact with the surface of the chalk or London clay in the London basin; and this appears as if the movement of water over the surface of the clay and chalk had been generally rapid at the commencement of the Thames deposits. No doubt there are many cases where the first deposit has been of fine materials, but I have not been able to meet with them.

Where concavities in the chalk and London clay exist in the neighbourhood of the Thames, and in its side-streams, and are so situated as to be favourable for quiet deposition, the lowest bed of gravel or coarse sand sometimes contains fossils derived from the Eocene beds, and rolled Mammalian remains of a later period. It is succeeded by a series of laminated clays and false-bedded sands, from 20 to 30 feet in thickness, with water-worn materials, and containing fluviatile and land shells, tranquilly deposited with Mammalian remains (only occasionally rolled). No natural sections exist of this class of deposits; but excavations for brick-earth have during the last 30 years opened out several of these fluviatile beds.

Non-fossiliferous gravels are often clearly contemporaneous with these fluviatile brick-earths. If on the same horizon, they are sometimes interstratified with the fossiliferous beds; but the non-fossiliferous gravels reach to much higher levels, and are deposited at much steeper angles, and contain materials not perfectly washed or water-worn. They include masses of Thanet sands, of plastic clay, of London clay, and of the fossiliferous bands of the Woolwich series, buried in great unrolled masses in the gravel, as well as water-worn masses of Druid Sandstone, derived from strata very near the spot of deposition. These materials were evidently washed in by heavy floods, not confined to the valleys, but passing over the whole surface of the land, tearing up the ground and carrying

it to lower levels into the valleys. At page 124, *Quart. Journ. Geol. Soc.* vol. xxiv., I described such a flood in Sind in the year 1866, when a rainfall of 24 inches occurred in ten hours, laying all the surface of the land under water, and moving very great weights a considerable distance. The ancient river-deposits of the Thames basin associated with this flood-borne detritus are generally covered by an upper bed of well-rolled gravel, not exceeding 10 feet in thickness. This bed is often only 3 or 4 feet thick, and lies on the frequently eroded surface of the stratified brick-earths and clays of the middle part of the Thames Quaternary series. Even when for a short distance this surface-bed contains few pebbles, there is no difficulty in identifying it; for it will soon be charged again with pebbles. This thin upper coating of well-rolled pebbles, enclosed often in a stiff brown clay, but sometimes in sand or loess, extends from the river-banks to near the tops of the hills over all the other parts of the series of gravels, and often over the London clay and chalk; it covers a most extensive area, forming a marked contrast to the middle series of laminated brick-earth, false-bedded sands, and stratified clays, which are more confined to the low levels near the rivers or brooks. This bed of upper clay and pebbles slopes to the lower ground (I estimate on an average) at 1 in 260, and falls therefore about 20 feet in a mile. It is washed over the surface of the land smoothly, removing all abrupt heaps of gravel, and leaving no greater irregularities of the surface in the London basin than occasional low terraces. The smoothness and perfect manner in which the gravel is deposited is shown by the fact that the surface of the London clay or chalk is covered, notwithstanding inequalities of the surface, by a bed only 25 feet thick over such a large district.

Fig. 1.—*Map of part of the course of the river Aire, Yorkshire.*

Map of the course of the River Aire, from Bingley to Castleford. The letters refer to the lines crossing the river, and are for reference to the longitudinal section (fig. 2) of the River Aire, or rather of a line supposed to be drawn at each point of its course 8 feet above the full-water-mark of the river. This map has been accurately reduced by the pentagraph from the one-inch Ordnance sheets.

The river Aire rises near Malham Tarn, in Yorkshire, in the Car-

boniferous Limestone, one branch coming from a cave near Gordale Scar, then flows over the very thick formation of Millstone-grit, reaches the Coal-measures between Shipley and Bingley, and, passing over the Magnesian Limestone and the New Red Sandstone, empties itself into the Humber, or a branch of the Humber, near Goole. The valley-gravel at Leeds is famous for the fine remains of Hippopotamus found in 1852. The bottom of the valley through which the Aire flows is from one-half to one and a half mile wide, and is coated over with a deposit of gravel and rolled boulders of unequal thickness and elevation, consisting principally of limestone at the upper part near Malham and Skipton, and of Millstone-grit and sandstone near Leeds, with some limestone boulders. The Bingley gravel consists of Millstone-grit, sand, rock, and limestone in large percentage. Some boulders of limestone are found near the mouth of the river near Goole; but at Bingley, fourteen miles from Skipton in a direct line, they form a considerable percentage of the whole gravel. Until the Leeds and Liverpool Canal was made between Leeds and Skipton, this valley-gravel was turned over for the purpose of removing the limestone boulders for burning into lime. There are numerous remains of old limestone-quarries and lime-kilns at Bingley, although it is ten miles from the nearest limestone rock.

Fig. 2.—*Map of part of the river Aire.*

Fig. 2 is a plan of the Aire valley between Bingley and Shipley, showing the lines of sections A B and C D from the sandstone-rocks on each side of the valley. (See Pl. IV. figs. 2 and 3, for details of the sections; and to these I will now refer.)

The section (fig. 4, Pl. IV.) through Ashfield, half a mile east of Bingley, exposes a thick gravel- and boulder-deposit, forming what are called the Old Hills. This gravel has been worked over for the blocks of limestone it contains, leaving an irregular surface.

The gravel is 50 feet above the river Aire in the thickest part; and no rock has been met with to prove how much lies below the level of the river-bed; but the town surveyor, who has carried out the drainage of that town, informs me that the rock is probably from 30 to 40 feet below the level of the river.

Considering the width of the valley, nearly three-fourths of a mile, we might estimate the gravel over a width of 1000 yards to be of an average thickness of 20 yards; 20,000 yards at a weight of $1\frac{1}{2}$ ton to the cubic yard, is 30,000 tons; there would be then 30,000 tons of rock and gravel to each yard run, or about 53 million tons of gravel to each mile of river, supposing the section, fig. 3, at Bingley to contain the average amount of gravel in that valley.

This quantity of 53 millions of tons of gravel per mile is apparently not out of proportion to the power and dimensions of a river able to flood its valley to a height of 50 feet over a surface a mile wide with a gradient of from 3 to 10 feet per mile.

The escarpments, or gravel-banks, 50 feet high and 200 yards wide, in fig. 4, appear to mark the ordinary limits of the ancient river, as the sides of the gravel-bank are laid in many parts in the form of a steep escarpment sloping to the current at an angle of from 25° to 35° , while the top of the gravel-bank is flat, except where it is hollowed out by another part of the river-channel more or less deeply. A bank of 10 feet above the Aire is sufficient now to retain the water, except in the heaviest floods. Contrasting the floods of this river at the present time with those in the period when the gravels were deposited, it would appear that a flood-line of 10 feet now replaces one of 50 feet. The cubes of these flood-lines would probably represent the proportion of the volume of water flowing down the valley in the two cases. Then 10^3 is to 50^3 as is 1 to 125. The volume of water would on that hypothesis now be only the $\frac{1}{125}$ of that in the gravel-period.

Section (Pl. IV. fig. 5) along the line C D, through Rye-Loaf Hill, one mile E.S.E. of Bingley, shows the outline of the gravel-deposit lying on Carboniferous sandstone and shales. This part of the Aire gravel, along the line C D, reaches also to a height of 50 feet above the present level of the river; it does not appear to have ever been worked for limestone, and is about the same height above the river Aire as in A B. The escarpment of the gravel at Rye-Loaf Hill is well marked, and slopes to the river at 25° ; the top is flat, and the form such as could only be produced by a river occupying during floods the whole width of the valley, the water rising at least 50 feet above the present flood-level. The limestone boulders forming a large portion of the Bingley gravel are so well rounded that they indicate long-continued rolling in a river-channel.

There is a swamp to the east of Bingley church, in which remains of oak-trees have been found in peat, and below the surface a stratum of freshwater shells of existing species, according to my informant. This part of the section appears to be identical with the Hippopotamus-bed at Leeds.

Mr. A. Harris, jun., of Ashfield, found, in 1868, pieces of rolled

chalk several inches long below the peat, during an excavation made for foundations of a chimney. This chalk may have fallen into fissures of the limestone during the denudation of the Cretaceous series, and have been washed out again at a comparatively recent period.

The quantity of gravel now left in the valley of the Aire may only represent the portion of water-rolled *débris* arrested *in transitu* to the sea, at a period when there occurred atmospheric changes of a character and importance sufficient to reduce the flow of water to its present insignificant amount.

My object in selecting the gravel-deposit of the river Aire as an illustration for examination is because that river flows over a variety of distinct strata, and yields evidence of the transport of a number of large boulders from its source to near its outfall. Then its source is in a district in which occur the celebrated caves of Kirkdale, so rich in the large Mammalia, such as Elephant, Rhinoceros, Hippopotamus, and Hyæna, and containing in the upper part of the cave-deposits human skeletons and pottery.

Although at Bingley itself no Mammalian remains have been noticed, yet at Leeds, twelve miles lower down the river, the following description of the discovery of *Hippopotamus* sufficiently identifies the Leeds gravel with the cave-gravel of the limestone district by means of its organic remains, and with the Bingley gravel by means of mineral character and stratigraphical position.

Mr. H. Denny, of Leeds, records the discovery, on the 3rd of April 1852, of great portions of the skeletons of two or three large specimens of *Hippopotamus major* at the angle formed between the river Aire and Wortley Beck, close to Holbeck Station, Leeds, at a height of 115 feet above the sea, and 20 feet above the banks of the Aire.

Mr. Denny writes, ('West Riding Geological Society Reports for 1854,' p. 325):—"The bones were discovered in a dark-blue sedimentary clay almost approaching mud, and appear from their condition to have belonged to animals which had lived and died in the immediate vicinity, and were subsequently drifted, together with fragments of trees to the bottom or lower part of a swamp, found only in this particular bed of clay, confined to one portion of the field; but the clay becomes much thicker, and reaches 10 feet in thickness as it approaches this spot, thus clearly indicating it to have been lower than the remainder of the brickfield."

Beds of clay of variable quality resting upon gravel, with occasionally large boulders, some of which contain impressions of *Stigmaria* and fragments of trees to the depth of 10 feet, yielded the bones of three individuals of *Hippopotamus major*, whose skeletons seem to have been complete when deposited. The canine tusk of one individual was 18 inches long, by $2\frac{1}{4}$ broad. Mr. Denny observed that the vertebral column of one Hippopotamus extended in a line across the trench; the ribs appear to have been *in situ*; and from its position he considered that the animal had been lying on its side. Portions of the *Elephas primigenius*, with molar teeth, and fragments

of *Urus* were also met with in the Wortley Beck Fields closely adjoining.

Section at Holbeck.

Greyish sandstone	2 feet.
Yellow sand	2 feet.
White sand with well rolled boulders of grit and sandstone, with an occasional fragment of limestone and chert.....	thickness unknown.

Mr. Filiter, the engineer of the Leeds Waterworks, informed me that the exact position where the remains were found was $53^{\circ} 47' 42''$ N., $1^{\circ} 34' 10''$ W.

It is uncertain whether the Aire-valley gravel represents exactly the beginning of the series of the Kirkdale-cave deposits; but the discovery of the Hippopotamus 10 feet below the surface of the gravel, and only 15 feet above the present level of the Aire, clearly proves that the great mass of the Bingley accumulations was anterior to the time at which the Leeds Hippopotamus lived. If the upper series of these Bingley gravels represented in time the last deposit of 10 feet at Leeds, they would be of the same date as the Hippopotamus, and all the lower part of the gravel at Bingley below the 10-feet level would be older than the Hippopotamus.

In fig. 2, plan of the course of the river Aire between Bingley and Shipley, a curve not very remote from the horseshoe-form is seen.

The section of the river-banks at many points near this curve (see Pl. IV. figs. 4 & 5) present steep escarpments to the river, and thus preclude any supposition of glacial or marine action, as neither sea nor glacier could arrange heaps of gravel with a regular slope facing a river following its course in a horseshoe curve.

When we find a set of gravels in the valleys of other rivers deposited in slopes and levels in a similar manner, and bearing close comparison in contour, condition, mineral contents, dimensions, and position with the mammaliferous gravel of the Aire, it may be safely inferred that we are examining valley-gravels of the same date as that of the Aire, although Hippopotamus and Elephant may not be found in every river-deposit.

The discovery of Hippopotamus in such a nearly perfect and unrolled condition in the superficial deposit of the river Aire, at Leeds, was a most fortunate thing for geologists who discuss the later Quaternary deposits, as it shows that the range of the extinct Mammalia extended to a time since which there have been no changes of importance in the level, nor disturbance, of the superficial deposits in Yorkshire.

At Cottingley Beck, near C, fig. 2, there is a natural section of gravel resting on shale and clay, and containing a block of sandstone moved from a distance. It is 12 feet long, and of very great weight, and only 6 feet from the surface. This could only have been transported down the small valley occupied by Cottingley Beck by a very heavy flood, and is an indication of the amount of rain which must have fallen at the date of the accumulation of the Aire-valley gravels.

Similar animal remains to those at Leeds have been found

associated with the works of man near Bedford, Abbeville, Salisbury, and other localities, so that we are enabled to correlate the gravel of the river Aire with that of a number of rivers which appear to have risen in times of floods from 40 to 80 feet above their present ordinary level in that part of the Quaternary period which I term the "Pluvial period" (Quart. Journ. Geol. Soc. vol. xxiv. pp. 105, 120).

The *Cyrena fluminalis* or *consobrina* is found in the gravels abundantly at St. Valery, at the mouth of the river Somme, in Picardy, and also in the estuary of the Humber at Kelsey Hill. The Aire flows into the Humber.

There is, as might be expected, great difference in the materials of which the respective gravels are composed, arising from the different character of the rocks over which the rivers flow, and from the greater fall of the Aire in its course, from its source to the sea, than that of the Somme. The important stratified valley-gravel at Abbeville, containing numerous species of shells, is represented by the Kelsey-Hill and Hessle beds near the mouth of the Humber at a similar height above the river. The width of the valleys is not so different as their depth; but this is due, as before stated, to the difference in the fall of water per mile in the two rivers Somme and Aire.

There is no good section of gravel now open at Bingley. The limestone boulder-pits have been filled up many years; and the railway is carried near the present river, or in an old secondary river-course of the gravel-period.

GLAMORGANSHIRE GRAVELS.

I shall now describe some measured sections of very thick and important river-gravels in the valleys of the rivers Taff, Rhondda, and Cynon, in Glamorganshire, South Wales.

A longitudinal section of the river Taff would show the rapid character of the river, which falls nearly 500 ft. between Merthyr Tydvil (A, fig. 3) and the sea at Cardiff, about twenty miles. The following heights of the bed of the river Taff, above the coping of Bute Dock, Cardiff, were kindly furnished me by Mr. George Fisher, Engineer in chief, Taff-Vale Railway. These heights were the only ones sent to me relating to the river Taff. Their exact position on the map was also laid down by Mr. Fisher himself, and fig. 3 is a carefully reduced copy of his map. The coping of Bute Dock is 9·82 ft. above the level of ordinary spring tides.

Plate V. is a plan of the river Taff, near Quaker's-Yard Junction. Pl. IV. fig. 1 represents a long section opened out by the Taff Vale Railway (ABC), the position of which is shown on the plan (Pl. V.). The heights at different points have been kindly determined by Mr. L. T. Lewis, F.G.S., of Aberdare, or by Mr. G. Fisher, of Cardiff, and are marked on the section.

The plan and section of the valley of the Taff, between Quaker's-Yard Junction and Aberdare Junction of Taff Vale, represent about one mile in length by two-thirds of a mile broad. The height of the point of the sandstone-rock forming the crown of the hills is about 275 feet above the level of the Taff at Aberdare

Junction; the river under the viaduct is 325 feet above the mean level of the tide at Liverpool. The rails of the Taff-Vale line are 96 feet above the Taff at Quaker's-Yard Junction, and 30 feet at Aberdare Junction, so that the rails are laid on a very steep incline of 1 in 50. On the average, between B and C the fall of the rail is 154 feet, and that of the river about 88 feet between the two stations. The rails cross the river on a viaduct about 96 feet high near the upper end of the incline, and on a bridge 30 feet high at the bottom of the incline.

Fig. 3.—Map of part of the course of the river Taff, Glamorganshire.

	ft.	
Bed of river Taff near Merthyr Station.....	493.0	A.
Bed of river at Quaker's-Yard Viaduct.....	297.50	B.
Bed of river near foot of incline at Aberdare Junction...	230.0	C.
Bed of river near junction of river, Clydach.....	199.50	D.
Junction of river Rhondda at Pontypridd Junction.....	144.0	E.
Bed of river Taff, at Taff's Well, between Llantrissant and Walnut-tree Junction	63.0	F.
Bed of river Taff at Pentyrch Bridge, between Walnut- tree and Penarth Junctions.....	49.0	G.
Bed of river Taff at Melengriffith, between Penarth Junc- tion and Llandaff Station.....	24.0	H.

The section given in Pl. IV. fig. 1 commences at Quaker's-Yard Junction with a bed of loose gravel lying along the sides of the valley, 140 feet high above the river, but probably not of greater thickness than 40 or 50 feet on any part of the rock out of which the sloping sides of the valley are excavated. The gravel is full of very large boulders of the local Carboniferous rock, which is an extremely hard, compact, massive-bedded sandstone of the Pennant series, several hundred feet thick. These boulders are sometimes angular, and sometimes weathered or rolled. There also occur quantities of Millstone-grit, brought ten miles down the valley of the Taff, these being always rolled, and boulders of limestone, with a small but conspicuous percentage of well-rolled pebbles of Old Red Sandstone from the Brecon Beacons and other distant hills. These red pebbles are still more worn and rolled than those of the Limestone and Millstone-grit, the source of which is nearer to Quaker's Yard than the Old Red Sandstone. The matrix is clay and sand derived from the Coal-measures, or from decomposed Millstone-grit.

After passing the viaduct, Pl. IV. fig. 1, the section is continued

through a deep railway-cutting, the rails being 80 feet below the surface of the ground, and 112 feet below the adjoining peak of rock at D.

The escarpment of rock is shown to slope about 43° N., at the northern end of the cutting, B, close to the viaduct, and only about 20° at the southern end, D. Gravel with boulders of Old Red Sandstone, and many other rocks, is spread over the surface of the rock at D, up to a height of 200 feet above the river Taff.

Fig. 4.—*Enlarged Section of Gravel Deposit.*

Fig. 4 is an enlarged view of the section at B along the line of railway, showing the gravel lying in a concavity of the rock forming a part of the Taff valley, at very high angles. Near the rock the dip of the gravel is 35° , gradually becoming less steep until it reaches 18° . The gravel below the rails is not exposed at this point. It will be seen that fig. 4 should have been reversed, to accord with Pl. IV. fig. 1; it is on the natural scale.

Fig. 5 is a transverse section at right angles to Plate IV. fig. 1, and fig. 4, showing the railway-cutting at the point D. As it is

Fig. 5.—*Section across the Taff-Valley Railway.*

not sloped, it gives a good opportunity of studying the position of the gravel, particularly as the surface of the rock is clearly exhi-

bited in the centre of the cutting. From this section it is evident that the Taff river flowed at one time in a straight line from near Quaker's-Yard Junction to Aberdare Junction, excavating the small valley shown in fig. 5, before it formed the horseshoe bend shown on the plan (Pl. V.), through which the river now reaches the same point, G; this valley at D is about 40 feet deep, and 200 yards across, and is excavated out of the wedge-bedded pennants, which dip south-east at a small angle, but have both horizontal and vertical joints well marked, dividing the rock into cubical masses at angles varying from 80° to 85° .

The railway engineers have taken advantage of this natural valley, and cut through it to a depth of 60 feet below the bottom of the gravel.

I wish to call attention to the position of this well-rolled loose gravel, containing many pieces of Old Red Sandstone and Millstone-grit, brought from above Merthyr and deposited at a very high level in the small valley cut out of the sandstone rock at Quaker's Yard. The rails are 393 feet above the datum-line at Cardiff, and the gravel is about 460 feet above the datum-line, and 163 feet above the river Taff, close by. If the horseshoe bend (Plate V.) had been gorged with ice at the gravel-period so as to prevent the passage of the river round the valley BHC, Plate V., then the river might have risen to a height which would have enabled it to pass along the line BDEFC; and the rolled gravel, which is a loose superficial deposit, may have been left by the last great flood.

Returning to Plate IV. fig. 1—before reaching E the rails cross a small lateral valley on an embankment 20 feet high, the surface of the ground dipping at right angles to the rails at an angle of 15° , and so reaching the river at an angle of 30° . The great gravel-cutting EF is now entered, and shows a maximum thickness of gravel of 70 feet.

The great cutting, EF, nearly in the line of the old incline, commences with a series of large blocks of sandstone from 6 to 9 feet long, laid at an angle of 1° N., and forming the last or topmost layer of the gravel. These great pieces of rock have evidently only descended a few yards from the mass from which they were detached. The gravel becomes 55 feet thick at a point (E) where the sand-rock is exposed (see fig. 6, p. 70). The sand-rock in the cutting is visible for nearly 60 yards to a height of 7 feet above the rails at the maximum. This gravel may be observed to be roughly parallel to the top surface of rock, which slopes gently to the south. The greatest thickness of gravel with large boulders occurs near F, reaching 70 feet. The slope of the surface is given at different points.

The larger blocks are often arranged in lines or planes of deposition, forming a distinct measurable angle. In the larger gravel-cutting, EF, they are not in any case more than 15° ; near the viaduct they reach an angle of 35° near the rock. These planes follow, and are nearly parallel with, the surface of the rock which crops out at E, and are so arranged that the last bed of gravel forming the surface, being also the last superficial deposit, is also almost parallel with the

top surface of the bed of stones on which it lies, just as the last bed is almost parallel with the mean angle of the surface of the rock on which it is spread by the action of water (see E). This section is to natural scale.

Fig. 6*.—Section along cutting of Taff-Valley Railway.

The exposure of rock at many different points of the horseshoe bend near the piers of the viaduct, and in the bed of the river, adds to the interest of the sections at Quaker's Yard. I shall now describe the character of this large rock-basin in which the gravel under consideration is contained. The first place at which I observed the rock was near the river, where a small quarry is worked close to one of the piers of the viaduct. The dip is slightly south-east, and the stone is divided by nearly vertical joints; there is another quarry of the same character on the opposite bank, between the tramway-bridge and the south-east end of the railway-viaduct, but close to the latter. In following the tramway round the horseshoe bend the rock is frequently visible; in the river-bed it is concealed by a coating of gravel as far as the junction of the stream at the north-east point of the bend. The gradient of this small stream is steep, and the water enters the river with some velocity, running nearly parallel with the Taff near the junction. The water of the Taff, up to the junction, is smooth, and flows quietly over a gravel bottom; but the lateral stream, coming in with added velocity, causes the stream of the Taff River almost to split into two different paths, one taking the east, and the other, or larger portion, crossing the bed of the river, and hollowing out a deep channel in the rock close to the west bank. The river flows below the junction at H entirely over rock, the gravel not being able to rest on account of the velocity of the current.

The action on the river-bed corresponds with what I have represented in other sections that I have made, which show that the work of denudation, or cutting out of the solid rock, is principally due to running water, either in the lateral streams themselves, or in the main river immediately after the side-stream has emptied itself into it. Of this I shall bring forward other examples at another

* Fig. 6 is also reversed. See Plate IV. fig. 1 at E.

time, to show that the work of denudation performed by the main stream flowing at the bottom of a deep valley becomes insignificant compared with that of the side-streams flowing into the main river.

The river itself only operates on a narrow slip of ground at the bottom of the valley, although, by the momentum it possesses and its constant action, it has a powerful denuding force. The side-streams flowing into the river, on the contrary, although more intermittent, act on the whole of the remaining surface of the valley, either through themselves or their smaller tributaries. They thus denude an area twenty times as great (or even more) as the river itself can touch. The side-streams are less fixed in position, and run with many times the speed of the river itself. Their power as an agent of denudation is not in proportion to the average amount of water which they contain, but to their speed in flood-times.

The side-streams occupy channels of infinitely greater length than that of the main river. The points where the greatest force of water is employed now at the bottom of the Taff valley are where the water in the side-streams and the main stream unite, as at the north-east corner of the horseshoe bend at Quaker's Yard at H.

At the south-eastern corner is a large quarry close to the road, in which 60 feet of massive-bedded sandstone is seen. It is also seen in the river-bed.

As the rails near the viaduct are 96 feet above the level of the river, and as rock is seen 120 feet above the rails, and 96 feet below them, we have at least 216 feet of solid sand-rock exposed in these sections, without any intermixture of clay or shale. This is an important fact, as we may observe that a horseshoe bend of as perfect a form as in alluvial soil may be excavated in solid rock. Mr. Fergusson has attempted to prove that the sizes of the bends and curves in rivers depend upon the velocity of the river, and that a curve in a river falling 10 feet in a mile will be smaller than one in a river falling 5 feet in a mile; but at Quaker's Yard we have a well-formed large horseshoe bend in a river falling 50 feet in a mile. The laws which regulate the flow of rivers may be different in sand-rocks from what they are in alluvial soil; but I still hold the view as to the cause of these curves which I mentioned to the Society in 1852, but which it is unnecessary to discuss at the present moment*.

At A, Plate IV. fig. 1, the gravels are exhibited very clearly in the cuttings at and near Quaker's-Yard Station; these are 200 yards long and 40 feet deep, and are represented at the north end of the section. The local sand-rock furnishes the largest, most numerous, and least-rolled pieces, many of them 8 and 9 feet long. The boulders of Millstone-grit are generally well rolled, and not more than 6 feet long; those of Old Red Sandstone are throughout well rolled, and not often more than 2 feet. Not more than one stone out of 500 is derived from the Old Red Sandstone or Millstone-grit.

The gravel is supposed to be from 30 to 40 feet thick at A, covering the side of the hill to a height of 200 feet or more above the river.

* See Phil. Mag. 4th series, vol. v. p. 275, 1852.

The stratification is generally flat where the sand is bedded at all; but the junction of the sand-rock and gravel is rarely seen at any part of this section north of the viaduct.

The gravel of the Taff retains its character as far as the sea-coast, and resembles that of the Neath river, described by Mr. Moggridge, Quart. Journ. Geol. Soc. vol. xii. p. 169. During the excavations for the Swansea Docks, Mr. Moggridge found part of this river-gravel, which he identified by the presence of river-rolled Old-Red-Sandstone, Limestone, and Millstone-grit boulders (which had come from the parent-rocks, more than twenty miles distant) above stratified beds of marine clay alternating with peat. The marine clay contained *Scrobicularia piperata*, a shell which now lives abundantly in the sands of the adjoining shore.

The river Taff and the Swansea and Neath rivers expose sections of gravel all along their course to the sea-coast; and the ancient banks and river-beds of these rivers consist of well-rolled boulders from the higher districts, deposited with those of local rocks in every part. Their ancient banks have a contour and escarpments like those of the Aire (see Plate IV. figs. 4 & 5). The gravel becomes rather smaller as the sea-coast is reached, and has then, no doubt, been removed and redeposited many times. The gravel near the level of each river is of similar age throughout its course, and must be considered very recent.

Mr. Godwin-Austen (Quart. Journ. Geol. Soc. vol. vii. p. 134, pl. 7) also describes this part of South Wales under the heading "Insulated Areas in periods of greatest depression," from evidence of a very different character from that offered by me.

Fig. 7 is accompanied by a scale of heights in feet, and of length in chains (66 feet each), and is a reduction by the pentagraph of a section taken for me in 1867 to determine exactly the thickness and position of the Hirwain gravel. (See Map, fig. 3, page 67.) The thick deposits of gravel on Hirwain Common are cut through by several rapid streams flowing from the lofty range of hills separating Hirwain Common from the Rhondda Valley. I have selected the section, fig. 7, along the course of one of the Hirwain streams, as an instance of the continuity of gravel from high ground to the bottom of the valley, and of the regularity of the action of denudation and also of deposition.

Fig. 8 is a reduction from a drawing made of an excavation in the gravel and the underlying coal-seam. The gravel is tinted to represent the lower, middle, and upper gravels. The lowest portion consists principally of the clays and shales of the Lower Coal-measures, slightly moved from their outcrop down towards the river Cynon. The middle division is stronger clay than the upper; but as the gravels approach the river they become sandy, and have been worked over for the limestone boulders they contain, and for sand for the use of the furnaces at the ironworks.

This sketch gives an actual view of the passage of the soft beds of the Coal-measures into gravel, at the end of the gravel-period. This excavation gives a better illustration of the actual process of denu-

Fig. 7.—Section of Surface of Lower Carboniferous Series, Hirwan Common.

The author is desirous of drawing the attention of geologists to the contour of the surface in this section, as illustrative of his hypothesis of parabolic curves enunciated in his paper read on November 11, 1868 (*supra*, p. 7).—Edwr. Jan. 28, 1869.

dition than can be obtained from natural sections in brooks, where the different gravels are moved and mixed up together by the recent action of water. The total thickness of the gravel is 18 feet.

Fig. 8.—Section showing Coal-seams squeezed into drift during the deposition of the lowest gravel on Hirwain Common.—Natural scale.

BACON HOLE, GOWER, S. WALES.

Near this well-known cave limestone-gravel lies on a very steep cliff of limestone, in some places sloping as much as 70° to the sea. The top surface of this gravel lies at an angle of 34° ; and the bottom fills up the concavity in the limestone rocks. The path to the cave passes over this gravel, which is entirely composed of sharp angular fragments of limestone, cemented together by carbonate of lime, a process which seems to have been common in the Quaternary period. I noted, on a previous occasion, the manner in which limestone-blocks had been observed by Mr. Pengelly to be cemented together near Torquay, at a considerable height above the sea.

The mass of limestone-gravel falling down this cliff, accumulating in concavities, and sloping to the sea at an angle of 32° S. on the surface, may be seen at a height of 120 feet above the sea. In its course from the high cliffs above, it seems to have sent off a vein of gravel at a point 70 feet above the sea, to the east, at an angle of 11° .

This vein of angular gravel thickens to the east, and expands to a thickness of 25 feet when it reaches the opening in the limestone cliff called Bacon Hole, a deep fissure penetrating a long distance into the limestone cliff, in a northerly direction, and having a roof with an acute angle, which has often been described.

The gravel here includes some enormous blocks of limestone, which have fallen into it from the rock above. When this stream of gravel reaches the mouth of Bacon Hole, it turns sharp round into the cave in a northerly direction, the angle of deposition changing to 25° N., the floor of the cave being about 30 feet above high water. The limestone-gravel is covered with the black bone-earth in which so many remains of Elephant and Rhinoceros have been found in the cave; and is mixed with marine shells of species now living on the coast and in the sea-sand. The whole is covered with stalagmite, from 2 to 3 feet thick, also sloping northwards at an angle of 25° N. There are layers of stalagmite included within the limestone-breccia (proving changes in the conditions of deposition at intervals) in the cave; but the gravel, although so much higher on the side of the cliff to the west, does not fill up the whole of the cave.

Looking at Dr. Buckland's drawings of caves, I think there is evidence that much of the loose materials which have formed the cave-deposit have passed downwards from the subaerial gravels on higher levels, and continued through and over the caves as far as the level of the sea, lake, or river below them.

All English bone-caves are at levels within 150 feet of running water at the present time, or of the sea; and the great majority are within the limit of 70 or 80 feet, their fossiliferous contents corresponding in level with all the known fossiliferous gravels containing mammalian remains, including the Crag.

In the period referred to, water ran apparently in the bottom of the valleys which are now dry.

All bone-caves seem, at the time of the deposition of their contents, to have been in situations where water traversed them from fissures above, and flowed into them from the side streams flowing along the valleys in which they are situated, and they offer evidence of an immense rainfall at the period of the deposition of the cave-gravels. The very washed condition of the materials found in bone-caves, and the manner of accumulation, indicate that their contents were more derived from above than below; and the enormous pieces of limestone which have fallen into the gravel at Kent's Hole and Bacon Hole show that there must have been not only the action of running water thereon, but the force of a column of water upon the cave to dislodge the pieces from the roof.

There are in all caves apertures in the roof, evidently worn by water coming through it. I observed a good instance of this in the Paviland Cave at Gower (close as it is to the sea), and another in Brixham Cave. Some years since, I believe, it was remarked by Dr. Buckland that there were always at least three openings in bone-caves, two of them from higher levels than the mouth.

The subaerial gravel comes down the steep slopes of the limestone cliffs at Gower from a great height, and falls into a stratified marine deposit, just as it falls down the steep cliffs into the Hopes-Nose raised beach, near Torquay, where it is mingled with the sea-sand, pebbles, and shells deposited almost horizontally on the limestone rock.

TRANSVERSE SECTION OF RHONDDA VALLEY.

The Rhondda river unites with the Taff river twelve miles below Quaker's Yard, and fifteen miles from its source.

The transverse section across C measures 1730 yards from the opening of the levels on the opposite sides, 84 yards on the west side, and 79 yards on the east side, above the level of the Rhondda river, at the point C.

The gradient from the west level (D) to the river is 1 in 13, that from the east level (C) is still steeper, being 1 in $8\frac{1}{2}$; so that the sides of the valley slope at 1 in 11 on the average (from C and D to the river), or at an angle of 8° . The point C is about four miles from the furthest watershed of the river beyond.

Below the level of the quarry, on the west side, above the river, there is no rock seen except at one point in a watercourse.

The dip of the Coal-measures at this point (D E) is about 3° north, and the section (Pl. IV. fig. 3, D E) is nearly east and west.

The escarpment of the high ground above D and E is nearly twice as steep and high as that of the South Downs, and pursues its course nearly north and south along each side of the Rhondda valley. The dip varies, sometimes being northerly, and at other times south. This is marked by the arrows in the Ordnance map; and the lines of outcrop of the coal also show the direction of the escarpment. The escarpment, therefore, in the Rhondda valley is in the direction of the dip, and not of the strike; and the dip changes without any alteration in the direction of the escarpment. In fact, in this district, we have in a short distance escarpments parallel with the dip and strike, and also at almost every angle to them.

The valley is at D E about 300 yards deep, but is not shown above the quarry in the diagram, as I do not allude at the present time to anything more than the portion of the gravel in the valley below the No. 2 vein of coal. I visited this pit several times during the progress of the sinking, and counted the proportion of rolled to weathered stones each time. About one-eighth of the gravel in this pit was well rolled, the remainder being generally weathered.

It is evident that if only one-eighth of the gravel is rolled by the river, seven-eighths of it must have been obtained from the sides of the valley, west and east of the pit, and only one-eighth from the north, and that the valley was then widening much faster than it was lengthening in the latter part of the Quaternary period. There are times in the denudation of valleys when pluvial action on the sides of the valley is much greater than fluvial action in the centre of the valley. I mentioned (Quart. Journ. Geol. Soc. vol. xxiv. p. 117) that even in the valley of the Somme, where the fall of the river is so small, a great mass of the material deposited in the concavities of the valley near the margin of the river is derived from the adjacent country, and has never been river-borne. Different as is the position of the gravel of the Rhondda, in this respect it resembles that of the Somme.

The gravel-deposit continues for the whole length of, and is also met with at a considerable height above the Rhondda river. The

cause of this great fall of rock and clay is due to the position of the permeable and impermeable beds, and the influence of the immense rainfall occurring, I believe, at former periods, which has detached large portions from the high ground on the side of the valley, and let it fall into the channel of the river below.

The character of the deposit in this valley is very simple; it is sand-rock, clay, and shale, derived from the adjoining higher beds. In the pit sunk near C all the stones were either weathered or rolled, including one block of several tons weight. As the Rhondda valley is separated by a hill 800 feet high in one part, and 600 feet from Hirwain in the lowest part, above the river-course, which flows off from the Millstone-grit and Old Red Sandstone, there are no boulders of these rocks in this valley; all is purely local.

In sinking the pit at C, which was 20 feet in diameter, the rock was reached on the west side of the pit 8 yards before solid ground on the east side was met with, so that the angle of the escarpment of the ancient bank of the river or side of the valley was about 50° to the ancient river.

By means of this sinking it is known that the gravel there is 101 feet thick; and, taking the average thickness to be 20 yards, and the width 1700 yards, there would be 34,000 cubic yards in each yard run (supposing that section C represents the average amount of gravel in the valley), which, at $1\frac{1}{2}$ ton to the cubic yard, represents 89,760,000 tons of gravel to the mile. The river flows over at least 50 feet of gravel, and probably much more.

The section of gravel passed through in the pit is as follows:—

	ft.	in.
Turf or peat	3	6
Blue clay with large stones	23	6
Sand	18	0
Blue clay.....	5	6
Gravel with large stones	26	6
Gravel with clay	13	0
Cliff.....	5	0
Gravel.....	6	0
	101	0

The fall of the Rhondda river in the distance of a mile before it reaches C is about 500 feet, in which distance there are two waterfalls.

The vast extent and the character of the Rhondda gravels proves that the denudation of the Rhondda valley was not dependent upon large rivers, or marine or glacial action, but might have been caused by intense pluvial action.

In a wet season, water pours over the escarpment along the edge of the hills into the Rhondda valley, and is not confined to a few distant watercourses; but there is no evidence of any accumulations of gravel now; perhaps the fall of stones along the escarpment equals the quantity carried down the river in floods.

The Salisbury pits are celebrated for their mammalian remains,

which, according to Dr. Blackmore, are associated with flint instruments of the Amiens type.

Fig. 9.—Section near Salisbury.

Land and freshwater shells are very abundant. The great shell-bed is near the chalk, lying upon a bed of gravel; and the Pupa-bed touches the lower surface of the covering bed of gravel, *e*. While the curve of the surface is on the average 3° between A and B, the dip of the lower gravel, *b*, where it touches the escarpment of the chalk, will probably lie at a very steep angle.

The Pupa-bed, *c*, a fine loess, is marked by a distinct band of colour; it is about 2 ft. thick, and slopes to the river in a gentle curve. I have marked its dip 16° , then 11° , then 9° . The bed *c* contains only six species of shells, *Helix arbustorum* (?), *Helix hispida*, *Zua lubrica*, *Lymnaea palustris*, *Valvata piscinalis*, *Pisidium obtusale*, besides the *Pupa* in immense abundance. The lower band *b* contains thirty-one species of terrestrial and fluviatile shells, according to the lists of the late Mr. John Brown and Dr. Blackmore.

The chalk, *a*, is 30 feet from the surface at A, but in 30 yards it reaches the surface; so that the escarpment in the part not yet opened is known to be at an angle of 30° , sloping towards the river. As is usual, the brick-earth is thickest a little way from the escarpment, and passes, on the same horizon, into gravel and sand as you approach the river.

The remains of another arctic mammal (the Musk-Ox) have been recently found in the loess at Salisbury. This animal is now living in a remote part of North America. The stratification at Salisbury, shown in fig. 9, is very similar to

that at Erith, fig. 17.

The mammalian gravel and Combe rock extends from Brighton, without interruption, to the point drawn, fig. 10, and presents a cliff of roughly stratified beds facing the sea, about 60 ft. in height.

The escarpment of the chalk is known to be not far distant in the rear of the present face of the cliffs at any point; and an extensive fall of gravel has exposed the escarpment of chalk. In the steepest

part it is 60° S., curving gradually towards high-water mark. At *b* the raised beach of flint-shingle, with masses of slightly rolled chalk, is seen reposing on it, dipping to the sea at about 6° , and above 8 or 10 ft. above the present high-water mark.

There are alternations of beds of very large flints beach-rolled, and layers of chalk débris, many pieces with the corners slightly rounded, varying from 6 in. to 2 feet in length. The fall of cliff at the spot represented in fig. 10 enables the geologist to measure the

Fig. 10.—Section about half a mile East of Sussex Square, Brighton.

angle of deposition of the gravel and Combe rock north and south, and also to observe the face east and west. The dips at several points have been marked in the sketch. The raised beach dips south at from

Fig. 11.—Section at Brighton.

6° to 8° , and the covering bed also south at 2° . We thus meet near the sea with the condition of the basement bed of gravel (*b*, fig. 10) and of the covering bed of gravel being deposited nearly horizontally, while

the intermediate series of stratified gravels, with the lower beds called Combe rock, a mixture of chalk and flint with yellow clay, have been pitched and washed by pluvial action over the escarpment. These materials, derived from the high ground behind the escarpment, and thrown over it, have not been sufficiently water-washed, nor perhaps even placed within the reach of water flowing sufficiently fast, to remove the angles of the soft chalk fragments; nor has the act of deposition abraded the edges of the flint, which are often splintered before being bedded in the Combe rock.

Fig. 11 is another view of the structure of the Brighton gravel-bed, at a point a little to the west of fig. 10, where the escarpment of the chalk must be very near the face of the cliff. The raised shingle-beach is 6 ft. thick, and falls at 8° south to the sea. The position of the present beach is shown 9 ft. below the ancient one. Dr. Mantell first obtained remains of *Elephas*, from these deposits, near Sussex Square; but no shells have yet been found in them.

Fig. 12 has been, by mistake, reversed in the drawing, and is

Fig. 12.—*Vertical Section of Brighton Cliff.*

a vertical section of the Brighton gravel-beds at a point intermediate between that shown in fig. 10 and Sussex Square. By means of a footpath up the cliff, I was enabled to determine the dip of part of the middle series of gravels (*d*) to be 25° west. The covering bed and gravel (*e*) and the raised beach deposited at the bottom (*b*) appear quite horizontal at that point.

The old beach (*b*) is about 7 ft. thick, and the chalk boulder-bands (*c*) interstratified with flints about 9 ft. thick, the cliff being about 63 ft. high.

Fig. 13, on a larger scale, exhibits a front view of the cliff 100 yards east of the Rottingdean landing-place, three miles east of Sussex Square, Brighton. This is near the termination of the Brighton gravel-deposit; and I have drawn it to show the manner in which the gravel reposes in a hollow of the chalk, formed by a stream which reached the sea at Rottingdean in the gravel-period.

There is a bed of marl on the 6 feet of solid chalk, shown in fig. 13,

Fig. 13.—*Section of the Cliff East of the Gap, Rottingdean, Sussex.*

and also continuous bands of flints. We can thus accurately observe the marks of the movement which has taken place in the upper surface of the denuded chalk. The bed of tabular flints, about 1 inch thick, touching the upper part of the artificial opening marked *cave*, is fractured and contorted, while the traces of a still higher band of large detached flints are visible in the gravel above the top of the cave, where it ends. Then comes a coarse ochreous flint-gravel 7 feet thick, with veins of sands and some chalk fragments. This is eaten into by the covering dark bed of sand with flints. The middle portion of this section is chalky, and must have been more altered by chemical than by mechanical action, as the flint-veins have been so little disturbed. At p. 113 of the last volume of this journal I gave some illustrations of the destruction of chalk *in situ* by some kind of chemical action, which was probably contemporaneous, if not identical, with that which has eroded the surface of the chalk, and formed pipes over such large areas.

Fig. 14, representing the highly-inclined chalk with flint veins 200 yards east of Freshwater Gate, Isle of Wight, is given for comparison. The covering bed is nearly horizontal, as usual. The character of the chalk and the disposition of the gravel closely resemble those at Rottingdean; and the latter is about the same height, from 25 to 30 feet. Mammalian remains have been found in this bed close by Freshwater Gate.

Fig. 15, Sangatte Cliff, near Calais, is given for comparison with these Brighton sections. The escarpment of the chalk, near the upper part is 43° , then it reaches 55° , and descends to 43° . The Sangatte raised beach is just above the level of high water at this point, and is composed of large flints and pieces of chalk, as at Brighton.

In the middle of the Sangatte series a line of chalk boulders, deposited at an angle of 12° , is shown in some places. At about half a

Fig. 14.—Section of Cliff East of Freshwater Gate, Isle of Wight.

Fig. 15.—Section of Sangatte Cliff from the beach.

mile from this point Mr. Prestwich found, just above high-water mark, abundance of freshwater shells of living species in a fine tranquilly deposited marl covered by 50 feet of gravel, continuous with the upper 30 ft. of the gravel shown in fig. 15. The slope of the covering bed of gravel (*e*) is not more than 1° or 2° . There is very high ground at the back of this section (which is near Cape Blanc-nez), from which the chalk and gravel were derived and thrown over the ancient chalk-cliff 80 feet high.

Near the village of Sangatte the covering gravel reaches high-water mark, and the whole of the middle series intercalated between the basement gravel and the covering gravel is seen.

While the materials of the gravel at Sangatte are principally brought over the cliff from the high land, they are interstratified with an ordinary beach and with a lacustrine deposit formed at the same time within a few feet of the present level of high water.

A similar beach, or a similar lacustrine deposit, might be formed

at the present time; but the physical conditions under which a gravel-deposit like that at Sangatte could occur do not exist at the present time in France. It would require the rainfall of a pluvial period to furnish the fresh water to lay down such a thickness of gravel as that of Sangatte in such a position on the sea-coast.

THAMES-VALLEY GRAVELS.

The section from the surface of the Thanet Sands, 86 feet above the Ordnance datum-line, to the river Thames, through the celebrated Cyrena-pits (Pl. IV. fig. 6), is drawn to an exaggerated scale of heights, in proportion to lengths, through the points A E I, approximately laid down on the map, Pl. IX. Pl. VII. is a copy of part of the Ordnance Parish Map of Grays reduced to half size.

An excavation into the chalk between B and C offers an opportunity of observing the Thanet Sands quite undisturbed at a distance from the surface of 27 ft. at C, and of 35 ft. at B; the dip of the chalk is there slightly south, the thin covering of gravel *e* dipping south. There is no opportunity of examining the structure of the ground between C and D, as it is not excavated; but I expect the section resembles that at Crayford, on the opposite side of the river.

Fig. 16 is a transverse section near the point D, showing about

Fig. 16.—Section at Grays, Essex.

18 ft. of yellow false-bedded sands, *d*, under the covering gravel, *e*, and lying upon the 4-feet pebble-bed, composed principally of pebbles from the Woolwich beds of the Eocene series. The *Cyrena* is abundant in the sand of this pebble-bed, and occurs, associated with other shells, in the shell-beds below at *e*, which are from 1 to 4 ft. in thickness, varying much in character, but dipping north to the

escarpment C. The beds below *c* are stratified clays, sometimes mottled, and are said to repose on the lower gravel. This lower gravel is said to be 3 ft. thick; and at 45 ft. from the surface the chalk is said to be found near D; but I have not seen this myself.

Some years since, an excavation was made near M, and chalk was found about 43 ft. from the surface. The false-bedded sands were 20 ft. thick at that point, and covered the upper-gravel series, *e*, very full of flints and pebbles*.

The Mammalian remains occurring in the gravel, *f*, are more rolled than those found in the brick-earth above.

A list of Grays species has been published by Prof. Morris, and also in the Geological Society's Journal, vol. xxii. p. 101, by Mr. Boyd Dawkins.

Returning to Pl. IV. fig. 6, the false-bedded sands are represented thinning out towards the river, as well as the Cyrena-beds. The depth of the chalk below the surface is ascertained by a well-section at F; and the covering gravel, *e*, comes in contact with the lower gravel, *f*, near the railway-cutting, *g*. Close to this point there was a section of coarse gravel, 12 ft. thick, open in a gravel-pit, reposing upon chalk without any beds of sand or brick-earth. The chalk is said to have been seen in the railway-cutting 9 ft. above the rails at the bridge I. In the tramway east of, and close to, Grays Station, the chalk is also well exposed 6 feet above the rails, with thick beds of gravel lying irregularly upon it, and some brick-earth beds and gravel above them containing comminuted shells derived from the Woolwich series, which beds have been entirely removed by denudation at Grays. In excavating for a cellar near Grays church, a very large mass of sandstone was found about 20 ft. above the Ordnance datum-line, many tons in weight, in the gravel close to the chalk. Such stones are abundant in the upper gravel in the Chalk and Thanet Sand west of the Grays Station.

Great changes have taken place at Grays, by removal of brick-earth, since Professor Morris first described the fossils from the Thames-valley deposits; and he is almost the only geologist who has watched the excavation from year to year. It will be seen from Pl. IV. fig. 6, that, like the gravels of the Somme, the fossiliferous brick-earths of Grays are deposited in a concavity of the chalk. The chalk is 15 ft. lower at D than at G, and 43 ft. lower at D than B. The old Thames river seems to have divided into two channels at Grays; and the fluvial deposits between C and G appear to have occupied one channel, thus forming the whole series of stratified beds which are intercalated between the gravels *e* and *f*.

The last deposit was the covering gravel-bed, *e*, which is continuous from the marsh to the point A, and beyond it.

The position of the section (Pl. IV. fig. 7, Erith) A B S T is laid down on the sketch map, Plate IX.; and A B, the part near the escarpment of the Chalk and Thanet Sands, is enlarged in fig. 17. page 85. Although Erith and Grays are not exactly opposite each other on the Thames valley, yet the sections, Plate IV. figs. 6 and 7,

* I obtained this information from Professor J. Morris, F.G.S.

will give an idea of the surface-configuration of the Thames valley and of the position of the fossiliferous gravels. The top bed of the Chalk and the basement bed of the Thanet Sands are well seen at G, Plate VIII. (fig. 18, p. 86), at a height of 46 feet, at Erith; and at B, Plate IV. fig. 6, Grays, at a height of about 47 feet above the Ordnance datum-line.

This basement bed is 70 feet at Purfleet, and 66 feet at Crayford (fig. 22) above the Ordnance datum-line.

The supposition of Mr. Searles Wood, jun., that the Thames valley is in a line of fault appears to me inadmissible, as far as Grays, Erith, and Crayford are concerned. The sections I now bring forward will show that the Thames valley has been excavated by river-action out of a mass of chalk and Thanet sands, lying nearly horizontally in the localities referred to. Beyond the districts referred to there is a flexure in the chalk which depresses the basement-bed of the Thanet sands from 46 feet at G, Plate VIII., Erith, to about 10 feet above the Ordnance datum-line at Erith station, without any fault. It is seen again at Woolwich and Lewisham, at the same height as at Erith, before it sinks under the London basin.

Referring to Plate IV. fig. 7. The chalk has been seen at A, 50 feet from the surface and 36 feet above the Ordnance datum-line, also in a well at S, 13 feet the above datum-line. It is said to be very near the surface at S, and that it is visible at low water for a mile between Erith and Crayford Ness, T. Although the ground along B S has not been excavated, like the other side of the Thames at Grays, it is known that the brick-earth thins out, and that the lower gravel (*f*), separated from the covering gravel (*e*) by brick-earth (*d*) and sands (*c*), together 50 feet thick between A and B, come together at S, just as *e* and *f* touch each other in fig. 6, Plate IV., as they approach the centre of the valley.

The escarpment of chalk and Thanet sands has not yet been excavated

Fig. 17.—Section of the Cyrena-pit, Erith.

west of A; but I have no doubt that the fossiliferous gravels, A B, will be found to lie in a similar concavity to those at Grays and Crayford, where the Thanet sands and chalk have been exposed.

The material for the beds *c* and *d* seems to have descended from A, and to have been brought by land-floods, and contains bones, but no shells except derived ones. On the contrary, *b* is full of shells deposited quietly. Many of the species are extremely delicate, and are found quite as perfect as recent specimens; the two series of deposits are interstratified together at D, the eastern bed (*b*) being of fluvial origin, and the western (*d*) being apparently of pluvial origin, and more due to land-floods.

The chalk below A in fig. 19, Erith, along the line A B, Plate VI., is 50 feet from the surface. The bottom-bed (*b*) shown in the section is false-bedded sand with a few veins of gravel. The *Cyrena* is very abundant between D and B in this sand (*b*). The mammalian remains are most abundant in the brick-earth beds (*d*).

These beds (*d, d*) dip near A at 25° E, then fall to 9°, 5°, and eventually dip 5° W. near B, forming a gentle curve. The covering bed (*e*) is 4 feet thick, and full of the Woolwich pebbles between B and D. It then changes its character, and contains only a few flints, but masses of the Woolwich shell-bed when it reaches A. The remains of this shell-bed occur through *b, c, d*, and *e*. The angles of deposition were observed both by Mr. S. Skertchly and myself in 1868, and the heights were measured by the former. Great removals of brick-earth have recently been made near A, so that the original section cannot be seen.

Fig. 18.—Section in Erith Pit. (Natural scale.)

Between B and C, Plate VIII., a long face of the *Cyrena*-sand-bed

is exposed at 45 feet above the Ordnance datum-line. I have found the *Cyrena* at about the same height at D; and this is the highest level at which the *Cyrena* is found either in the Thames valley or in the Humber or Somme.

Fig. 18 is a section exhibiting the excavation of the chalk and Thanet sands by the old River Thames. The chalk (*a*) reaches a height (fig. 18) of about 45 feet above the Ordnance datum-line; and the Thanet sands (*b*) have been partly denuded, so that only 15 feet, out of 60 feet seen in the Station ballast-pit at Erith, remain here. The gravel (*c*) reposes conformably on the top of the Thanet sands at G, the upper part of the section; but the gravel follows the denuded surface of the chalk, filling up the concavities. It contains much rearranged Thanet sand; and a large lump (*b*) is conspicuous in it. The 6-feet step in the chalk is well marked and sharply cut out by the action of the river flowing at right angles to the line G K.

Fig. 19 is a section in the same direction as fig. 18; but steps from

Fig. 19.—Section in Erith Pit. (Natural scale.)

1 to 2 feet deep, cut by water in the Thanet sands, are distinctly seen. The gravel is then deposited in the concavities formed by the water. Brick-earth, with veins of gravel (*d*), follows on for 15 feet, and the covering gravel (*e*) succeeds, sloping to the river at a gradient of 1 in 200 only. The Thanet sand is a somewhat incoherent mass; and the action of the water must have been very gentle to have formed such perfect steps. The deposition of gravel and brick-earth must have at once followed the denudation.

Fig. 20 is a section along part of the line G E. The chalk, with a basement-bed of Thanet sands and green-coated flints, is seen at *d*, *in situ*. The Thanet sands, in fig. 20, were excavated by one of the side-streams which formerly flowed into the ancient Thames. The edges of the sand (*b*) are so sharply cut in this transverse section of a river-bed that it would be difficult to believe it was not an ar-

tificial excavation, if other sections, such as fig. 19, did not exist close by. The gravel (c) fills up the inequalities in the sand.

Fig. 20.—Section in Erith Pit.

Rearranged Thanet sand is frequently met with in these gravels, which were formed out of materials derived from this bed during the excavation or deepening of the Thames valley and its tributaries at this period. The large bull's-head flints from the basement-bed are carried down to lower levels (owing to their weight) and deposited on the new surface of the chalk.

The pebbles from the Woolwich bed are most abundant in the middle part of the Thames Quaternary series, and have been carried a great distance from and below the escarpment before they were redeposited in the gravel.

Fig. 21 is from a drawing on wood by Mr. S. Skertchly of a mass

Fig. 21.—*Erith Pit*.

of Thanet sands (b) 38 feet in length by 7 feet 10 inches in width, with a portion of the Woolwich pebble-bed attached to it, lying upon and against one piece of the Woolwich shell-bed 12 feet by 6 feet

5 inches (*c*), and near another fallen piece,—the whole enveloped in a mass of purple clay (*d*), which may be either Plastic or London clay rearranged, and which often contains pieces of rolled chalk, like what is called Till or Boulder-clay. The situation is near M, Plate VIII., Erith; and the height of *b*, fig. 21, is about 65 feet above the Ordnance datum-line. Large pieces of the Woolwich shell-bed, with masses of purple clay, occur in the ordinary covering bed (*e*) near I, at a height of 55 feet. This covering-bed loses its pebbles at J, and at L M N assumes the character of a purple clay enclosing large unrolled fragments of the Eocene beds and small pieces of chalk. The Woolwich shell-bed, with the pebble-bed, is visible *in situ* at the south-west corner of the ballast-pit at Erith Station, at a height of 60 feet above the chalk, the whole series of Thanet sands being exposed there below it. It is probable that the escarpment of Thanet sands and chalk is only a little to the west of M. It is evident that this particular mass (*b*, fig. 21), must have been, when *in situ*, at least 110 feet above the Ordnance datum-line; for part of the pebble-bed is still attached to it, and that pebble-bed is 60 feet above the basement-bed of the Thanet sands, which sands would be, if *in situ*, 50 feet above the datum-line at the point M, according to my calculation. The positions of the top and bottom of the Thanet sands can be determined, as they are visible *in situ* between J and I, and also along the face of the chalk-pit between G and E.

The difficulty is, to understand what kind of wash of water removed this incoherent mass of Thanet sands and deposited it again (without breaking it up into sand) 35 feet lower down, upon a mass of the Woolwich shell-bed (*c*), which must have descended 45 feet and changed places with *b*.

As the Thanet sand has been removed from the large area of chalk to the west, the mass of sand (fig. 21) may have been derived from a distance.

The Thanet sands before their denudation dipped towards the north, from the Kentish hills to Erith, at a slope of 1 in 100 on an average of ten miles. It may have been owing to the existence of a rainy period, and to high land to the west, that water could have moved the Thanet sands. Eastward, towards the low ground near the Thames, on the opposite side of the river, at Grays, I have only seen one small mass of Thanet sand imbedded in the covering gravel (*e*) near F, Plate VII.; but to the east of Grays the chalk soon dips eastward*, and therefore falls below the level of the river; so that there was no high land to give transporting-power and velocity to the water that fell upon it. The nearest English deposit in physical character, and perhaps in age, to that represented in fig. 21, is at Cromer. It is also probable that without the high background near Cromer such a deposit as that in the Cromer cliffs could not have been formed. The great mass of chalk at Trimmingham is an instance of an unbroken rock having been detached and moved down to a lower level in the Quaternary period, and enveloped in clay of a different formation. The stratified fossiliferous gravel on the Norfolk coast has been tranquilly deposited in front of, or near, the great slipped masses

* And northward.

of what is there called boulder-clay, just as the stratified fossiliferous gravels at Erith are close to and interstratified with the non-remanié but moved Tertiaries at Erith.

Fig. 22.—Section in Crayford Pit.

This section is to natural scale, but is reversed.

The basement-bed of the Thanet sands and a few feet of the sands are exposed along the face of the cutting, A C, Plate VI. The surface of the chalk is seen to be eroded by chemical action into pipes and basin-shaped cavities. The Thanet sands are broken through and moved into the pipes in the usual manner, and are covered by a top bed of sand and gravel, which fills up all the hollows of the underlying sands and chalk, and smooths over all previous inequalities.

The beak of chalk under *a* has protected small masses of Thanet sands (B and B') below and behind it from being remanié and mixed up with the gravel. The covering bed of sand and gravel is smooth and nearly horizontal. *c* dips at 10° towards the Thames, and in one point at 12°. The chalk near A is 70 feet above the Ordnance datum-line, although only 50 feet is shown here. At P, Plate VI., 400 feet to the south-east, the chalk is seen in a pond only 10 feet above the Ordnance datum-line.

The inclination of the escarpment between C and P is 1 in 6, or 15° on the average; and this is entirely due to the excavation of the channel by the ancient River Thames. The lowest bed exposed in fig. 22 consists of from 3 to 4 feet of coarse gravel lying on the eroded chalk.

Fig. 23 represents rather higher ground; and 15 feet of Thanet sands are seen *in situ* (*b*). The gravel and brick-earth are well seen, dipping as much as 22° east at the steepest point.

The covering bed of sand and gravel, 5 feet thick (*a*), lies on the edges of the gravel (*c*) and brick-earths, and slopes to the river at a

gradient of 1 in 25 in that part. Only 48 feet in height is shown in fig. 23. The sketch is to natural scale; so that the angles of the escarpment of chalk are not exaggerated.

Fig. 23.—Section in Crayford Pit.

E. D 80.

88 C. W.

Fig. 24 is a view near the curved line E F, of the eroded surface of the chalk (A), with rearranged Thanet sands (B) 4 or 5 feet thick,

Fig. 24.—Section in Crayford Pit.

enclosing large flints from the basement-bed. The deposition of 6 feet of coarse flint-gravel (C), on the bed B, in a hollow of the chalk excavated prior to the gravel-period, is well exposed just where the tramway cuts through it.

The brick-earths above the gravel have been excavated. The brick-earth generally is more valuable than the sand and gravel, and the workmen often leave the lower beds in the state shown in fig. 24

for a long time. The chalk is excavated at the tramway to 20 feet above the Ordnance datum-line, and is 55 feet high at E, where the Thanet sands (B) are nearly *in situ*.

The curve in the line of section gives an appearance of flexure to the veins of flints in figs. 24 and 25 which does not really exist. These bands are not far from horizontal.

Fig. 25.—Section in Crayford Pit.

Fig. 25 is along the line T U, Plate VI., and shows the eroded surface of the chalk in an oblique direction to the old river-channel. The gravel (B) is very conspicuous, and the brick-earth series (D) is thinner than usual; but this is due to the distance of U from the escarpment. To the east of U the brick-earths gradually thin out, and the upper and lower gravels unite, forming a bed of coarse gravel 8 or 10 feet thick lying on the chalk. The height of the point T is about 40 feet, and the chalk under U about 20 feet, above the Ordnance datum-line.

Fig. 26, a section along the line I J, offered good opportunities

Fig. 26.—Section in Crayford Pit.

for examining the dip of the gravel, and showing the relation of the thick brick-earth beds deposited at angles as high as 19° , 17° , 14° , 11° , and 9° , in curves falling to the river. The beak of chalk is shown at about 18 feet above the Ordnance datum-line. It is about 10 feet long; and the junction with the escarpment of chalk is not seen.

This drawing is by Mr. S. Skertchly, to natural scale.

Fig. 27 is from a drawing I made in 1866, near the line I J.

Fig. 27.—Section in Crayford Pit.

The mass of Thanet sand B, below the beak of chalk near A, was then well seen; patches of these sands also occur in the brick-earths and gravels C. The dip was 15° where marked, and that of the covering bed only 2° . Fig. 27 is to natural scale, and represents a greater length of section than fig. 26. This part of the pit has recently

Fig. 28.—Section of part of the Cyrena-bed, Crayford.

been excavated; so I give both sections for reference. The coarse gravel in front of the beak is pushed out horizontally into the finer beds of sand, and divides into fine points. There were other beaks of chalk visible at G and H.

Fig. 28 is a careful sketch of part of the Cyrena-bed, by Mr. S. Skertchly. The section is as follows, in ascending order:—sands 4 feet, wavy laminated brick-earth beds 6 feet, brick-earth 5 feet, a bed of pebbles from the Woolwich series, brick-earth, and clay 6 feet; the Cyrena sand-bed, C, passing upwards into laminated brick-earth, gravel, and loam, 25 feet thick.

Fig. 29 is a section of the most fossiliferous part of the Crayford pit,

along the line K L (see Pl. VI.). It represents about 30 feet in height by 150 in length; so that the horizontal scale is about five times the vertical. It is intended to show the variation in the pebble-beds, C, and the thickening of the whole series in the direction of the escarpment. The pebble-beds, C, are in three divisions, and the great Cyrena-bed, D, rests on one of them; but shells are found in B, C, D, and E, though not in the covering bed F, which contains flint-pebbles distributed through it in the usual manner. The lowest beds are not seen in fig. 29; but at X, near M N, there are several openings at a depth of 35 feet from the surface, where a bed of coarse gravel is visible, which is near the eroded surface of the chalk. The chalk itself, with a covering of gravel or sand, is seen in the ponds marked P in the Crayford pit, where the water stands a little below the level of high water.

The Cyrena-bed varies from 4 feet to 6 feet (D, fig. 29) in thickness, and consists of false-bedded sand containing a great number of small and delicate shells in a perfect state of preservation.

Fig. 30 exhibits the series of gravels in the Stoke-Newington pit,

which is 43 feet deep. The bottom consists of yellow false-bedded sands, passing upwards into a series of stratified brick-earths and clays with veins of gravel. The covering bed, *k*, is indented into the brick-earth, *j*; it is from 6 to 8 feet thick, composed of large flints in stiff brown clay, and forms a marked contrast to the finely laminated loams and clays forming the 30 feet below it, *c-k*. The bed *f* is a black peaty clay; and the shells in the list from Stoke Newington are found in brick-earth and clay above this black bed in the adjoining pit. The covering bed reaches a height of 125 feet above the Ordnance datum-line, and is within 300 yards of the escarpment of the London Clay to the north. Where the London Clay is visible to the north,

Fig. 31.—Section at Grays, illustrating the rearrangement of the Thanet Sands in Pipes in the Chalk near L, Plate IX.

at a height of 140 feet, it is only covered by 6 feet of gravel, while in fig. 30 it is covered by about 50 feet of the Quaternary series.

Fig. 31 is a drawing to natural scale, showing the relations of the Thanet sands to the covering bed of gravel, where these sands have fallen into pipes in the chalk. The Thanet sands have not been faulted by any general movement, but have dropped down where they have been undermined by the removal of chalk by chemical action.

It would appear as if this chemical action had been contemporaneous with the denudation of the Thames-valley district by pluvial and fluvial agency, and had preceded the deposition of the covering bed of sand and gravel. The proof of this is that the fissures in the sand are filled by veins of gravel from $\frac{1}{2}$ inch to $1\frac{1}{2}$ inch wide to a great depth.

When the sands fell by gravitation into the great pipe (A), (which penetrates the chalk 25 feet, and the Thanet sands an equal amount), they filled it up to a height of 20 feet. This must have left a large funnel-shaped hole on the surface of the sands. It will be seen that, although the gravel is only 5 or 6 feet thick over the undisturbed sands at B, it has been able to fill a cavity 30 feet deep in the Thanet sands and chalk; and yet the top of the gravel is so smooth that no one could

tell from the appearance of the surface that such a cavity had existed. There must have been a wash of gravel all over the sand to have furnished the material to fill up such cavities as that shown in fig. 31; and since that wash of gravel there is no indication of any subsequent deepening of the pipes in the chalk.

The Thanet sands (*b*), 25 feet thick at B, at the north of fig. 31, are seen lying *in situ* on the chalk.

Where the formation of the great pipe A has undermined the Thanet sands, *b*, the latter have fallen down to the bottom of the pipe A, which is funnel-shaped, and filled up 20 feet without any mixture of gravel whatever. The sands moved from their proper position are indicated by the letters *b' b'*; the bottom part of the pipe A is not shown in this drawing. Under C, masses of sand, *b b*, are shown falling into the pit by gravitation, from the removal of the chalk by chemical action. The gravel series *c c* is 30 feet thick above A, and is similar in mineral character to the covering bed *e*.

Mr. Prestwich has explained the formation of pipes by chemical agency in the chalk.

Fig. 32 is a drawing of other beak-like projections in the chalk

Fig. 32.—Section near Knighton Mill, Isle of Wight.

where it has been acted upon by denudation previously to the deposition of the gravel near Knighton Mill, Isle of Wight.

Fig. 33 (p. 98) is a section of a fissure or pipe in the highly inclined chalk on Brading Down, Isle of Wight, which is a remarkable contrast to the ordinary pipes which are filled with sand or gravel.

The general conclusions to be drawn from the sections of the Thames-valley fossiliferous deposits appears to be that they are a series of variable sands, loams, and clays intercalated between the upper and lower gravels.

The false-bedded sands are below the loams and clays at Erith, and above them at Grays, while at Crayford they are in the middle of the series. This may simply arise from the principal current changing from one side of the old river to the other during the deposition of the Thames Quaternary series. By this means, where the current became more rapid, sands would be deposited, and where

the current was reduced, the finer materials might be deposited. There is no evidence of any change of level during these deposits; and the river must have risen 50 feet at least in times of flood.

Fig. 33.—Section on Brading Down, Isle of Wight, showing blocks of Chalk in Gravel.

In conclusion I may mention that the beds described above as pluvial have been referred to by Mr. Godwin-Austen under the name of Head, or Subaerial beds; and I quite admit the propriety and importance of his terms in many respects; but I was obliged to use the term "pluvial," in order specially to introduce the notion of rain-water action, as I was treating of deposits made on slopes by the action of water (probably by a non-continuous action) and thus necessarily to be separated from the ordinary aqueous action, such as is performed by rivers, lakes, and seas; as well as from ordinary subaerial action. While the pluvial beds, as I have shown in my sections, pass down into the aqueous beds, they pass upward into Mr. Godwin-Austen's subaerial beds, and the stones which have been *head* at one time, and weathered by subaerial action, pass down into what may be termed pluvial beds, which in their turn are being constantly despoiled, and carried into the purely aqueous beds at lower levels. Although Mr. Godwin-Austen's term subaerial is an excellent one, I submit that it does not sufficiently define the kind of action for the purpose I have in view, and still less so for the paper I have prepared on the formation of valleys, and the denudation of the surface.

When the Crayford and Erith beds have been as fully examined as those at Grays, the numerical superiority of the Mollusca in the latter deposit (as shown in the following Table) will probably disappear.

The list of shells from Salisbury is principally on Dr. Blackmore's authority. The Grays list is compiled from Mr. Pickering's table, the Ilford one partly from Mr. Morris's; the remainder by Mr. A. Bell, who has revised the list, and omitted the doubtful species.

N^o 1 Section - Horizontal Scale same as Vertical between A and E; between E and G the Horizontal Scale is one fourth the Vertical.

The Heights have been obtained by Survey.

Cottage Well North End Crossing
S. E. Rail.

MAP

To illustrate Memoir by A. Tylor F.G.S.

PART OF KENT AND ESSEX.

Key showing position of Plans, Plates VI, VII & VIII - and the River Thames.

The list from the Stoke-Newington fossiliferous pit, discovered by me in 1868, has been prepared by Mr. J. Wood Mason, F.G.S., who has added several species to the list first published.

The Land and Freshwater Mollusca of the Fossiliferous Gravels at localities referred to in this Memoir.

Species.	Grays.	Tlford.	Erith.	Crayford.	Stoke New- ington.	Salisbury.
<i>Achatina acicula</i> , Müll.	*?					
<i>Ancylus fluviatilis</i> , Müll.	*	*	*	*	*	*
<i>Anodonta cygnea</i> , Linn.	*	*	*	*		
<i>Balæa fragilis</i> , Drap.	*?					
<i>Bithynia tentaculata</i> , Linn.	*	*	*	*	*	*
<i>Carychium minimum</i> , Müll.	*	*	*	...	*	*
<i>Clausilia biplicata</i> , Mont.	*	...	...	...	*	
<i>Cyclas cornea</i> , Linn.	*	*	*	*	*	
<i>Cyrena fluminalis</i> , Müll.	*	*	*	*		
<i>Helix arbustorum</i> , Linn.	*	...	...	...	*	*
— <i>concinna</i> , Jeff.	*	...	...	...	...	*
— <i>fulva</i> , Müll.	...	*	...	...	...	*
— <i>hispida</i> , Linn.	*	*	*	*	...	*
— <i>nemoralis</i> , Linn.	*	*	*	*	*	*
— <i>pulchella</i> , Müll.	*	...	...	*	*	*
— <i>pygmea</i> , Drap.	*	...	...	...	...	*
— <i>rotundata</i> , Müll.	*					
— <i>runderata</i> , Stud.	*					
— <i>rufescens</i> , Pen. (Morris) ...	*	...	...	...	*	*
— (Zonites) <i>crystallina</i> , Müll.	...	...	...	*?		
— (—) <i>nitidula</i> , Drap.	*	...	...	*		
— (—) <i>radiatula</i> , Ald.	*	...	...	*		
<i>Hydrobia marginata</i> , Mich.	*	...	*	*		
<i>Limax agrestis</i> , Linn.	*	...	...	...	...	*
— <i>marginatus</i> , Müll.	*					
<i>Lymnæus auricularius</i> , Linn. ...	*	*	...	*		
— <i>palustris</i> , Müll.	*	*	...	*	*	*
— <i>pereger</i> , Müll.	*	*	*	*	...	*
— <i>stagnalis</i> , Linn.	*	...	...	*		
— <i>truncatulus</i> , Müll.	*	*	...	*	*	*
<i>Paludina</i>	...	...	...	*		
<i>Pisidium obtusale</i>	...	...	...	...	...	*
— <i>amnicum</i> , Müll.	*	*	*	*	*	
— <i>fontinale</i> , Drap.	*	*	*	*		
— <i>pusillum</i> , Gmel.	*	...	...	...	...	*
<i>Planorbis albus</i> , Müll.	*					
— <i>carinatus</i> , Müll.	*	...	*	...	*	*
— <i>complanatus</i> , Flem.	*	...	...	*	*	
— <i>contortus</i> , Müll.	*	...	...	...	*	
— <i>corneus</i> , Linn.	*	*	*	*		
— <i>glaber</i> , Jeff.	*	...	...	...	*	
— <i>nautilæus</i> , Flem.	*					
— <i>nitidus</i> , Müll.	*	...	...	...	*	
— <i>spirorbis</i> , Müll.	*	...	...	*	*	*
— <i>vortex</i> , Müll. (Morris)	...	...	*			

Table (*continued*).

Species.	Grays.	Ilford.	Erith.	Crayford.	Stoke New- ington.	Salisbury.
<i>Pupa marginata</i> , <i>Drap.</i>	*	*	*	*	...	*
— <i>umbilicata</i> , <i>Drap.</i>	...	...	...	...	*	
— (<i>Vertigo</i>) <i>antivertigo</i> , <i>Drap.</i>	*	*				
— (—) <i>pusilla</i> , <i>Müll.</i>	*					
— (—) <i>substriata</i> , <i>Jeff.</i>	*					
<i>Succinia putris</i> , <i>Linn.</i>	*	*	...	*	*	*
<i>Unio litoralis</i> , <i>Lam.</i>	*	*	...	*		
— <i>pictorum</i> , <i>Linn.</i>	*	*	...	*		
— <i>tumidus</i> , <i>Phil.</i>	*	*				
<i>Valvata cristata</i> , <i>Müll.</i>	*	...	...	*	*	
— <i>piscinalis</i> , <i>Müll.</i>	*	*	*	*	*	*
<i>Zua lubrica</i> , <i>Müll.</i>	*	*	...	...	*	*
<i>Velletia lacustris</i> , <i>Müll.</i>	*	...	...	...	*	