observed. Unilateral castration, unilateral vasectomy, and feeding the gland-substance had no influence whatever upon the prostate. Caspar disagrees with Sakur, who claims that with the loss of the testis-function something is withdrawn from the system that is necessary to the maintenance of the prostate. As agreeing with experimental and clinical facts, he adopts the view of Pezewalski, who holds that the division of the nerves of Cowper which accompany the vasa brings about an atrophy of the prostate, but not of the testis. The author declines to draw any practical conclusions from his experiments, and thinks it is too early for any accurate deductions from the clinical material so far at hand.

Cholecystitis Typhosa.—VON DUNGERN (*Münchener medicinische Wochenschrift*, No. 26, 1897) has been able to find four cases of cholecystitis in which the Eberth bacillus could be shown to be the cause. To these he adds the following: A woman, aged forty-six years, had typhoid fever in 1882; in 1887 she began to have severe attacks of cardialgia with vomiting. The same trouble recurred again the next year, and again in 1889, this time with icterus. No symptoms in 1895, when there was a periostitis of the lower jaw with necrosis. In October, 1896, there was colic, and in the region of the gall-bladder a tumor the size of a child's head, which proved to be an abscess connecting with the gall-bladder. The pus contained typhoid bacilli in pure culture. The blood-serum of the patient was very active, causing a complete Widal reaction in a dilution of 1 to 80, although the patient had had no typhoid for fourteen years. The author thinks it most likely that the bacilli entered the gall-bladder during the typhoid fever fourteen and a half years ago, and remained entirely latent for five years, causing the various attacks since.

It is possible that the primary infection may have been gall-stones, and that the gall-bladder was secondarily infected, forming a "liver typhoid," which did not involve the intestine because the germs were of low virulence and because the patient had an increased resistance due to the previous typhoid.

Etiology of Multiple Sclerosis.—BLUMBEICH and JACOBY (*Deutsche medizinische Wochenschrift*, No. 28, 1897) have studied the histories of the cases of multiple sclerosis in Gerhard's clinic, with especial reference to trauma as an etiological factor in this affection. In all twenty-nine cases were analyzed; twenty-three were men and six women. The fact that several times the affection began in childhood serves to bear out the theory of Ziegler, that multiple sclerosis arises upon the basis of a congenital disposition. Many of the cases had had infectious diseases; in only one, however (influenza), could any connection be made out between it and the onset of the multiple sclerosis. There was but one undoubted syphilitic. Six times there was a possibility of an intoxication, and two were alcoholics. All were carefully questioned as to trauma; eleven had been subjects of injuries. Five times the disease followed promptly upon the injury, in a sixth the multiple sclerosis followed at once upon the trauma; but this latter was due to a sudden dizziness. In the other five cases the trauma occurred a long time before the onset. Although the etiology of multiple sclerosis is not yet clear, the authors

think there can be no doubt that a relation exists between acute infections, intoxications and trauma. The trauma may cause the anatomical changes upon which the disease depends, or it may only occasion the breaking out of an already existing disease; again, it is possible that the trauma only plants the germ of the affection which may require a second influence to give it typical characters. On the other hand, it may be that the relation is the reverse, and that the giddiness and uncertain gait of the disease cause the trauma. A knowledge of the causes of multiple sclerosis will be of value in the way of prophylaxis and for treatment.

Röntgen Rays and Acute Dilatation of the Heart.—SCHOTT (*Deutsche medicinische Wochenschrift*, No. 31, 1897) in 1890 published a series of experiments showing that an acute dilatation of the heart could be caused by severe exercise. He was at that time obliged to depend entirely upon methods of physical diagnosis. Now he publishes a series in which the results of physical examination are corroborated by means of skiagraphs and diagrams from screens. The results showed that after exercise the heart was enlarged in all diameters, in one case as much 3.5 cm. laterally. The right ventricle not only showed the greatest increase in size, but its enlargement came on first and disappeared before that of the left ventricle. A bicycle ride of 5½ km. in twenty-three minutes caused a dilatation of 2 cm. in the right ventricle, and the apex to move outward 1½ cm.; a walk of 1½ km. in thirteen minutes, on the other hand, while causing a sweating as marked as the riding, brought about no increase in the size of the heart.

Anatomy of Gout.—RIEHL (*Wiener klinische Wochenschrift*, No. 34, 1897), opposing the view of Ebslein, says that in gout urate crystals are deposited in unaltered living tissues, and that the assumption that necrosis of the tissues must precede the deposit of the salts must be given up. Whether the tissue increase about the urate deposits is due to mechanical or chemical irritation is not yet clear. Gouty nodules are probably formed about as follows: First, a deposit of crystals in the lymph-spaces and vessels and in the connective tissue itself, due to this inflammation of the surrounding tissue with granulation tissue and giant cells. The author thinks that the failure of others to find urates in the healthy tissues is due in part to their having examined tissues from the cadaver, and in part to faulty methods of hardening.

Stomach Tetany and Autointoxication.—This affection is tolerably rare, about forty cases having been reported, of which not more than eleven are considered pure. The cause of the spasm is in doubt. The view that it is an intoxication has the most adherents, and is the one entertained by the author. If this theory is correct one would expect to find the poison in the stomach-contents and in the urine. GUMPRECHT (*Centralblatt für innere Medizin*, No. 24, 1897) has studied these secretions in a nineteen-year-old workman, who had dilatation from pyloric stenosis. The patient had typical tetany spasm lasting several hours and slowly disappearing through several days. The author reviews the literature of the subject and gives in detail the methods used in his own work. In conclusion, he says: "Neither