will develop very marked differences in oils which originally are excellent in quality, within from four to ten days. The author has found that masticating the oil with ordinary white bread will bring out any petroleum taste that may be present. This test is very delicate and makes it possible to eliminate oils with this objectionable taste. Brooks does not believe that various chemical tests that have been proposed to test the purity of liquid petrolatum are of any value whatever. He believes that the simple tests he has suggested are easily sufficient to determine the relative merits of different oils. The freezing test for detecting the presence of paraffin has also been proposed; but if paraffin does not separate from the oil at room temperature, it certainly could not do so in the body. The presence of a small amount of paraffin indicates that the oil has probably been prepared from one of the so-called "paraffin base" crudes, and does not indicate an adulteration.

Sodium-salvarsan.—WECHSELMANN and DREYFUS (*Munch. med. Wchnschr.*, 1915, lxii, 177, 178) report their clinical tests of a new salvarsan preparation given them by Ehrlich for trial. This preparation is called sodium-salvarsan and chemically is practically the same in fixed form as is obtained by adding alkali in the preparation of solution of salvarsan for intravenous injection. Their experience with this new preparation extends over a period of one and a half years. They both recommend it very highly on the ground that it has the same clinical action as the original salvarsan while it is readily soluble and therefore lends itself easily to intravenous injections. Sodium-salvarsan contains about 20 per cent. of arsenic and is a yellow powder which darkens on exposure to air and becomes insoluble and very toxic, hence it must be used in fresh solution as is the case with neosalvarsan. Solutions of sodium-salvarsan are alkaline and for that reason they are not to be recommended for intramuscular use. Both authors recommend the giving of sodium-salvarsan in doses of from 0.3 to 0.6 gram, two or three times a week. They have given in certain cases as high as forty and even fifty doses, but as a rule, from 4 to 7 grams are sufficient. They say that sodium-salvarsan injections give rise to fewer reactions than either of the other two forms of salvarsan. Weichselmann says he uses it freely in a great many conditions in which previously it was considered risky to try salvarsan therapy. He mentions among such complications chronic nephritis, optic neuritis, etc., and says that in certain cases he has noticed that albuminuria has improved following the use of sodium-salvarsan. Weichselmann considers that it is very important to use pure salvarsan therapy as he believes that the combined treatment has a tendency to increase the toxicity of the salvarsan to the danger point. He also does not approve of giving sodium salvarsan in concentrated solution by a syringe but prefers to give it in a solution, using 10 c.c. of a 0.4 per cent. salt solution for every 0.1 gram of the drug. Dreyfus, on the other hand, prefers to give it in concentrated solution in distilled water by means of a glass syringe, the total amount of fluid being about 30 c.c. He also strongly favors the combined treatment, using two doses of sodium-salvarsan and one of mercury per week. They both claim that sodium-salvarsan combines the more intensive action of salvarsan on syphilis with the simple technic of neosalvarsan.