Commercial Iron Silicides with a High Percentage of Silicon: Their Properties, Manufacture, and Uses. BY G. DE CHALMOT. *J. Am. Chem. Soc.*, **21**, 59-66.—The author has produced in the electric furnace iron silicides containing from twenty-five to fifty per cent. of silicon and probably consisting of the two compounds FeSi_2 and Fe_3Si_2 . These silicides are crystalline, white or gray, hard and brittle, unalterable in the air, and can be cast well with sharp outlines and corners. These alloys are recommended for castings, on account of the smooth finish and high luster they are capable of taking; for anodes for electrolysis in aqueous baths; and for abrasives. Their manufacture is carried out by heating iron ore, river sand, and coke, in a special form of electric furnace.

The Preparation of Zirconium Nitrides. BY J. MERRITT MATTHEWS. *J. Am. Chem. Soc.*, **20**, 843-846.—The compound $\text{ZrCl}_4 \cdot 8\text{NH}_3$ heated in a current of nitrogen decomposes and leaves as a gray residue the nitride Zr_3N_8 . By heating zirconium tetrachloride in ammonia gas there is decomposition with the formation of the nitride Zr_3N_8 .

Method of Preparing a Strictly Neutral Ammonium Citrate Solution. BY A. D. COOK. *J. Am. Chem. Soc.*, **20**, 585-586.—The author recommends stirring the citric acid solution after adding ammonia, the heat of neutralization being sufficient to drive off the excess of ammonia.

On the Segregation of Carbon in a Piece of Boiler Plate. BY HENRY FAY AND HENRY P. TALBOT. *J. Am. Chem. Soc.*, **20**, 614-617.—The authors have examined a piece of boiler-plate steel which, after having been twice annealed, cracked on flanging. On polishing and etching the surface, it was found that the steel was highly crystalline, and that the crystalline layer extended only part way into the plate on either side. Microscopical examination showed the carbon to be much segregated, and analysis of the borings taken from different parts of the plate varied from 0.03 to 0.156 per cent. of carbon. The probable cause of the brittleness was due to the annealing which was done at a temperature, 600° - 750° C., at which granulation of mild steel takes place, as has been shown by the experiments of Stead.

ORGANIC CHEMISTRY.

J. F. NORRIS, REVIEWER.

On Some Double Halides of Mercury. BY J. N. SWAN. *Am. Chem. J.*, **20**, 613-633.—The double chlorides of mercury with aniline and the toluidines were prepared and studied. Aniline, *o*-toluidine, and *p*-toluidine gave salts of the composition, $2\text{HgCl}_2 \cdot \text{BHCl}$, $\text{HgCl}_2 \cdot \text{BHCl}$, and $\text{HgCl}_2 \cdot 2\text{BHCl}$ where B repre-

sents the base. With *p*-toluidine but one salt was obtained which had the formula $\text{HgCl}_2 \cdot \text{BHCl}$. A mixed double halide of *o*-toluidine, $\text{HgBr}_2 \cdot \text{BHCl}$ and the following inorganic salts were also prepared : $3\text{HgCl}_2 \cdot \text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, $3\text{HgCl}_2 \cdot \text{SrCl}_2 \cdot \text{XH}_2\text{O}$, $3\text{HgCl}_2 \cdot \text{BaCl}_2 \cdot 8\text{H}_2\text{O}$.

The Double Halides of Tin with Aniline and the Toluidines.

BY ROBERT L. SLAGLE. *Am. Chem. J.*, 20, 633-646.—A systematic search for the possible double chlorides of tin with aniline and the toluidines led to the preparation of the following salts which are fully described : With aniline, $\text{SnCl}_2 \cdot \text{BHCl} \cdot \text{H}_2\text{O}$, $\text{SnCl}_2 \cdot 2\text{BHCl}$, $\text{SnCl}_4 \cdot 2\text{BHCl}$, and $\text{SnCl}_4 \cdot 2\text{BHCl} \cdot 3\text{H}_2\text{O}$; with *o*-toluidine, $\text{SnCl}_2 \cdot \text{BHCl} \cdot \frac{1}{2}\text{H}_2\text{O}$, $\text{SnCl}_4 \cdot 3\text{BHCl} \cdot 2\text{H}_2\text{O}$; with *m*-toluidine, $\text{SnCl}_2 \cdot \text{BHCl} \cdot \frac{1}{2}\text{H}_2\text{O}$, $\text{SnCl}_4 \cdot 2\text{BHCl} \cdot \text{H}_2\text{O}$; and with *p*-toluidine $\text{SnCl}_2 \cdot \text{BHCl} \cdot \frac{1}{2}\text{H}_2\text{O}$, $\text{SnCl}_4 \cdot 2\text{BHCl}$, $\text{SnCl}_4 \cdot 2\text{BHCl} \cdot \text{H}_2\text{O}$.

On Double Halides of Zinc with Aniline and the Toluidines.

BY DANIEL BASE. *Am. Chem. J.*, 20, 646-664.—The author gives a complete list of all the double halides of zinc with organic and inorganic bases which have been described. A study of the double chlorides of zinc and ammonium showed that but two salts are possible although eight have been described. These salts have the composition $\text{ZnCl}_2 \cdot 2\text{NH}_4\text{Cl}$ and $\text{ZnCl}_2 \cdot 3\text{NH}_4\text{Cl}$. A careful study of the possible double chlorides and bromides of zinc with aniline and the toluidines resulted in the formation of the following salts : With aniline, $\text{ZnCl}_2 \cdot 2\text{BHCl} \cdot \text{H}_2\text{O}$, $\text{ZnCl}_2 \cdot 3\text{BHCl} \cdot 2\text{H}_2\text{O}$, $\text{ZnBr}_2 \cdot 2\text{BHBr} \cdot \text{H}_2\text{O}$; with *o*-toluidine, $\text{ZnCl}_2 \cdot 2\text{BHCl} \cdot 2\text{H}_2\text{O}$, $\text{ZnBr}_2 \cdot 2\text{BHBr} \cdot \text{H}_2\text{O}$; with *m*-toluidine, $\text{ZnCl}_2 \cdot 2\text{BHCl}$, $\text{ZnCl}_2 \cdot 3\text{BHCl}$, $\text{ZnBr}_2 \cdot 2\text{BHBr} \cdot 2\text{H}_2\text{O}$, $\text{ZnBr}_2 \cdot 3\text{BHBr}$; and with *p*-toluidine, $\text{ZnCl}_2 \cdot 2\text{BHCl}$, $\text{ZnCl}_2 \cdot 3\text{BHCl}$, $\text{ZnBr}_2 \cdot 2\text{BHBr} \cdot \text{H}_2\text{O}$.

Sulphonation of the Paraffins. BY R. A. WORSTALL. *Am. Chem. J.*, 20, 664-675.—The normal paraffins hexane, heptane, and octane are readily sulphonated when they are heated with fuming sulphuric acid in an open flask with a return-condenser. A yield of 30 to 40 per cent. of the monosulphonic acid was obtained when successive portions of the acid were used and the temperature was kept at the boiling point of the hydrocarbon. When the vapors of sulphur trioxide are passed through the boiling hydrocarbons disulphonic acids are produced. The yield of the latter was about 40 per cent. of the theoretical. The mono- and disulphonic acids are thick sirups which form barium and lead salts, which are very soluble in water and deliquesce rapidly.

The Formation of Hydrazides by the Action of Phenylhydrazine upon Organic Acids in the Cold. BY VIRGIL L. LEIGHTON.

Am. Chem. J., **20**, 676-679.—Acetylphenylhydrazide was formed when a solution of phenylhydrazine acetate in acetic acid was allowed to stand four days. The formation of hydrazides from the phenylhydrazine salts of acids dissolved in the smallest quantity of the respective acid was shown to be a general reaction by the preparation of the hydrazides of formic, acetic, propionic, isobutyric, isohexylic, and heptylic acids.

Aliphatic Sulphonic Acids: II. Ethylenesulphonic Acids.

BY ELMER P. KOHLER. *Am. Chem. J.*, **20**, 680-695.—Ethylenesulphonic acid was prepared by decomposing 1,2-ethanedisulphone chloride with boiling water, by distilling 1,2-bromethanesulphone chloride under an increased pressure, and by heating the potassium salt of the acetic ester of isethionic acid at 185°. The acid forms well-crystallized ammonium, potassium, sodium, barium, and lead salts. The chloride of the acid prepared by treating the potassium salt suspended in chloroform with phosphorus pentachloride can be distilled under diminished pressure. Ethylenesulphonic acid is oxidized readily, the products being carbon dioxide, sulphuric acid, and water. Reduction is effected with difficulty: sodium amalgam in alkaline solution and zinc in acid solution are without effect. When the acid is heated with phosphorus and iodine, ethanesulphonic acid is formed. Ethylenesulphonic acid combines readily with water, especially in the presence of alkali, forming isethionic acid. When heated at 130° with hydrochloric and hydrobromic acids, addition-products are formed in which the halogen atoms are in the 2-position. With bromine, 1,1-bromethylenesulphonic acid is produced, the substitution being rapid and complete even at 0°.

The Ethers of Toluquinoneoxime and their Bearing on the Space Isomerism of Nitrogen. BY JOHN L. BRIDGE AND WM. CONGER MORGAN. *Am. Chem. J.*, **20**, 761-776.—Toluquinone-*o*-oxime obtained by the action of nitrous acid on *m*-cresol forms well-characterized acetyl, benzoyl, and methyl ethers. The benzoyl ether of toluquinone-*m*-oxime, prepared in the same way from *o*-cresol, was shown to consist of a mixture of two isomeric substances, which were separated by fractional crystallization. The same ethers were obtained from the oxime formed by the action of hydroxylamine on toluquinone, and as no toluquinone-*o*-oxime was formed under these circumstances it seems highly probable that the isomerism is due to the space relation of the oxime group. Isomeric acetyl and methyl derivatives of toluquinone-*m*-oxime were also obtained. The compounds mentioned and a number of their bromine derivatives are described fully.

Camphoric Acid. By WILLIAM A. NOYES. *Am. Chem. J.*, 20, 789-814.—By an error in his thermometer the author working with E. B. Harris, was led to think that the oxime of a ketone prepared from cis-campholytic acid (Harris, *this Rev.*, 3, 12), and the oxime of dimethyl-(1,3)-cyclohexanone (2) made by Kipping (*J. Chem. Soc.*, 67, 357) were identical. If the oximes were the same cis-campholytic acid would have the structure of 4'-tetrahydroxylic acid, $C_6H_4(CH_3)_2(1,3)COOH(2)$ and accordingly the synthesis of the latter acid was attempted. Although the preparation of the xylic acid will probably throw no light on the structure of cis-campholytic acid, the synthesis, which has been carried as far as the preparation of the neighboring xylic acid, is described, as a large number of new compounds has been prepared. The nitrile of the acid was obtained by the diazo reaction from 2-amino-*m*-xylene, but its saponification could not be effected. The xyloidine required was obtained from commercial xyloidine, in which it is present to the extent of about 2 per cent. An attempt was next made to prepare the xylic acid from dimethylcyclohexanone. In the preparation of the necessary dimethylpimelic acid, the dimethylpentanetetra-carboxylic acid obtained as an intermediate product was isolated and found to have properties different from those which had been assigned to it. Dimethylcyclohexanone was prepared from dimethylpimelic acid, but the attempts to convert it into the hexahydroxylic acid through the cyanhydrin or cyanide were unsuccessful. The next method of preparation studied was the conversion of ethylidenediacetic ester into a cyclic compound by distillation under diminished pressure with chloroform and subsequent treatment with bromine (Knoevenagel, *Ann. Chem. (Liebig)*, 281, 121). As only dimethyl-(1,3)-phenol-5-methyl-(2) acid, and its ethyl ester were obtained the method was abandoned. Unsuccessful attempts to prepare the xylic from acetobromxylene and acetoiodoxylene led to the preparation of 4-chlor-2-brom-*m*-xylene, 4-aceto-5-brom-1,3-xylene, bromxylic acid, $C_6H_4(CH_3)_2(1,3)COOH(4)Br(5)$, acetoiodoxylene, $C_6H_4(CH_3)_2(1,3)COCH_3(4)I(5)$, iodoxylylglyoxalic acid, $C_6H_4(CH_3)_2CO.COOH$, iodomethylterephthalic acid, $C_6H_4CH_3(1)(COOH)_2(2,3)I(5)$ and two iodoxylic acids. The preparation of the neighboring xylic acid was finally effected from acetomesitylene. Improved directions for the preparation of mesitylene from acetone are given by which the yield is greatly increased. The mesitylene was converted into the acetyl derivative which, when oxidized, gave a mixture of about equal parts of 2,6-dimethyl terephthalic acid and dimethylphthalic acid. The acid ester of the former acid was converted into the amide, which was changed into an amine by means of sodium hypobromite. The amido group was replaced by iodine, which in turn was removed

by zinc dust and ammonia. The resulting acid was the neighboring xylic acid sought. The author proposes to reduce this acid.

On the Non-existence of Four Methenylphenylparatolyl Amidines. BY H. L. WHEELER AND T. B. JOHNSON. *Am. Chem. J.*, 20, 853-861.—The authors have carefully studied the four methenylphenylparatolyl amidines prepared according to Zwingenberger and Walther (*J. prakt. Chem.*, 57, 209), and have found that the substances are not pure compounds. By fractional crystallization the amidines were shown to consist of two or more substances. By crystallizing together mixtures of the constituents found, crystals were obtained which had the properties of the amidines described and which formed double platinum salts and picrates agreeing closely in melting-points with those given by Zwingenberger and Walther. The fact that but one methenylphenylparatolyl amidine exists destroys the conclusions of Walther (*J. prakt. Chem.*, 55, 43; 57, 229) as to the structure of the anilides. Methenylmethylphenylamide-phenylimidine, $C_6H_5(CH_3)N-CH=NC_6H_5$, was prepared by mixing methylaniline with methylisoformanilide, by adding a mixture of methylaniline and formanilide to an excess of phosphorus trichloride, and by heating methenyldiphenylamidine with an excess of methyl iodide. The compound is a colorless oil, which forms a hydrochloride and a double chloride with gold.

Creatin and Its Separation. BY E. F. LADD AND P. B. BOTTENFIELD. *Am. Chem. J.*, 20, 869-870.—The amount of creatin in ox flesh was determined by the methods of Neubauer, Stradler, and Liebig and was found to be 0.250, 0.234, and 0.215 respectively. The same methods of analysis applied to the flesh of a dog gave 0.242, 0.230, and 0.218 per cent. The authors consider the high results obtained by Neubauer's method the most accurate.

A Double Citrate of Zirconium and Ammonium. BY S. H. HARRIS. *Am. Chem. J.*, 20, 871-872.—By adding an aqueous solution of ammonium citrate to a solution of zirconium chloride a double salt of the composition $Zr_2(C_6H_5O_7)(NH_4)_4$ was formed as a curdy white precipitate which was very deliquescent when dried.

On the Constitution of the Phenylhydrazones. BY PAUL C. FREER. *Am. Chem. J.*, 21, 14-64.—Two possible constitutions can be assigned to the compounds of phenylhydrazine with ketones and aldehydes:

When phenylhydrazine reacts with an aldehyde or a ketone there is, in all probability, a direct addition-product formed, which immediately loses water and passes into the hydrazone. This elimination of water could take place in two ways and would yield either a true hydrazone (formula 1) or a hydrazo compound (formula 2). That such addition-products are formed was shown by the fact that ethereal solutions of aldehyde and methyl-ethyl ketone, when cooled to -20° and mixed with phenylhydrazine, gave precipitates which dissolved and changed into hydrazones. The phenylhydrazones can be divided into two classes: those which are readily oxidized in the air and change to non-crystallizable tars with the evolution of gas, such as the phenylhydrazones of acetone, other aliphatic ketones, and acetophenone; and those which are perfectly stable or do not yield oxidation-products, such as the derivatives of aceto-acetic ester, pyruvic acid, and benzophenone. It is concluded that the hydrazones of the first class are probably hydrazo compounds and have the structure represented by formula (2) while those of the second class are true hydrazones (formula 1). This conclusion is justified by the fact that acetone-*p*-bromphenylhydrazone, when carefully subjected to the action of the air, is quantitatively changed to a red explosive azo compound, which is reduced by sodium amalgam to the original hydrazone. The azo compound exhibited the characteristic properties of true azo bodies, forming a crystalline perbromide, which decomposed in the manner typical of such compounds. Since the formation of an azo compound from a true hydrazone by oxidation seems highly improbable, formula (2) must be assigned to acetone-*p*-bromphenylhydrazone. The phenylhydrazone of acetophenone formed derivatives analogous to those described above. The two classes of hydrazones show a marked difference in behavior with benzoyl chloride. Acetonephenylhydrazone, which is readily oxidized to an azo compound, and has the hydrazo structure, readily reacts with benzoyl chloride and forms a product containing two benzoyl radicles, whereas benzophenone, having the hydrazone structure, reacts with difficulty and the resulting compound contains but one benzoyl group. This method of deciding structure can not be applied to the large number of hydrazones which do not react with benzoyl chloride. If the phenylhydrazones are formed by the elimination of water from an addition-product of the ketone and phenylhydrazine, it is possible that one ketone might yield two isomeric derivatives represented by the above formulas. Smith and Ransom (*Am. Chem. J.*, 16, 108), have proved the existence of two benzoinphenylhydrazones and considered them to be geometrical isomers. The author's method of study shows that the two substances have different structures. One is readily

oxidized to an azo compound by nitrogen trioxide with which it forms a red crystalline addition-product, which in turn forms a characteristic perbromide. The other, when treated in the same way, gives a white tetrazone. β -Benzoïnphenylhydrazone is converted into an isomeric substance by benzoyl chloride. The two acetaldehydrazones were studied in the same way but the results were not conclusive, since the two compounds change from one form to the other with great readiness. Doubt is thrown upon the structure of the compound prepared by oxidizing acetoaceticesterphenylhydrazone with mercuric oxide. The facts that the compound does not give a perbromide, that it is not reduced to the original hydrazone by sodium amalgam, and that it can be heated above its melting-point without decomposition, show that it is not benzene-azo- β -oxycrotonic ester. Chloraldehyde hydrate and phenylhydrazine yielded a substance of the structure

Note on the Action of Liquid Hydriodic Acid on Ethyl Ether.

By F. G. COTTRELL AND ROY RAVONE ROGERS. *Am. Chem. J.*, 21, 64-67.—When ether and liquid hydriodic acid were mixed a vigorous reaction took place and water and ethyl iodide were formed. By conducting the experiment in a specially designed piece of apparatus, an almost quantitative yield of ethyl iodide was obtained. This is noteworthy since Silva has shown (*Compt. rend.*, 81, 323) that with gaseous hydriodic acid alcohol was also formed.

Certain Alkaloidal Periodides and the Volumetric Estimation of Alkaloids as Higher Periodides. By A. B. PRESCOTT AND H. M. GORDIN. *J. Am. Chem. Soc.*, 20, 706-728.—By pouring a solution of atropine in dilute hydrochloric acid into a large excess of Wagner's reagent a periodide is precipitated containing nine iodine atoms. Under similar conditions strychnine, brucine, and aconitine give periodides containing seven atoms of iodine, while with morphine the precipitate contains but four atoms of iodine. The authors show that the formation of these perhalides is quantitative, and describe the volumetric estimation of the above alkaloids based on this reaction. Two double iodides of atropine and mercury are also described.

On Tetraphenylmethane. By M. GOMBERG. *J. Am. Chem. Soc.*, 20, 773-780.—Improved directions for the preparation of tetraphenylmethane from triphenylmethanehydrazobenzene are given. A molecular weight determination of the hydrocarbon and an analysis of the tetranitro-derivative furnish further evidence for the composition which has been assigned it. This is also shown by the fact that the nitro-derivative does not

form a sodium salt, whereas dinitrodiphenylmethane and trinitrotriphenylmethane give highly colored sodium compounds.

Hydrazo- and Azo-derivatives of Triphenylmethane. By M. GOMBERG AND A. CAMPBELL. *J. Am. Chem. Soc.*, 20, 780-790.—Well-crystallized hydrazo-compounds were formed by treating triphenylbrommethane with a large number of hydrazines. They oxidize in the air and form salts with dry hydrochloric, oxalic, picric, and other acids. The hydrazo-compounds are readily oxidized by amyl nitrite to azo-derivatives, which give off their nitrogen quantitatively when heated.

A Periodide of Triphenylbrommethane. By M. GOMBERG. *J. Am. Chem. Soc.*, 790-793.—Triphenylbrommethane unites with iodine in benzene solution and forms a periodide of the composition $(C_6H_5)_3CBr.I_2$, which crystallizes in hollow hexagonal crystals.

Perhalides of Quinoline. By P. F. TROWBRIDGE. *J. Am. Chem. Soc.*, 21, 66-72.—A large number of perhalides of quinoline containing one, two, and four added halogen atoms are described. Halogen addition-products of the compounds formed by the union of quinoline and methyl chloride, bromide, and iodide were also prepared.

Analysis of Aqueous Alcohol. By CHESTER B. CURTIS. *J. Phys. Chem.*, 2, 371-375.—The amount of alcohol in an aqueous solution can be estimated by determining the quantity of toluene necessary to cause a cloudiness in the solution. The method can be used only when the solution contains above eighty-five per cent. of alcohol and gives results as accurate as those obtained by the pycnometer test.

The Physical and Chemical Properties of Lithium Benzoate and Lithium Salicylate. By LYMAN F. KEBLER. *Am. J. Pharm.*, 71, 57-64.—The properties of samples of lithium salicylate and benzoate, which are on the market, and of the carefully purified salts, are described in detail.

The Electric Current in Organic Chemistry. By EDGAR F. SMITH. *Proc. Am. Assoc. Advancement Sci.*, 1898, 179-194.—This paper is a historical address delivered by the author as chairman of the Chemical Section of the Association.

GEOLOGICAL AND MINERALOGICAL CHEMISTRY.

W. O. CROSBY, REVIEWER.

Notes on North Carolina Minerals. By J. H. PRATT. *J. Elisha Mitchell Sci. Soc.*, 14, pt. 2, 61-83. The notes on four of