to be about, the apparatus is strapped to the body as already stated.

In using the same to drain the peritoneal cavity, I make use of the ordinary glass drainage-tube generally employed, which has a glass tube fitted within it, by means of which the fluid is siphoned away as rapidly as it accumulates within the drainage-tube proper. Where large areas have been denuded, I combine gauze drainage with this, bringing the end of the gauze up into the lower end of the drainage-tube. I believe that in this class of cases this method of drainage is most marked in its efficacy, owing to the fact that it acts continuously and cares for itself, thus obviating the constant attention which would otherwise be necessary on the part of the nurse, and where the risks of infection through incompetence in aseptic technic on the part of careless or ignorant attendants is considerable, as must frequently be the case, its advantages seem to me invaluable, as infection is absolutely guarded against.

While the principles involved in this method of drainage are well recognized and not new, the application of the same in the manner that I have detailed has not hitherto been made use of to my knowledge.

DISCUSSION.

DR. J. B. BULLITT, SR., of Louisville, Ky.—The device proposed by Dr. Jepson is certainly to be desired, provided it would work. I have experimented with a somewhat similar device myself and am prepared to say that it would not work in practice, and if it would, it would require a condition which the device is intended to prevent—the accumulation of fluid in a cavity to be drained under a certain amount of pressure; otherwise the siphonage would not occur until this pressure was supplied from behind, which would preclude using this apparatus in conditions where it would be desirable to use it as in suprapubic cystotomy. This apparatus would not be sufficient. It would presume that you could occlude this entrance and make it air-tight, which is difficult or impossible. The same would be true of all operative wounds. Unless an apparatus of this kind be operated with an artificial reservoir to work the siphonage apparatus, it is impossible to make it work. I have seen such an apparatus work for days at a time by means of an artificial reservoir of which I speak. The time can be graduated. This apparatus was devised by one of our Louisville surgeons, Dr. Chenoweth. I do not claim any priority for him, for perhaps we might find several hundred years ago some other surgeon might have done that same thing—but Louisville has done it again!

DR. FRANK WARNER of Columbus—I think that the gentleman has produced an apparatus that is very ingenious in its construction. Yet at the same time in one of the places in which he suggests its use, a very much better means of drainage is iodoform gauze, or the ordinary glass or rubber drainage-tube. I think we have been too much prone to drain, instead of closing our wounds. Yet while a large number have to be drained, we could use iodoform gauze in a vast majority of cases.

DR. D. S. FAIRCHILD of Clinton, Iowa—Referring to iodoform gauze, it has not been my experience that it serves well for drainage purposes. All who have drained chest cavities have been really inconvenienced by the constant soiling of dressings, and if this arrangement suggested by Dr. Jepson will work, it will certainly be a convenience, if nothing more.

DR. R. C. DUGAN, Eyota, Minn.—I rise to second Dr. Fairchild's remark as to iodoform gauze not making good drainage material. In doing emergency work I have used nothing but gauze, and I have come to the conclusion that unless it is wrapped in a rubber tube it is a first-class thing to stop drainage in pus cavities. The gauze very soon becomes filled with fibrin and makes a splendid plug.

DR. J. B. EAGLESON of Seattle, Wash.—I would state that the instrument referred to was described by Dr. Keen of Philadelphia, and was a modification of one by an English surgeon.

DR. JEPSON—In closing the discussion of my paper, there is little for me to say, except in reply to Dr. Bullitt's statement that the means which I employ in draining normal cavities will not work. I am surprised to learn that the doctor is so well acquainted with its inutilty, as I do not understand that he has ever employed the method. His opinion is certainly at variance with my experience, as I have by this method drained the peritoneal cavity in the instance of tubercular peritonitis

for sixteen successive days, with no attention on the part of the nurse other than emptying the receiving chamber when it was filled.

There is only one condition which can prevent the proper working of the method and that is the entrance of large quantities of air into the cavity to be drained; but as entrance of air can only take place about the drainage tube, it certainly occurs very rarely, if the wound is properly dressed.

ANATOMIC VARIATIONS OF THE NASAL CHAMBER AND ASSOCIATED PARTS.*

BY M. H. CRYER, M.D., D.D.S.

PHILADELPHIA.

No man who spends any considerable portion of his time in the study of practical anatomy, i. e., actual dissections, can fail to note how great is the number of anatomic variations he meets. So common are these variations that it can not be exactly said which is typical normal anatomy, and which are the anatomic variations. In plain words, anatomy is not a study to be classed among exact sciences. I do not mean by this that there is not such a regular basis of anatomic science that safe anatomic rules can not be made, but the more closely we study the subject the more variations as to details we are compelled to record. To be sure, given the mandible of an animal, a femur, or even a tarsal bone, the nature of associated bones, their size, positions and forms can be deduced, and by further study even an entire animal can be reconstructed. Admitting this, however, there are still as many variations in the internal anatomy as there are differences in the external appearance. Especially is this true in the anatomy of the human head. This special regional anatomy is modified by age, disease, occupation, climate, race, and many other conditions.

The skull consists of bones of both cartilaginous and membranous origin. In the fetus and infant the bones are soft and yielding; they receive deposits of calcific material until the bones become harder and harder as age advances. In the upper and lower jaws there are constant changes caused by development, eruption, and the loss of the teeth and the alveolar process.

Disease causes anatomic changes in the bony structures as well as in the other tissues of the body. In some disorders of the nutritive system the bones may not become incorporated with the full amount of calcific material, which would have the effect of leaving them soft and yielding; or, on the other hand, an undue quantity of calcium salts may be incorporated into the bones, with the opposite effect of making them hard and unyielding, thus modifying the physiologic functions that should take place in and about them.

Occupation will modify the general shape and character of the face and head, especially in youth. Those who are studious and pass an indoor life will have more delicate development of the face, with a larger brain-case, than those who are brought up with a laborious outdoor life. The mastication of coarser foods will develop the muscles of mastication and their bony attachments. Numerous other factors might be cited to show the influence of personal habit on the course of anatomic development.

Climate and its consequent environment has great influence on modifying the development of the bones of the head, as is demonstrated in the different formations of the skulls of the great races of the world.

There are also great variations in the shape, size and

*Presented to the Section on Laryngology and Otology, at the Fiftieth Annual Meeting of the American Medical Association, held at Columbus, Ohio, June 6-9, 1899.

markings of one side of the face from the other in the same individual. In bilateral bones, such as the frontal, sphenoid, vomer, and lower jaw, one side is generally found to differ from the other. This being the case it will be readily understood that the internal openings and spaces, such as the mouth, the nasal chambers, and the orbits, the maxillary, frontal and sphenoidal sinuses, the ethmoid and other cells, will differ accordingly.

The density of the bone varies also in different persons and in different periods of life in the individuals.

The illustrations I present herewith are made from my own dissections. I wish to apologize for showing

Plate 1 is what might be termed a typical skull of an adult person showing perfect occlusion of the teeth.

Plate 2 gives a front view of the same skull. These

Figure 1.

a few that have already been published; they are used on this occasion to demonstrate new points or to more fully establish points that I spoke of before this Section two years ago.

Figure 2.

In order to thoroughly appreciate the delicate symmetry and shades of differences which occur in the construction of the anatomy of the face, it is necessary first to study those skulls and bones that are considered typical at various periods of life, and to give some idea of typical bones I have introduced the following:

Figure 4.

two slides present an almost ideal anatomy of the external bony structure of the face. There is but the slightest variation of the two sides. It may be noticed that the upper right molar stands out slightly more than the left. The septum of the nose is seen to be deflected in the same

Figure 5.

direction, and on examination of the internal structures of the nose it is found that the bulla-ethmoidalis is enlarged on the left side, projecting toward the concavity of the septum. It may be taken as quite a constant anatomic law that when the mouth, palate and dental arches are bilaterally symmetric, the outer cranial

structures exhibit a like condition; even more, this bilateral symmetry will be found to extend throughout the body. However, this law, as all others relating to anatomic science, is open to exceptions.

Plate 3 (Fig. 4†) represents an interior lateral view of the skull, before published. It is now shown to draw attention to the close relations the upper teeth have to the antrum and the floor of the nose.

Plate 4 is taken from a skull about 6 years of age. The external wall of the alveolar process of the upper and lower jaw has been removed, along with some of the cancellated tissue, exposing the roots of the deciduous teeth and the crown of the permanent ones. It will be noticed that nearly all of the spaces of the maxillary bones are occupied by the dental organs, and even at this period of life there is but little room for maxillary sinus. By any interference with these organs at this period it will be seen how easily these teeth can be deflected in an abnormal position, interfering with the sinus or nasal chamber, and by these chambers being changed, the associated cavities and adjoining structures will be changed also.

Figure 6.

It will be noticed that the position of the developing second and third upper molars is very close to the under and posterior portion of the orbit.

In the paper on "The Relations of the Nasal Chamber and Accessory Cavities and Dental Lesions," that Dr. Kyle read before the Section (see JOURNAL, October 7) he referred to a patient of mine whose third upper molar tooth is situated in the upper portion of the distal wall of the maxillary sinus. Dr. Stout thought it strange that a tooth should pass upward instead of downward. In my opinion the tooth referred to has never passed downward from its place of development; through some inflammatory process the tooth with its capsule has become adherent to the posterior wall of the maxillary bone, the "intrinsic force" being insufficient to force the tooth into its proper position at the time it should have made its descent. It is somewhat of a similar nature to an adherent testicle in the abdominal cavity or in the inguinal canal.

† The numbers in brackets refer to illustrations given in a paper read by me before this Section in 1897, which see in JOURNAL, April 2, 1898, pp. 781-789.

There is no more reason to think that the tooth passed upward than to say a testicle has passed upward from the scrotum into the canal or abdominal cavity.

Plate 5 (Fig. 3†) is made from the skull of a seven-months' old fetus. It is a transverse bilateral section,

Figure 7.

passing just within the floor of the orbit. In the upper portion of this section will be seen two openings into the brain-case, with the crista galli and falx cerebri between them; below this is the nasal chamber with its septum; projecting from the upper walls of the chamber are the

Figure 11.

middle turbinals. In the middle meatus there is seen the unciform process forming the inner wall of the hiatus semilunaris; passing upward and a little forward from the base of the inferior turbinal to the outer side of this process is the hiatus semilunaris, from which there is a small opening communicating with the maxillary sinus, which is very small at this period of life. The greater portion of the maxillary bones, both superior and inferior, is occupied by the developing tooth organs. The section will show how nearly symmetric the skull is in the region at this time of life.

Plate 6 shows a perpendicular transverse bilateral section passing through the anterior portion of the orbit, maxillary sinus and first molar of each jaw on both sides. The eye has been divided in front of the crystalline lens. Between the upper portion of the nasal cavities are seen portions of the middle ethmoidal cells. It will be noticed how nearly symmetric the parts are. The maxillary sinuses are not quite alike, as the saw passes a little farther back on the right side than it did on the left. The floor of the antrum descends between the roots of the molars—a condition much more common among the white race than among the negroes, according to my observation. Abscesses on the roots of the teeth may, and sometimes do, cause perforation of the floor

Figure 12.

of the antrum, and the extraction of a tooth may be accompanied by the fracture of the floor of the sinus, a portion of it coming away with the tooth.

We admit that pus or infected matter will pass in the direction of the least resistance; we might suppose that by the close approximation of the apical foramen of the roots of the molars, when an abscess is formed at their points, the infected matter would pass into the antrum instead of through the alveolar process into the mouth.

Figure 13.

My clinical experience teaches me otherwise, i. e., an abscess caused from an infected tooth usually makes its fistula into the mouth. The only reason I can give for this is that Nature usually protects weak, important points in various ways. In this case I believe the mucoperiosteum lining of the sinus takes up its defense, and through its osteoblastic layer builds and protects its walls against absorption and perforation.

Plate 7 is a front view of a skull that is not symmetric. It will be noticed that the canine fossa of the

right side is lacking, the face being bulged out at this point. The teeth have not been in normal position. The septum is deflected over to the right side. In a skull of this character the internal structure will usually be very much out of symmetry. One might suppose that a large antrum would be found under the projection, but in this particular case it is not so.

Plate 8 presents a transverse section of the latter skull. It will be seen that the right antrum is smaller than the left, the fulness in the region of the infraorbital foramen and canine fossa being due to the thickness of the bone. The frontal sinuses of this specimen extend downward lower than usual, between the orbits.

Plate 9 (Fig. 11†) exhibits an entire lack of symmetry between the nasal fossa and the sinus of the right and left sides, the inferior meatus of one side being closed anteriorly by the deflected nasal septum and "spur" passing over and against the inferior turbinal and carrying it toward the outer wall of the fossa.

Figure 14.

In Plate 10 (Fig. 12†) it will be noted that a line or instrument passing through the axes of the alveolar process of the right side would perforate the nostril instead of the front of the maxillary sinus.

Plate 11 is from a different skull and has almost a straight septum and a bilateral symmetry as to the nasal chamber; the maxillary sinuses, however, vary throughout the depth of the skull. On the left side the antrum would not be reached by drilling through the alveolar process.

Plate 12 is also from a different skull, the transverse section made in the region of the first premolar. The septum is almost straight, the nasal chambers nearly symmetric on each side, but there is a great variation in the maxillary sinus. The outer wall of the nasal chamber of the right side is the outer portion of the maxillary bone, the floor of the antrum being on a much higher plane. In the floor of the left nasal chamber there will

be noticed an elevation which covers a tooth root, more than likely a supernumerary one.

In Plates 9 and 10, it may have been noticed that the nasal walls of the right inferior meatus, and in Slide No. 11 the left meatus, passed outward under the maxillary sinus to the facial portion of the maxillary bone; and in the distal portion to the zygomatic wall the same condition will be observed in Slide No. 12. We often read and hear of surgeons opening into the maxillary sinus through the canine fossa; if such an operation were performed on the skull of the specimen from which these photographs were made, the surgeon would go into the nasal chamber instead of the sinus, and I have known such to have taken place.

Figure 15.

Plate 13 is a transverse section made through a skull in the region of the second molar. The septum is straight, the nasal chamber and its appendages and the maxillary sinuses are almost bilaterally symmetric. The maxillary sinuses are small and do not extend downward in the direction of the teeth and alveolar process.

Plate 14 is a section from the same skull as is No. 13, cut more anteriorly, in the region of the premolars; here we find the parts are almost symmetric. The crista galli has been cut transversely, showing within the walls a cell of considerable size; its opening is into the frontal sinus.

Figure 16.

Plate 15 is made from the skull of an old person where the bones have been much resorbed; it is comparatively symmetric, with the floor of the sinus much higher than usual and the nasal fossa passing outward to the external portion of the maxillary bone.

Plate 16 is also from a skull where the floor of the nasal chamber extends outward over the alveolar process until it reaches the outer wall of the maxillary bone. The points of the palatal roots of the first and second

molars will be noticed in the floor of the nasal chamber. The floor of the sinus is well up on the side of the bone.

Plate 17 is from a transverse horizontal section through both orbits and the upper portion of the nasal chamber, giving a good view of a cross-section of the anterior, middle and posterior ethmoidal cells and sphenoid sinus. This picture also illustrates the ethmoidal cells situated principally between the two orbits in a horizontal section. Many of them are formed by the union of the orbital plates of the frontal and the ethmoid bone; others are alone associated with the ethmoid. The anterior ethmoid cells are the smallest of the three divisions, and open by several small orifices into the anterior portion of the hiatus semilunaris. The middle ethmoid cells vary

Figure 17.

more in size than the other two. The inner covering or wall is usually sphenoidal in form, and is known as the bulla-ethmoidalis. It is situated in the upper portion of the lateral wall of the hiatus semilunaris, and extends down and inward toward the unciform process. The openings of the cells are in the inner walls of the bulla-ethmoidalis and discharge into the hiatus semilunaris.

It is through the increase in the size of the bulla-ethmoidalis that disturbance is caused in the anterior and superior portion of the nasal chamber, the frontal sinuses, and the antrum of Highmore; by its enlargement it presses toward and against the septum, closing

Figure 18.

the space. When the enlargement is downward it presses more upon the unciform process and into the hiatus semilunaris, closing this semicircular groove and preventing the passage of the fluids from the frontal sinuses and the anterior ethmoidal cells into the middle meatus; consequently they enter the maxillary sinus, and through general inflammation of the parts there is an excess of fluids that can not find exit. This would interfere with the vitality of the teeth through the pressure on the nerves

and vessels passing through the sinus. It is from these nerves and vessels that branches pass into the apical foramen of the roots of the various teeth. The excess of fluids without an outlet will cause fulness of all the anterior cells as well as the frontal sinus, and possibly interfere with the anterior portion of the brain-case. It is this blocking of the hiatus semilunaris which causes the secondary or associated openings between the antrum and nasal chambers.

Figure 19.

It will be noticed that there is a lack of bilateral symmetry in the two sides of the septum. The anterior and middle ethmoidal cells of the right side are more numerous than the left at this point of the section, the posterior ethmoidal cells and sphenoidal sinuses varying in shape and size.

Figure 19 1/2.

Plate 18 is from a section made about the maxillo-molar articulation through which the maxillary sinus passes far into the malar bone.

Plate 19 gives a lateral view of the left nasal chamber; As far as I am acquainted with the text-books used in American medical colleges, they teach that there are three meati; this point should be corrected. Zuckerkandl has stated that about 62 per cent. of the skulls examined have four meati. I find fully that number in the skulls that I have made sections of, and in many

cases I find five. There are five in the illustration herewith. I do not know of five being mentioned by writers.

Plate 19 1/2 gives a view of a section cut near the posterior nares looking forward; it will be observed that there are six meati, the only case I have noticed showing six, in my section cutting.

Plate 20 gives a typical view of the lateral walls of the nasal chambers from the same skull. They are almost symmetric. It will be noticed that the ethmoid cells are

Figure 21.

exposed in the right picture. In both these, four meati are plainly seen and in one there are five.

The general teachings of our schools is that the fluids from the lachrymal glands flow through the lower meatus. The fluids from the frontal sinus, anterior ethmoidal cells and the maxillary sinus flow through the middle meatus, and the fluids from the posterior cells and the sphenoidal sinus flow through the third superior meatus. There should be added to the middle meatus the

Figure 23.

fluid from the middle ethmoidal cells. In the third meatus the most constant cell that empties its fluid is the one found in the orbital process of the palatal bone, while in the fourth meatus, "The Supreme," there are the fluids from the sphenoidal cells and the posterior ethmoidal cell. Where there are five meati, the fluids from the posterior ethmoidal cell flow into the fifth.

Before dismissing this illustration I wish to call attention to a pathologic condition. It will be noticed, in the region of the frontal sinuses, that there is a deposit

filling the space. It was spongy at the part shown. In one I have cut the outer wall away in order to more fully expose and examine it, and I found that the spongy condition became more and more solid as it extended outward until it was much harder than ordinary bone.

Plate 21 shows the outer portion of this hardened structure projecting into the open space of the lateral portion of the frontal sinuses.

Figure 25.

The two maxillary sinuses of this skull are smaller than usually found. It will be noticed that there are slight ridges in the floor of the sinuses suggesting the formation of septi. Complete bony septi of the maxillary sinuses are spoken of. I have been unable to find such conditions, although I have specimens which, when looked at in a casual way, could be classed as such.

Figure 26.

Plate 22 (Fig. 9†) shows a membranous septum extending well up into the sinus, with a round opening communicating with the posterior section of the antrum.

Plate 23 (Fig. 17†) is shown on this section to give an idea of a large sinus in proportion to the bone. It will also be noticed that there is a ridge or partial septum; this is due to a root having been left in, the action of the re-absorption at this point being arrested.

Plate 24 is also shown to give an idea of the size of the maxillary sinus of an ordinary skull.

Plate 25 shows what might be a bony division of the

maxillary sinus; but close investigation reveals that the crescent-shaped cavities situated on the upper posterior corner of the sinus are the cells of the orbital process of the palate bone, and passing a small probe from this opening it would enter the superior meatus of the nose.

Plate 26 is a section of a negro's skull, made through

Figure 27.

the molar teeth and the middle of the orbit. It shows a sinus at the upper posterior corner of the maxillary sinus. This opens into the superior meatus of the nose and belongs to the palate bones. It will be noticed that the antrum is of very small size.

Figure 28.

Plate 27 is from the left side of another skull. The section is made in the same region as the last, a very small, peculiarly-shaped antrum and a crescent-shaped cell which opens into the superior meatus.

Figure 29.

Plate 28 is from the left side of another skull showing two sinuses. The posterior one passes around the posterior border to the external surface of the anterior sinus.

The bony septum of this specimen might very easily

be looked on as a division dividing the antrum into two; but I would not thus classify it, as this posterior sinus also opens into the superior meatus, as do the others.

It is a well-established fact that the maxillary sinus is developed by an invagination of the mucous membrane of the middle meatus into the body of the superior maxillary bone. If we should have two of these invaginations from the meatus, I could easily accept as a fact that these cells were a divided maxillary sinus; but as the outlet of the posterior one is into the superior meatus, and, as the cells of the orbital process of the palate bone open into the superior meatus, I claim that this is an enlarged palatal sinus or cell and not a divided maxillary sinus.

Plate 29 is another section made through the molar teeth and the center of the orbit. Posterior to the maxillary sinus we find another sinus of a different character, and, to judge by superficial observation, it might be said that it was related to the maxillary sinus or those belonging to the palate bone. When passing a probe into this it leads into the supreme or fourth meatus of the nose, showing that it is related or connected with the sphenoidal sinus. In fact, it is a very large sphenoidal sinus extending out laterally in a line almost to the outer parts of the maxillary band.

Plate 30 gives a view of the external wall of the nasal fossa of the same skull, the sphenoidal sinus in the right hand picture extending outward until it passes into the

Figure 30.

opening posterior to the maxillary sinus in the preceding slide. It will be observed that this large sphenoidal sinus extends well forward toward the frontal bone and backward to the basilar process of the sphenoid bone.

Plate 31 gives an outer and inner view of the same section. In the right picture the cut is made passing through the premolar teeth and a little to the inner side of the middle of the orbit, exposing the inner wall of the maxillary sinus, the cell of the palatal bone, and the sphenoidal sinus, over which will be seen the sella turcica.

In the anterior clinoid process it will be noticed that there is a triangular opening which leads into the sphenoidal sinus; in the left picture the external wall of the nose will be observed, and in the region of the body of the sphenoidal bone a very large sinus at the bottom of which will be noticed a space under the sella turcica shown in the right picture.

By the illustrations shown it has been noted that we have many marked variations in the general character of the bony structures. It is evident that there must be some general principles underlying these changes. It may be well to define certain laws that influence them.

At the commencement of the growth of the embryo, and continuing throughout life, there are two forces constantly acting on the body, known as the intrinsic and extrinsic. If these two forces be normal in strength

and application throughout fetal life, the result will be a fully developed child; but if these forces be interfered with in any way by lack of nourishment or undue external pressure, the child will not be of a normal physique.

If for some reason there is an insufficient quantity of lime-salts assimilated into the bony tissue, the bones will be soft and will not give proper shape to the body, they being the framework, such as the steel framework is to our lofty buildings. The brain-case is liable to enlarge when intrinsic growth forces out the soft yielding structure, while, on the other hand, an over-amount of calcic salts will harden the bone, and cause it to resist the intrinsic forces and prevent them from giving proper shape. At times the deposit of lime-salts will close up the sutures of the brain-case and absolutely prevent the expansion of the brain. Microcephalic skulls are sometimes caused in this way. By the use of the surgical engine, artificial sutures have been made which allow the brain to extend, as these sutures permit the case to expand.

The slopes and forms of the skull of the various races are influenced by the growth of the brain. In other words, the brain, by its intrinsic forces acting on the bone-tissue, will cause the skull to develop according to the character of the individual or race. If the anterior lobes of the brain are of large size, the forehead will be

Figure 31.

carried upward and forward; if, on the reverse, the cerebellum is large, the occipital region extends backward with a low receding forehead.

After the birth of the child muscular action and various forces have direct influence over the change of the shape of the bones, according to the following general laws:

The normal application of forces to developing bone results in the normal development of the form of the bone. The abnormal application of forces under the same circumstances results in the development of an abnormal form. Abnormal applications of forces to bone in adult life will also change and modify the shape and character. The bones becoming changed, the whole body is modified according to the change in the bone tissue.

It is to be understood that the factor behind these anatomic variations leading to symmetric development is necessarily nutritional. Some logical interference with normal nutrition has brought about functional disturbance of a part, and this in turn a corresponding modification of anatomic form.

PROFESSOR LUBARSCH of Rostock, Germany, has been placed in charge of the department of pathologic anatomy in the new State Institute of Hygiene at Posen.