DIPHTHERIA.

BEFORE the New York Academy of Medicine, Dr. C. E. Billington read a paper on this disease and its treatment, which was especially valuable, as it was based entirely upon clinical and personal experience; his observations having been made with great care, and extending over a large number of cases. The records of the Bureau of Vital Statistics showed, said he, that in 1873 there were over four hundred deaths from diphtheria in this city; in 1874 over one thousand, and in 1875 no less than two thousand three hundred and twenty-nine. This terrible epidemic he thought could not be checked by any therapeutic methods, but could only be stamped out by the most revolutionary and active sanitary reform. Dr. Billington has enjoyed unusual facilities for the study of the disease, as he is one of the District Physicians of the Demilt Dispensary, and has seen altogether about three hundred cases, of which he has careful records of about one half.

As a result of his observation and study, he has become fully convinced that diphtheria is a local disease, at least primarily; and, though this is the opinion of a minority of the authorities on the subject, he is glad to have his views corroborated by such observers as Drs. Jacobi and J. Lewis Smith. This conclusion is based upon the following points:

- First. The local affection commences first.
- Second. The gravity of the general symptoms is in proportion to the severity of the local manifestations.
- Third. The results of treatment seem to substantiate this view.

In the study of the nature of the disease, he said, three elements were to be considered:

- (1) The contagium, which he did not propose to discuss on this occasion.
- (2) The inflammation, denuding the fauces of epithelium, and resulting in membranous exudation; and
- (3) The effects reflected from the inflammation upon the system in general are, to a greater or less extent, septicæmic in character.

Dr. Billington's treatment consists mainly in local disinfection, together with the most careful and unremitting watching and attention. The agents which he regards as most useful are the following, in the order in which they stand in his estimation: Tincture of the chloride of iron, lime-water and glycerine; and after them, salicylic and carbolic acids, sulphite of sodium, chlorate of potassium, etc. One formula which he uses in almost every case is as follows:

R—Tinct. ferri chlor., f dr. iss.
Glycerinæ.
Aque, aa f oz. j.—M.
S.—Teaspoonful every hour or half hour.

Besides being very effective, it has the merit of being pleasant to the taste, which is a great desideratum for children, especially when the dose has to be so frequently repeated. If the child is under two years, one drachm of the tincture of the chloride of iron is enough, and if vomiting follows the administration of the medicine, it should not be given so often.

In connection with the above, Dr. Billington formerly employed the following:

R—Potass. chlor., dr. iss.
Glycerinæ, f oz. ss.
Liq. calcis, f oz. iiss.—M.

A teaspoonful of this was alternated with a dose of the former; so that the patient would receive one or the other every half hour. As a substitute for the chlorate of potassium mixture, he now generally uses the following:

R—Acid salicylic, gr. x—xv.
Sodii sulphit., gr. xxx—xlvi.
Glycerinæ, f oz. ss.
Aque, f oz. iiss.—M.

Here the salicylic acid is rendered soluble by the addition of three times its weight of sulphite of sodium (borax also has the same effect), so that in this prescription we have the advantages of both these reputed antiseptics, which are indicated theoretically, and really seem to be of considerable practical benefit. It is of great importance that in every case in which it is practicable some sort of spray should be used upon the throat; and the most convenient instrument with which to accomplish this is the ordinary little perfumery spray-apparatus now in such general use. In order to annoy the child as little as possible, it is best to employ the spray immediately after a dose of the medicine is administered. The combination generally used by Dr. Billington is the following:

R—Acid, carbolic, m. x.
Liq. calcis, f oz. iv.—M.

He believes that the nasal douche or syringe has saved many lives; and even when the nasal passages, apparently, do not seem affected, it is often useful in reaching portions of the mucous membrane inaccessible to the spray. If, therefore, the breath should remain fetid after the employment of the latter, it ought to be resorted to; and the mixture mentioned above, containing the salicylic acid, is as good as any other for the purpose.

In adults or large children it may occasionally be of service to apply carefully strong tincture of iron (say two parts of the tincture to one of glycerine) to circumscribed patches of membrane; but, as a rule, topical applications of caustics or astringents by the probang or camel's-hair brush do much more harm than good, as they cause exhaustion of the little patients from their struggles to resist, excite an increased flow of blood to the part, and really occasion further thickening and spread of the membrane.

Dr. Billington expressed the opinion (which is hardly substantiated by other observers) that quinine is worse than useless in diphtheria in children; being objectionable, if for no other reason, on account of its bitter taste, which makes every dose dreaded by the patient.

In cases attended with high secondary fever, a full dose of quinine worked better in his hands. He can not subscribe to the prevalent opinion that diphtheria will never bear antiphlogistic treatment.

Dr. Billington then proceeded to give an interesting summary of the cases which he had personally observed, prefacing his statement with an allusion to the well-recognized disadvantages to be encountered in dispensary practice. According to his observations, about sixty-five per cent of all cases of diphtheria occur in persons under five years of age, and it is quite a rare affection among adults (except in the peculiar experience of certain irregular practitioners), even when individuals are constantly and to the fullest extent exposed to the disease. He has also found that about sixty per cent of all the cases will recover without any treatment at all, and that about five per cent will prove fatal whatever plan may be adopted. Out of one hundred and two carefully tabulated

dispensary cases treated by him, fourteen died, and eighty-eight recovered; while of seventeen cases in private practice, one died, and sixteen recovered.

The usual duration of the attack, from the commencement of the treatment to the disappearance of the diphtheritic membranes, was only from four to six days. Twenty-four cases in private practice, treated on the same principles by Dr. Wm. Darken, house physician to the Demilt Dispensary, show even a better result; as not a single death occurred directly from the disease, though one of the children died several weeks after the acute attack, from some unexplained cause.

A still later series of fourteen cases treated by Dr. Billington in conjunction with Dr. W. E. Bullard (in order that the patients might receive the fullest possible amount of attention) all recovered, so that we have fifty-five cases altogether, with only one death directly attributed to the disease. In a large number of these the attack was of very great severity.

From his observations, Dr. Billington has been induced to believe that a laryngeal complication can often be prevented or aborted by the use of the spray, and that even after the membranes have been fully formed in this locality it is of very great service. Calomel has also proved useful in many cases. The inhalation of hot vapor, he thinks, renders the system more favorable to the absorption of septic materials, and therefore injurious.

He did not express a positive opinion as to the identity of croup and diphtheria, but apparently seemed to hold to the former view.—*Canada Lancet*.

SULPHIDE OF CALCIUM.

By T. CURTIS SMITH, M.D., Middleport, Ohio.

I TAKE pleasure in calling attention to this remedy for action in a field where an agent that will produce the effects claimed for it is much needed—namely, in discussing glandular swellings which threaten suppuration, in furuncular eruptions, carbuncles, and chronic induration of various tissues, the result of inflammatory action. Sidney Ringer, of London, has done much to bring this prominently before the profession as a valuable therapeutic agent. He, however, signally fails to give the rationale of its mode of action upon the physical economy, further than to claim great alterative action for it. This is but another method of covering up our ignorance of the real actions of various drugs now in general use.

I have used this agent more or less for a year past, and am well satisfied that it possesses some valuable properties for difficulties of the classes above named.

In cases where glandular abscesses are threatened in scrofulous children, it has the effect to hasten their removal by either rapidly increasing their tendency to suppuration and rapid healthy granulation, or of removing them quickly by absorption without suppuration. This seems like a contradiction of terms, but we may well remember that other and apparently less potent agents have similar therapeutic value. Very often warm, moist applications are used to assist in alleviating inflammation, and the same kind of applications are also used to hasten suppuration. So we claim no more for this agent than has long been practically claimed in these different actions for a common poultice, but we hold that it is far more efficient.

We not unfrequently meet with glandular swellings in strumous children and adults, not of specific origin. These often become very large. Sometimes, after weeks of suffering and inconvenience from the swellings, they suppurate, and after opening are very slow to heal, and for weeks will discharge an unhealthy-looking pus; and when they have healed, too often leave very ugly cicatrices about the neck. If in such cases the sulphide of calcium is given, and without any other aid, these swellings, however large they may be, will soon disappear, unless suppuration has already begun. In the latter case, instead of a long, tedious course of waiting through days and nights of suffering, and increased general debility, they will rapidly hasten to a termination by the suppurative process. The pus formed under its use seems of a more laudable character than that commonly observed in strumous abscesses, and after being discharged, healthy granulations spring up; soon the swelling has disappeared, and the point of opening has closed, leaving very little trace of the former trouble. If no pus has already formed in these large, threatening, indurated glands, then under the sulphide of calcium they soon become less in size, and in a remarkably short time disappear.

Another use of no mean value for this agent is in the treatment of boils. Several times in the last year have I met with cases where a succession of crops of furuncles would appear. As one crop would commence to decline another would put in an appearance. In all of such cases, where the sulphide of calcium has been resorted to, they have speedily recovered. The boils already formed would quickly suppurate, and discharge profusely, the "core" melting away with astonishing rapidity. The little boils just commencing to form would rapidly disappear without ever becoming painful, and no trace of them could be observed. The points were notably gained in the case of an old lady who was so much annoyed from the boils that she could not sleep day or night, nor move in any direction without pain, often severe. In a few days she was relieved completely. Also in a recent case where a succession of these painful abscesses had occurred, those present rapidly disappeared, and have not yet returned.

In carbuncles I have had a more limited experience. Sidney Ringer is loud in his praise of the remedy in this very painful affection, and especially mentions the rapid solution of the indurated "core," and the formation and discharge of healthy pus, followed by rapid granulation, and not followed by a repetition of the swellings at other points.

In those children that are, from some constitutional vice, subject to a constant repetition of ill-conditioned sores, often greatly annoying them, and, from their very slow tendency to heal, last for weeks and months, the sulphide of calcium will soon effect a rapid recovery and prevent a tendency to their return.

I have recently used this remedy in a few cases of visceral indurations of long standing, with the effect of a rapid removal of the morbid condition and a speedy improvement in the general health. I am now trying its influence over some very intractable diseases, but can not yet speak of its power in these new directions.

I have noticed, in the use of this agent, that there has been a general improvement in the health and vigor, and that especially the nervous system seemed to become more efficient, and through this and its alterative power there would be a general improvement in the secretions; the skin and kidneys would act more freely; the bowels, if costive, would become at once regular; any tendency to acidity of the stomach would be partially or completely removed. I believe its influence in toning the nervous system is an effect fully as important as its alterative power. Time and an extensive use of

this agent will decide whether it will maintain the high stand it now seems to claim.

Ringer claims the same effect for the sulphides of potassium and sodium. I have only used the sulphide of calcium. Dr. Ringer recommends doses of one tenth of a grain repeated every two hours. It has been my habit to give to young children a half grain every four to six hours, and to an adult, from a half to two grains in sugar four to six times a day. Its odor is disagreeable, and similar to that given off by the Blue Lick mineral waters of Kentucky, and not unlike sulphuretted hydrogen. The taste, however, is easily concealed by mixing with sugar or making into pill. The solution in water can not well be attained, and if it could, it would be very unpleasant to the patient to take.

I have no reason to believe the agent would have any special power over syphilitic indurations of any kind, but have not extended the use of the agent in this direction to ascertain what power it might exert, except in a single case, where it was used in connection with mercury. The results were good, but whether the agent had any influence or not, I can not say.—*Southern Medical Record*.

CHLORIDE OF SODA IN CANCEROUS ULCERS.

By G. M. RIVERS, M.D.

ABOUT three years ago, I was requested to visit an old lady with an ugly ulcer of the left breast. The whole gland was involved in the disease, hard and uneven to the touch—the ulcer three inches in diameter, quite deep, with raised, ragged and irregular edges; in fact, presenting all the appearance of true cancer. I had previously attended two cases of cancer of the breast, both of which presented the same appearance and proved fatal. I held out no hope of recovery, but advised and instituted, as I believed, a purely palliative treatment. I directed the ulcer to be covered with simple cerate, and, to destroy the odor, to be washed twice a day with a solution of chloride of soda—Labarraque's solution, in proportion of three ounces of the chloride to five of water. In three or four months, with no other treatment whatever, except small doses of morphine to relieve pain, the ulcer had perfectly healed, the swelling and inflammation subsided, and the patient cured. It has now been more than two years, and the disease shows no disposition to return.

The second case upon which I tried this remedy was a young negro woman, about twenty, with half the left nostril and a large portion of the upper lip destroyed. This woman had been to the hospital in Charleston, and the physicians there had pronounced it cancer. Upon this case I used the chloride very freely, injecting into the nares as well as applying it externally. In four months the patient was cured, the last integument restored, leaving less deformity than I thought possible. In this case, also, I used no other remedy, except to protect it from the air with simple cerate.

The question will naturally occur, Were these cases true cancers? Their recovery alone makes me doubtful. I certainly had no doubts when I first saw them. But may not the chloride of soda be a remedy? It is as good as any thing else. Why not try it? Some of our most useful remedies have been discovered by accident.—*Southern Medical Record*.

A MULTIPLE ANTIDOTE.

DR. JEANNEL, of Paris, has endeavored to answer the question, Is it possible to prepare an agent which shall be official, that is susceptible of indefinite preservation, capable of neutralizing chemically all poisons in the stomach or in the intestine, or, at least, of transforming them into compounds relatively inoffensive, and then determining their prompt evacuation? He discusses the applicability of the agents suggested by Dorval for this purpose, pointing out what appears to him to be the excellences and defects of each, and finally proposes the following:

	Grammes.
Solution of ferric sulphate, s. g. 1.45.....	100
Water.....	800
Calcined magnesia.....	80
Washed animal charcoal.....	40

Let the solution of ferric sulphate be kept in a bottle by itself, and the magnesia and charcoal mixed with the water in another bottle. When required, pour the ferric solution into the other bottle, and shake violently. The mixture should be given repeatedly in doses of fifty to one hundred grammes.

According to Jeannel's experiment, this antidote, employed in suitable proportions, renders completely insoluble the preparations of arsenic, zinc, and digitaline, but not the oxide of copper. It leaves in solution notable quantities of mercuric oxide and appreciable quantities of morphia and strychnia, and does not decompose or precipitate either the cyanide of mercury or tartar emetic. It completely saturates free iodine, but acts only partially on solution of the alkaline hypochlorites.

THE PRESERVATION OF ICE AT THE BEDSIDE.

CUT a piece of flannel about nine inches square, and secure it by ligature round the mouth of an ordinary tumbler, so as to leave a cup-shaped depression of flannel within the tumbler to about half its depth. In the flannel cup so constructed pieces of ice may be preserved many hours—all the longer if a piece of flannel from four to five inches square be used as a loose cover to the ice-cup. Cheap flannel, with comparatively open meshes, is preferable, as the water easily drains through it, and the ice is thus kept quite dry. When good flannel with close texture is employed, a small hole must be made in the bottom of the flannel cup; otherwise it holds the water and facilitates the melting of the ice, which is, nevertheless, preserved much longer than in a naked cup or tumbler. A reserve supply outside the bedroom door can be secured by making a flannel cup, on the plan above described, in a pitcher, and filling it with little lumps of ice, care being taken that there is space enough below the bag to allow the water to collect and leave the ice dry. This provision will allow ice to be used during the hottest nights without the supply failing or the patient being disturbed—two very important considerations. The real therapeutic benefit of ice is only produced in some cases by its free use, and its soothing and stilling effect must be aided by the most perfect surrounding quiet.

FELIZET, of Elbeuf having observed that in epidemics of foot-and-mouth disease no beast affected with cow-pox is ever stricken with the former disorder, vaccinated thirty oxen, and not one of the twenty-five beasts effectually vaccinated showed any sign of foot-and-mouth disease, even after living for months among animals largely affected with it.

* This woman's father died with cancer.