will be much more readily obtained in cases in which a large perforation of the membrana tympani permits the water to come into direct contact with the wall of the labyrinth. This stimulation is also the more readily obtained by using water quite warm or quite cold. The force of the stream has naturally no influence on this stimulation.

The study of the occurrence of vertigo and nystagmus, both that observed in disease of the labyrinth and that brought on by syringing the ear with hot and cold water, has established the following facts, which are of great assistance in diagnosis not only of the labyrinth but of the cerebellum as well:

1. If the semicircular canals are normal and the ear is syringed with cold water, vertigo will result, and there will be set up a nystagmus increased by directing the eyes toward the opposite side. If the ear is syringed with warm water the same symptoms will occur, but the nystagmus will be toward the same side.

2. Should there exist an irritation of the endings of the vestibular nerve in the labyrinth, such as may be occasioned by a circumscribed suppuration in the labyrinth, there will be spontaneous nystagmus directed toward the same side. Syringing the ear with cold water will produce a positive reaction.

3. If there occurs a sudden destruction of endings of the vestibular nerve, such as would be occasioned by a diffuse suppuration in the labyrinth, there will be set up a spontaneous nystagmus directed toward the opposite side, but lasting only from a few days to several weeks. This nystagmus has its origin in the opposite normal ear. Syringing the affected ear with hot and cold water produces no response.

4. In case of long standing destruction of the nerve endings in the vestibular nerve, such as occurs in chronic diffuse labyrinth suppuration, there will be no spontaneous nystagmus and no reaction can be obtained by syringing the ear with hot or cold water.

5. In case of cerebellar disease, such as cerebellar tumor, cerebellar abscess or a meningitis in this locality, there will occur a spontaneous nystagmus directed toward the affected side.

It follows, therefore, that in case there exists pronounced rotating nystagmus, and the membrana tympani is normal, showing an absence of an inflammation in the middle ear, the presence of a cerebellar tumor should be suspected. On the other hand, should there exist pronounced spontaneous rotating nystagmus in a case of suppurative otitis media, without fever, but with severe deafness, and this nystagmus is directed toward the affected side, while the syringing of the ear with cold water produces no response, the diagnosis of a cerebellar abscess is the probable one. Here the complete destruction of the hearing in the affected ear would indicate a probable diffuse suppuration of the labyrinth. The failure to get the caloric response on syringing points to destruction of end-organs in the semicircular canals. The only spontaneous nystagmus which this lesion could produce would be directed toward the opposite side. The absence of rise of temperature would in most cases eliminate a meningitis as the source of the nystagmus, while a cerebellar abscess pressing on the vestibular nerve could produce a spontaneous rotating nystagmus, which would be increased by directing the eyes toward the affected side.

100 State Street.

INTESTINAL LOCALIZATION

WITH CERTAIN OTHER SUBJECTS RELATING TO THE
SURGERY AND SURGICAL ANATOMY OF THE SMALL
INTESTINE AND ITS MESENTERY

GEORGE H. MONKS, M.D.

Surgeon to the Boston City Hospital; Lecturer in Surgery at the
Harvard Medical School
BOSTON

This article is, with a few changes, the text of a demonstration on the cadaver which I gave in Boston, May 1, 1908, by invitation, before the Society of Clinical Surgery. The subjects considered have little to connect them, except that all of them relate to the surgery, or to the surgical anatomy, of the small intestine and its mesentery. The procedures described are taken mostly from cadaver studies which I have made from time to time. Some of these studies have already been published at greater length in former articles.¹ The object of the present article is to present, in a form as condensed and practical as possible, some of the procedures with which these studies were concerned. An effort has been made to elucidate the text by a free use of illustrations—a method which, useful as it may be in many instances, is unfortunately not wholly satisfactory as a substitute for actual demonstration.

Fig. 1.—Diagram showing on the abdominal wall a surface marking, which corresponds approximately with the root of the mesentery. (This diagram and the next one are taken from a former article, but a different figure has been used.)

Fig. 2.—Diagram showing on the abdominal wall three oblique lines. The dotted line (the same as already shown in Fig. 1) corresponds approximately with the root of the mesentery; the two heavier lines indicate the division of the abdominal cavity into thirds.

While some of the procedures described have repeatedly been employed with success in operations, others—which have not been used in operations—are presented merely as procedures which, from an anatomic point of view, are possible. Others, again, are simply anatomic demonstrations.

THE MESENTERY

As the mesentery will frequently be referred to in this article, it is desirable at the outset to make a few general remarks about this structure, especially in regard to its root, its shape, and the disposition of its different parts.

The Root of the Mesentery.—The part of the mesentery where the vessels enter is, as pointed out by Treves,

1. Intestinal Localization, *Ann. Surg.*, October, 1903; Studies in the Surgical Anatomy of the Small Intestine and its Mesentery, *Ann. Surg.*, October, 1905; Experiments in Flushing the Intestinal Canal with Salt Solution Through Multiple Enterotomy Openings, *Ann. Surg.*, June, 1908.

the real root of the mesentery. For the reason, however, that the mesentery is attached to the back part of the abdominal cavity for about six inches below this point, the term "the root of the mesentery"—in practice, at least—is usually considered to include the six inches of attachment. This root lies obliquely in reference to the spine, and, as shown in Figure 1, can be marked out with a fair degree of accuracy on the abdominal wall.

The Shape of the Mesentery and the Disposition of Its Different Parts.—Though the root of the mesentery measures only about six inches, the intestinal border of

Fig. 3.—A loop, with attached mesentery, from the small intestine of a thin muscular (male) subject. It belongs to the uppermost part of the bowel, as is shown by its large size, its obvious thickness, its vascularity, and by the presence of large and numerous valvulae conniventes; and also because of the presence of large, long and straight vessels in the mesentery radiating to the gut from arches mostly primary. There is very little fat in the mesentery, and the transparent spaces between the vessels are very extensive.

this structure measures thirty to sixty times as much. This means, of course, an enormous expansion—an expansion which, however, is greatest in the outer third or fourth of the mesentery. It is with this greatly elongated outer border, or "ruffled border"—a term I have applied to it—that the surgeon has most to do.

It is obvious that such a structure as the mesentery, in order that it may be accommodated in the abdominal cavity, must be thrown into many folds—folds which are even more numerous in cases in which the length of the intestine is greater than usual. These folds are, naturally, simplest near the root of the mesentery, where the diameter of that structure from above downward is comparatively little increased. Of these main folds, the first one, arising from the upper part of the mesenteric root, is generally directed to the left side of the abdomen; the one next lower down to the right; while those arising below proceed somewhat indefinitely from the lower part of the mesenteric attachment to both sides of the abdomen and the pelvis. The various coils of intestine, naturally, occupy the same regions as the corresponding parts of the mesentery.

INTESTINAL LOCALIZATION

"Intestinal localization" (a term which I suggested some years ago) means the determination by the surgeon of the part of the intestine to which any given loop, presenting in an abdominal wound, belongs.

In order to make this determination, the surgeon depends principally on two factors:²

1. A knowledge in regard to the part of the intestine he is most likely to meet with in a wound which has been made through any part of the abdominal wall.

2. A familiarity with the general characteristics of the intestine in the different parts of its course, and also with the characteristics of the adjacent portion of the mesentery.

1. *The Part of the Intestine Most Likely to Be Encountered in Wounds in Different Parts of the Abdominal Wall.*—Although the intestines may readily change their position, because of any one of a number of causes, and, although they may not be placed in exactly the same way in any two abdominal cavities, nevertheless the different parts of the intestine have their usual lodging-places, and it would seem to be worth while for the surgeon to know where these places are. Now, in a general way, the uppermost third of the intestine usually occupies the large cavity on the left side of the abdomen, high up underneath the ribs; the middle third occupies the middle part of the abdomen and the left iliac fossa; and the lowest third the right iliac fossa and the pelvis. Occasionally a loop from the lowest part of the intestine may be found high in the abdomen, but this is unusual; it is even more unusual for a loop from the highest part of the intestine to lie near the pubes—a statement the truth of which any one may test by attempting, on cadavers, to draw down such a loop.

The three compartments which the different parts of the small intestine usually occupy can roughly be

Fig. 4.—A loop, with attached mesentery, from the small intestine of the same subject as shown in Fig. 3. It belongs to the lowest part of the bowel, as is shown by its smaller caliber, its comparative thinness, the absence of vascularity and of valvulae conniventes; and also by the presence in the mesentery of comparatively short, small and somewhat tortuous vessels which radiate from arches mostly secondary or tertiary. The fat in the mesentery is more abundant than in the specimen shown in Fig. 3, and it approaches nearer to the intestine. For this reason the mesentery here is much more opaque, the vessels are somewhat obscured, and the transparent areas small.

indicated on the surface of the abdomen by means of two lines, drawn at the extremities of the oblique line already shown in Figure 1, and at right angles to it.

2. The absence of such pathologic conditions as might interfere with the process of localization, here described, is, of course, assumed.

This division of the abdomen into three compartments is shown in Figure 2. Now, speaking generally (for one should not forget that there are occasional exceptions) the compartment above the first line lodges the uppermost third of the small intestine, that between the first and second lines lodges its middle third, and that below the second line its lowest third. The surgeon may, therefore, knowing what compartment his incision has opened up, roughly determine what part of the intestine he is most likely to encounter there. If his incision is in the upper part of the abdomen, especially if it is in the middle line or to the left of it, any loop of small intestine he meets will belong probably to the uppermost third of the intestine; if his incision is in the middle part of the abdomen or in the left iliac fossa, the loops will belong probably to the middle third; and, finally, if his incision is in the right iliac fossa or to the right of the rectus muscle, or even in the median line between umbilicus and pubes, the loops will probably belong to the lowest third of the intestine.

2. *The Characteristics of the Intestine (and Attached Mesentery) in the Uppermost Third of the Bowel Compared with Those of the Intestine (and Attached Mesentery) in the Lowest Third.*—If the surgeon picks up a loop of small intestine which belongs to the uppermost third of the bowel (Figs. 3 and 5) he will notice that it is thick,³ and that in many cases it feels even "fleshy." Holding it now between the thumb and fingers of one hand, and stroking it downward in the direction of its axis, between the thumb and fingers of the other, he will distinctly feel the valvulae conniventes, as his fingers pass over them. He will probably observe

that the intestine is of large size and very vascular, being covered with numerous branching vessels. Indeed, everything about the loop will probably suggest to him a high degree of physiologic activity of this part of the bowel.

The amount of information which the surgeon will get by an examination of the attached mesentery depends largely on the amount of fat there. If, as one would expect in a thin subject, there is little fat in the mesentery, he will be able to see the blood vessels, to note their characteristics, and thus to get considerable

Fig. 6.—A loop, with attached mesentery, from the small intestine of the same subject as the loop in Fig 5. It belongs to the lowest part of the bowel, as is shown by its thinness, its small caliber, the absence of vascularity, and of valvulae conniventes; also by the fact that the vessels in the mesentery are hidden in a thick layer of fat. Little tabs of fat encroach on the wall of the intestine itself.

information as to the part of the mesentery he is examining; but, if there is so much fat as to conceal the vessels entirely, he will, for localization purposes at least, not secure very much information, unless, by transmitted light, he is able to get some idea as to the course of the vessels. Luckily, however, the fat is not often present to such a degree as entirely to conceal the vessels (Fig. 5), and they—at least that part of them nearest to the intestine—are usually fairly well defined.

In a case, then, where the vessels are visible, the surgeon will probably notice at once that these vessels are large, long and straight and that they radiate from the depths of the mesentery directly to the intestine. He will probably also be able to see, especially if he retracts the edge of the wound, that these vessels arise mostly from primary arches, deep in the mesentery. A few secondary arches may be seen, but these are usually more characteristic of the mesentery opposite lower parts of the intestine. The parts of the mesentery between the vessels will probably be more or less translucent, or, near the intestine, even transparent. These transparent spaces between the vessels appear like little windows, as it were, through which one may look; because of their shape and transparency I have

Fig. 5.—A loop, with attached mesentery, from the small intestine of a subject (female), which had less muscle and more fat than the subject which furnished the specimens in Figs. 3 and 4. This loop belongs to the uppermost part of the bowel, as is shown (just as in Fig. 3, though most of the characteristics are less marked here than there) by its large size, its comparative thickness, its vascularity, and the presence of large and numerous valvulae conniventes; also, by the large, long and straight vessels in the mesentery, which vessels radiate to the gut from arches mostly primary. There is a good deal of fat in the mesentery, and, because of this, the vessels are somewhat obscured, and the transparent areas near the intestine are very small.

3. The word "thick" is used here in a relative sense. It means that this loop of bowel is thick as compared with a loop from the lowest part of the same bowel. Vigorous males would be apt to have a thick muscular intestine, while in less strongly developed individuals, as, for instance, in women, the intestine would be thinner throughout. In any intestine, however, the upper part of the bowel is usually appreciably thicker than its lower part. Most of the other characteristics spoken of are also relative.

called them "lunettes." Sometimes these little spaces are so small that they can not be seen without firmly drawing up the loop of bowel, when, the mesentery being tense, they can usually be made out very close to the intestine.

Fig. 7.—A method of determining the real direction of the intestine by palpation of the root of the mesentery.

The liberal supply of large vessels in this part of the mesentery suggests at once to the surgeon's mind that more blood is going to the attached loop of intestine than is necessary for the nourishment of this loop; and, again, the idea of the high degree of physiologic activity of this part of the intestine is forced on his attention. Occasionally the surgeon may be able to see the lacteals in this part of the mesentery; for these little vessels, when filled with chyle, appear as milk-white streaks proceeding, in an irregular course, from the intestine to the deeper parts of the mesentery.

Turning now to a consideration of a loop of bowel—with its attached mesentery—from the lowest third of the intestine, the surgeon sees a different picture (Figs. 4 and 6). This loop of bowel is usually thin, and is generally of a smaller caliber than that of a loop from the uppermost part of the bowel.⁴ Few valvulae conniventes can be felt through its walls, or even none at all. Finally, it is probably less vascular than is the loop higher up in the bowel.

Fig. 8.—Method of using two sheets of rubber tissue for the protection of intestinal coils while they are outside of the abdominal cavity.

The vessels of the mesentery, if they can be seen at all through the fat, are comparatively short, small and often more or less tortuous in their course. If the edge of the abdominal wound is strongly retracted these vessels may, in case the mesentery is very thin, be seen to rise from secondary or even tertiary arches. It is unlikely that the surgeon will see any lacteals in the mesentery opposite this part of the intestine. Everything, in fact, will probably suggest to him a low degree of physiologic activity. This low degree of physiologic activity probably explains why such great lengths of this part of the intestine may be excised without seriously interfering with the nutrition of the patient. Excision of only a few feet of the upper part of the jejunum would probably kill him.

There is almost always more fat in this part of the mesentery than there is in that part of it attached to the upper portion of the bowel. For this reason the mesentery here is much more opaque. Moreover, the fat usually lies nearer to the intestine, a common characteristic of this part of the mesentery being, in fact, that

Fig. 9.—Method of using a single sheet of thin rubber tissue for the purpose of protecting the coils of small intestine during the period while they are outside of the abdominal cavity.

little tabs of fat project from it on the bowel itself (Fig. 6)—a condition rarely found in the mesentery opposite the upper part of the intestine, except in subjects with an unusual amount of fat.

Recapitulation.—The distinguishing characteristics, then, of a loop of bowel near the upper end of the intestine, and of the mesentery attached to it, are as follows:

The bowel is usually thick, of large caliber, very vascular, and it contains large and numerous valvulae conniventes, which can easily be felt through the walls of the intestine. The attached mesentery—if there is not too much fat in it—exhibits large, long, straight vessels radiating to the intestine from arches (principally primary) deep in the mesentery. There may be so much fat in the mesentery as to obscure the vessels, or even to hide them, but ordinarily the mesentery here is likely to

4. One should not forget, however, that, on the living subject, the apparent size of a loop of bowel as well as the thickness of its walls may vary in accordance with its state of contraction or distension. In some cases of chronic obstruction, also, the size of the intestine above the obstruction may be greatly increased, as well as the thickness of its walls.

be thin and to show the vessels, and between them translucent, or even transparent, areas close to the intestine.

On the other hand, the distinguishing characteristics of a loop of bowel near the lower end of the intestine, and of the mesentery attached to it, may be summarized as follows:

The bowel is usually thin, is likely to be of comparatively small caliber, and only moderately vascular; while probably no valvulae conniventes can be felt through its walls. The attached mesentery has relatively small, short, tortuous vessels, which spring from secondary, or even tertiary loops deep in the mesentery. There is usually more fat in this part of the mesentery than there is in that part opposite a loop near the upper end of the intestine, and, for this reason, the mesentery here is usually thick and opaque and the vessels are obscured, or, in very fat mesenteries, entirely hidden. In many cases little tabs of fat can be seen extending from the mesentery a short distance on each side of the bowel.

Fig. 10.—Method of drawing to one side the small intestine (enveloped in thin rubber) and retracting the abdominal wall on the other side, so as to bring into view the great left fossa of the abdominal cavity.

The characteristic features of the intestine (and its attached mesentery) in its upper part are thus contrasted with those of the intestine (and its mesentery) in its lower part, in order to emphasize the differences between them. The transition which these features undergo from one end of the intestine to the other is a gradual one; and it is by an estimate on the part of the surgeon as to the stage in this transition, which any given loop represents, that he is assisted in getting a fair idea as to the position of this loop in reference to the rest of the intestine. The thicker, the larger, the more vascular the loop, the more distinctly the valvulae conniventes can be felt through its walls; and, in the mesentery, the larger, the longer, the straighter the vessels, and the greater the transparency, or, at least, the translucency, of that structure, the more probable it is that the loop in question belongs to the upper part of the bowel.

On the other hand, the thinner, the smaller, the less vascular the loop, the less the valvulae conniventes can be felt through its walls; and, in the mesentery, the smaller, the shorter, the more tortuous the mesenteric

Fig. 11.—Method of inspecting the pelvis while the abdominal wound at its lower end is broadened and drawn downward and the small intestine, protected by a thin rubber covering, is gently drawn upward.

vessels, the thicker and more opaque the mesentery itself, especially if tabs of fat project from it on the bowel, the more certain it is that the loop belongs to the lower part of the bowel.

It is not by the presence of any one of these characteristics that the surgeon is able to get a satisfactory idea as to the real position in the intestine of any loop he may examine, for there is no single sign so characteristic of any part of the intestine as to enable him to determine what that part is, except at the two extremities where the intestine is fixed. He must, therefore, estimate the position of a given loop by a combination of characteristics just as he makes a diagnosis of a disease by a combination of symptoms. In making this estimate he is also influenced, as already stated, by knowing what part of the intestine he is most likely to meet with in the wound he has made.

Fig. 12.—The elevation of all the coils of the small intestine on an instrument specially designed for this purpose.

It is in this way that the surgeon localizes a loop of bowel, and, though he may never do so exactly, yet, with a little practice, he ought to do so near enough for his purpose. When he needs the information, this is certainly better than no localization at all.

THE DETERMINATION OF THE REAL DIRECTION OF THE BOWEL IN ANY GIVEN LOOP

To know the real direction of the bowel in any given loop (or, in other words, to discover which is the proximal end of the loop and which is the distal) is, of course, often useful to the surgeon; therefore, it seems appropriate here to describe a method by which, in cases in which the abdominal wound is large enough to permit palpation of the mesenteric root, he can, without much trouble, determine it.⁵ The procedure is as follows (Fig. 7): The loop is drawn well out of the wound, and its two extremities held upward by the assistant, the axis of the loop being kept by him parallel to the known axis of the root of the mesentery. The surgeon's right hand is now applied to the loop in such manner that his thumb is on one side of the mesentery

Fig. 13.—Small intestine suspended on a rod, after having been gathered up on it loop by loop.

and his first two fingers are on the other. The thumb and fingers are then gently pushed downward in the direction of the root of the mesentery. If, by palpation of the mesentery, the surgeon detects no twist in it, he at once knows that the upper end of the loop (as it is then held) will, if followed up, conduct him to the duodenum, and that the lower end will conduct him to the ileocecal valve. If, however, there is a twist in the mesentery, the surgeon should withdraw his hand, remove the twist by turning the bowel, and examine again. When, finally, no twist remains in the mesentery, the surgeon knows, just as in the first instance in which no twist was found, that the end of the loop which is uppermost is the one which leads to the duodenum, and that which is lowest to the ileocecal valve.

The same determination may, in certain cases, be

5. When I first made tests in reference to this procedure I thought that the principle on which this determination rested was new: but, later, I learned that it had already been suggested by Dr. R. F. Rand (Lancet, Dec. 22, 1883, p. 1083). I have also found a few references to the principle in text-books—of later date, however, than that of Dr. Rand's article. But, so far as I know, the idea of the surgeon's thumb and fingers "straddling" the loop, as it were, so as to allow palpation of both sides of the mesenteric root at once, is not suggested in any of these places.

made by inspection alone, if the abdominal wall on one side of the wound, and the coils of intestine under it, are strongly retracted.

PROTECTION OF COILS OF INTESTINE OUTSIDE THE ABDOMINAL CAVITY

As it often happens that the surgeon brings out a considerable part of the intestine through an abdominal wound, or that, in spite of his efforts to keep coils, when they are distended, within the abdominal cavity, they still protrude through the wound, it is obviously desirable to protect them so long as they remain outside of the abdominal cavity. Towels, which have been wet with hot saline solution and wrapped about them, are heavy, rough and easily displaced; they also soon get cold. In a number of operations on patients, as well as on animals, I have found the use of two sheets of thin rubber to be of much value in protecting these intestinal coils from injury and infection, and in keeping them warm and moist (Fig. 8). One of these sheets, with a short slit near its center, is laid on the abdomen in such position that the slit is directly over the abdominal wound. The intestines are then drawn out through the wound and through the slit, and are made to rest on the smooth upper surface of the sheet of rubber. If this sheet is large enough—it should be at least eighteen inches square—the bowels, on account of the restraining effect of the mesentery, can not reach beyond its borders. The second sheet of rubber is used to cover the bowel. If the surgeon wishes for any purpose to open the bowel, he can perforate the second sheet, and, through the opening thus made, he may draw out the loop of the bowel on which he intends to operate; or, if he prefers to do so, he may, without perforating the second sheet, draw this sheet around the loop of bowel, constricting the loop as much as he wishes by drawing a piece of the rubber sheet tightly about it, and fastening the ends of this piece together with the hemostatic forceps.

By means, then, of these two sheets of thin rubber the bowels may be kept moist and warm and protected from injury and infection.

If the lower sheet of rubber is large enough it may be turned up over the bowels in the way indicated in Figure 9, and thus be sufficient protection to them, without the second sheet being needed.

Even in cases in which, for the purpose of opening the intestine, only a single loop of bowel is drawn outside of the abdominal cavity, this loop may, before it is opened, first be drawn through a small slit in a sheet of this rubber, and the peritoneal cavity, by this maneuver, be sufficiently protected from the escaping contents of the bowel. It would seem that surgeons might well learn a lesson from the dentists in regard to the use of the rubber dam.

It is well to have these sheets of rubber sterilized, and, like rubber gloves, kept ready for use. They may be soaked in warm salt solution just before using.

INSPECTION OF THE ABDOMINAL CAVITY WHEN THE SMALL INTESTINE LIES OUTSIDE OF THE ABDOMINAL WALL

If all of the movable part of the small intestine is outside of the abdominal cavity, and rests between the two rubber sheets, the great cavities of the abdomen (that is, the fossa on the left side, that on the right, and, also, the cavity of the pelvis) may be inspected with comparative ease—and such of their contents as

remain examined—by gently drawing the rubber sheet and the bowels first to one side, then to the other, and finally upward, a broad retractor being used at the same time to draw the abdominal wall away from the mesentery (Figs. 10 and 11).

This same procedure has been carried out (on the cadaver, at least) even more satisfactorily by the use of a large instrument, shaped like a tuning-fork, and consisting of two parallel arms and a broad handle (Fig. 12). The instrument is well rounded and smooth, so as to minimize the possibility of damaging the intestine or mesentery. The arms are eight and one-half inches long, and the interval between them is a little less than three-quarters of an inch. The whole instrument is about fifteen inches long. It is intended for the purpose of elevating and supporting the intestines; during which process the mesentery, in a continuous flat sheet, occupies the long narrow interval between its two arms. This instrument is placed in position in one of two ways:

1. It may be held flatwise over the abdomen, and in such position that the long space between its two arms lies parallel with and directly over the abdominal wound. A loop of small intestine near the ileocecal valve is drawn out of the wound and pulled through the space between the two arms of the instrument, and then turned to one side or the other. The upper end of the loop is then gently pulled on, and as much more of the intestine as it is desired to take out is gradually drawn up through the opening and turned over one of the arms. Nearly the whole of the small intestine can in this way be drawn up between the arms of the instrument and made to rest on their upper surface. When the intestines are on this instrument, they can easily be moved *en masse* to one side or the other, thus making it easy to inspect the large fossæ of the abdomen, the edge of the abdominal wound being first, of course, strongly retracted.

2. The instrument may be passed at once under the mass of intestines, its two arms slipping along the two sides of the mesenteric root, the mesentery itself entering the space between them. This is done by taking up the lower end of the ileum into the space between the arms of the instrument, and then working them gently upward along the back part of the abdominal cavity, so as to include the whole of the mesentery along its base. This procedure will probably be found much easier to execute than the first method in cases in which the intestinal coils are much distended with gas.

DEMONSTRATION OF INTESTINE AND MESENTERY AT THE SAME TIME

For the purpose of demonstrating the mesentery and small intestine at the same time I have frequently gathered up the intestine on a rod, the rod being introduced into the intestine through a small wound a few inches above the end of the ileum, and the whole, or nearly the whole, of the intestine drawn over the end of the rod.

When this has been done and the rod, with the intestine on it, is drawn upward so as to stretch the mesentery, a very satisfactory view of intestine and mesentery together may be obtained (Fig. 13).

INTESTINAL TUBAGE

It was several years ago (1903), while demonstrating the intestines gathered on a rod, as already described, that the idea occurred to me of using a rigid tube in

much the same manner that I used the rod, the tube being for the purpose, however, of emptying an intestine full of gas and liquid feces. Since then I have experimented on many occasions with such a tube on the distended intestines of cadavers, and, in a number of instances, I have used the method on patients. The tube can not be pushed into the intestine, but the intestine must gradually, by means of a piece of gauze, be drawn over the end of it (Fig. 14). The tube opening into the intestine, for passage of the tube, may be made near one end of the intestine, and the intestine slowly drawn over the end of the tube, or it may be made somewhere near the middle of the intestine, and the intestinal canal emptied in both directions.

Fig. 14.—Intubation of the intestine on the cadaver. The intestine is being drawn over the end of the tube by means of a piece of gauze. Some of the tubes used in these experiments were straight and some were curved. The one shown at the lower left-hand corner of the picture has a curved extremity.

THE PELVIC FOLD OF THE MESENTERY

I can not close this article without referring to certain surgical procedures which, directly or indirectly, have to do with the lowest fold in the mesentery, which fold—for lack of a better name—I have ventured to call “the pelvic fold of the mesentery.” This fold consists of that part of the mesentery which is attached to the intestinal coils in the pelvis. The posterior border of this fold forms a definite landmark there. From a wound in the right iliac region, it can easily be reached and palpated by the forefinger of the operator; in fact, if the wound is large enough all the pelvic coils of intestine can be lifted out of the pelvis by pressure upward on this fold, to which, of course, the coils of intestine are all attached.

In case it is thought necessary to irrigate the abdomen from a wound in the right iliac region, it is well to know that the irrigating nozzle may best reach the great fossa on the left side of the abdomen by being carried behind this pelvic fold and then upward.

In connection with the technic of irrigating the abdominal cavity, it is well to remember that, in a way,

the mesentery may be looked on as a sort of partition, or fence, separating the two sides of the abdominal cavity. Though this partition may be curved, twisted and reduplicated in many different ways, yet it still remains a partition, and, in a sense, it separates one side of the abdominal cavity from the other. Now, for those operators who practice irrigation of the peritoneal cavity, it is well to know that the fossæ on both sides of this partition may easily be reached by the nozzle of the irrigating tube, through almost any wound where a loop of small intestine presents itself; for the nozzle may be passed down on each side of the mesentery attached to the loop toward the mesenteric root, care being taken to follow the surface of the mesentery closely. In this way the tip of the irrigating tube may, in almost any abdominal wound, be carried first to the bottom of one abdominal fossa and then to that of the other.

There is one more point concerning the pelvic fold of the mesentery which seems well to deserve the surgeon's attention. It is that whenever, through an incision in the right iliac region, the operator desires at

Fig. 15.—Method of picking up the lowest end of the ileum through an abdominal wound. The forefinger is seen, as it rounds the pelvic fold of the mesentery. The thumb (indicated in dotted lines) is on the other side of the mesentery. (Reproduced, with a few changes, from a figure in a former article.)

once to find the lower end of the ileum (as in perforated typhoid ulcer, for instance) he can do so by the following method: He passes his index-finger (two fingers are sometimes better) along the parietal peritoneum at the back of the pelvis, and around the base of the pelvic fold of the mesentery (Fig. 15). He now hooks his finger about the pelvic fold and closes the tip of the thumb on the tip of the finger, so as to grasp that fold and the loop of intestine which is with it. If he now allows the fold of mesentery gradually to slip away from between his thumb and finger, and then brings up the loop of bowel attached to it, he will find, in almost all instances, that he has in his grasp the ileum at a point close to the ileocecal valve. From this point he can then follow the ileum to any distance from below upward. If, on the other hand, he wishes to find the cecum, or appendix, he can be conducted to them by following the ileum for a few inches in the other direction.

SUMMARY

* The special points which I wish to emphasize in this article are as follows:

1. The surgeon can, approximately at least, tell to what part of the small intestine any given loop belongs.

2. He can tell the real direction of the bowel in that loop, provided the wound is large enough to allow him to palpate the mesenteric root; or, after the bowels and the abdominal wall have been retracted, to see it.

3. Intestinal coils outside of an abdominal wound can be efficiently protected between two thin sheets of rubber.

4. When the intestines are enveloped (in the sheets of rubber) the abdominal fossæ and the pelvis may be inspected much more thoroughly than when the coils of intestine are within the abdominal cavity.

5. As a demonstration, the mass of bowels, comprising nearly the whole of the small intestine, can be supported well above an abdominal wound by means of a special holder.

6. As a demonstration, the intestine and mesentery may be satisfactorily shown at the same time by gathering the intestine on a rod, and then elevating the rod horizontally above the abdominal wound.

7. In order properly to introduce a long rigid tube into the intestines for the purpose of emptying them, the intestines should be drawn over the end of the tube, and "gathered" on it in a manner similar to that in which, for demonstration, the rod was used.

8. The base of the pelvic fold of the mesentery is an important landmark in the lower abdomen, and it can usually be palpated from a wound in that part of the abdomen.

9. The great fossa on the left of the abdomen may be reached from a wound in the lower abdomen by carrying the tip of an irrigating nozzle well around the pelvic fold of the mesentery, and then pushing it upward.

10. Finally, the lowest coil of the ileum can be picked up at once between the surgeon's thumb and finger by causing them to grasp the base of the pelvic fold of the mesentery and the loop of bowel which is attached to it.

BIBLIOGRAPHIC REFERENCES

I wish to acknowledge assistance in regard to the anatomic disposition of the mesentery and the course of the intestine, from the excellent articles by Trevers, Henke, Sernoff, Weinberg and Mall, and also—concerning the arrangement of the mesenteric vessels—from the valuable paper by Prof. Thomas Dwight.

Treves: Three Lectures on the Anatomy of the Intestinal Canal and Peritoneum, Brit. Med. Jour., Feb. 28, 1885, p. 415 et seq.

Henke: Arch. f. Anat. u. Entwicklungsgesch. (His), 1891, p. 89.

Sernoff: Zur Kenntniss der Lage und Form des mesenterischen Theiles des Dünndarmes und seines Gekröses, Internat. Monatschr. f. Anat. u. Physiol., 1894, xl, 437.

Weinberg: Internat. Monatschr. f. Anat. u. Physiol., 1896, xlii, 89.

Mall: Development of the Human Intestine and Its Position in the Adult, Johns Hopkins Hosp. Bull., 1898 ix, 197.

Dwight: Anat. Anz., 1903, xxiii, 184.

67 Marlboro Street.

Deaf-Mutism in Idiots.—H. B. Sheffield, New York, states that congenital deafness is associated with mutism, but that even when hearing is intact few idiots are able to speak. Some of them by imitation learn to utter a few words, but their expressions usually bear no relation to any distinct desire or action, and they understand words spoken to them no better than what they speak. Moreover, their power of imitation is very much delayed in development or may never become manifest—all depending on the faculty of attention. In partial idiocy, such as mild forms of infantilism, cretinism, encephalitis or microcephalus, the power of conversation may reach a certain degree of potency, but the vocabulary is very limited and fragmentary.—*Post-Graduate*, February.