lant and opiate treatment ; but when it comes on after an occasional violent debauch, such remedies must be extremely improper. Adopt here everything calculated to remove gastric irritation. We have facts to show that most decided advantage may arise from the application of leeches, even where the symptoms of gastritis are absent.—*Ibid.*

EFFECTS OF IODINE.

BY DR. GRAVES, OF DUBLIN.

“If it be true,” says Dr. Graves, “that mercury may be made to affect the mouth more rapidly by means of giving sulphate of quinine to the patient during or after its use, as has been asserted by Doctor Harty of Dublin, there must be some remarkable difference between the effects of sulphate of quinine and iodine on the animal economy, a difference well worth investigating, inasmuch as these remedies are usually considered very similar in their mode of action, and are often ordered indifferently in the treatment of chronic diseases, such as scrofula. As Dr. Kluge observes, the modifying powers iodine exerts on the action of mercury, opens a new field of inquiry in the treatment of the venereal disease, and renders it an object of great interest to discover to what species or stages of syphilis one or both of these remedies is adapted.

Doctor Kluge’s experiments are of particular interest, when viewed in combination with the great encomiums recently bestowed in France and in London on the efficacy of iodine and deutiodide of mercury in the cure of syphilis and various other diseases. We have seen that two remedies, usually classed together as tonics, differ most materially in their effects, sulphate of quinine and iodine. The former stops the paroxysm of ague and induces salivation ; the latter does not stop ague, and checks mercurial ptyalism. Again, when during a course of iodine, the patient becomes subject to pain in the stomach ; this, as Lugol has shown, is best remedied by sulphate of quinine !

What are the therapeutical relations of another tonic, arsenic and iodine ?—The former certainly exerts a powerful influence in controlling ague, and I have seen it produce salivation. Would iodine be useful in guarding against or mitigating its occasional bad effects ?—If experience answered in the affirmative, no greater boon could be conferred on the practitioner, who would then be enabled to use long-continued arsenical courses in lepra, psoriasis, and various other diseases of the skin, without fear of producing unpleasant or dangerous effects on the constitution.”

In this brief passage there is much to give us pause. Dr. Graves has founded some curious conjectures on more curious statements. Our business is chiefly with the latter.

Dr. Harty, of Dublin, would seem to have asserted, that sulphate of quinine, given before or after the exhibition of mercury, accelerates the affection of the mouth by the latter. That this is opposed to common observation is merely an obvious truism. The experience of all surgeons and physicians has shown them that measures of depletion always augment the influence of mercury. Bleed a man and keep him low, and

mark how soon salivation is effected. Reasoning would lead the reflecting practitioner to suppose that converse circumstances would produce the opposite effects. And facts appear to warrant the suggestions of experience and judgment. We suppose it will be granted that few situations afford more ample opportunities of watching the effects of mercury than the wards of a Lock Hospital. Observation in those wards does not confirm the ideas of Dr. Harty. If a patient is weak, he is ordered the sarsaparilla or the quinine, in order that he may *bear* mercury, and not be rapidly salivated by it. Again and again have we given quinine with this view and with the same result. We join issue with Dr. Harty. We maintain that, *cæteris paribus*, quinine tends to prevent and not to produce salivation.

Dr. Graves observes that iodine and quina are usually classed together as tonics. They may be *classed* together in the schools, but the brittle and unsubstantial junction is like that of Nebuchadnezzar's image, one part iron and the other clay. They may be riveted—they never can amalgamate.

The best mode of testing the real action of a remedy, is surely to push it to the point of producing injurious effects. Apply this test to iodine and quina. The patient who takes the former grows pale, emaciated, nervous—has flutterings at the epigastrium—a tendency to syncope—a tremulous and frequent pulse. He rallies under stimulants. Is this the action of a tonic? The man who is under the influence of quina displays very contrary phenomena. The tongue is probably loaded—the face is flushed—the head may ache, and fulness in it is experienced—the pulse is full—the vascular system is excited and replete.

In these two series of effects we perceive no resemblance nor any analogy. The obvious deduction from their observation is that, *cæteris paribus*, iodine debilitates and quina strengthens.

Perhaps Dr. Harty or others may urge that though iodine, pushed to a certain extent, occasions the results we have described, smaller and more guarded doses possess less exceptionable tonic properties. They may probably refer to the instance of opium, which, in moderate quantities, is said to be a stimulant, and a powerful sedative in larger. We cannot allow them the benefit of this analogy, nor the use of this conjecture. We have seen iodine employed extensively, but we never yet saw it act as a tonic. We have indeed seen patients received into a hospital, every circumstance altered in their favor, and iodine given in small doses. Such patients have gained flesh and acquired strength, but we could not avoid entertaining the suspicion that the results were vicious, the experiment deceptive. Even in those instances, the most favorable we have witnessed, the mineral was seldom continued long without the superposition of unpleasant symptoms. So soon as the beneficial influence of better or of regulated diet, of repose, and of appropriate attentions had worn off, the depressing action of the iodine was more and more apparent.

There is a fallacy attending the operation of remedies like iodine, mercury, &c. which deserves to be adverted to. If they tend to effect the removal of a disease which is wearing down the powers of the patient, they act *pro tanto* and *pro tempore* as tonics. This is not unfrequently

the case with mercury, which, under ordinary circumstances, undoubtedly displays no tonic properties. Nothing is more common than to see a person affected with secondary, or even neglected primary, symptoms, acquire strength and embonpoint so soon as mercury is properly administered.

Our readers may gather from these remarks that we do not consider iodine a tonic in the fair and general acceptation of the term; nay, we deem the notion pregnant with danger, and lift our voice most earnestly against it. The inexperienced practitioner should know that in using iodine he wields a potent and a deadly weapon. If he employs it as he would a tonic, he will probably find occasion to lament his error.

There are two particular statements to which we shall allude before we close these imperfect observations. The first relates to the success derived from iodine in cases of secondary symptoms. Dr. G. observes that great encomiums have recently been bestowed on it, in reference to these affections, by surgeons in London and in France. Of this we are aware, and we had the pleasure of conversing with some who have advocated the remedy in public and in private. We have given it in several cases of node, and in some of cachectic ulceration of the skin and of the throat. We have also seen it frequently exhibited by others. Yet such has been our evil fortune, that we never *saw* it prove of service. It was lately prescribed for a patient with node, in the Lock Hospital. The man took five minims three times daily, and had a meat diet in addition. After the medicine had been continued for ten days or a fortnight, the patient became so debilitated and depressed, that the iodine was instantly desisted from, and brandy and ammonia substituted for it. In many instances it has seemed to do neither good nor harm; but in those the dose was small, and the patients had beer and good diet, which might probably counteract any mischievous effects. On the whole, we feel no confidence in iodine as a remedy in syphilitic symptoms.

The second statement relates to the administration of iodine in cases of mercurial salivation.

Now this is observed under three sets of circumstances. In one, the patient is in tolerable health, and salivation results from a very considerable quantity of mercury. It is said to depend, in such a case, on peculiar individual idiosyncrasy, and possibly the notion is correct.

In a second instance, salivation is the consequence of mercury, actively pushed for an acute inflammatory malady, as hepatitis or phrenitis.

In a third, salivation is induced by a moderate quantity of mercury, given as a course for syphilis, or for any other disease. In this case, the patient is almost always weak, or his constitutional powers are impaired, or he is kept too low, or his room is warm, or some analogous cause exists. We had under our charge a case which illustrates this position. A female who had drunk great quantities of gin was affected with tubercular eruption, after a primary sore. We put her on the pilula hydrargyri, five grains twice daily, and conjoined with it a diet of animal food. In a very few days she became salivated, and had some symptoms of the mercurial erythismus. We discontinued the mercury of course; but as soon as we thought she could again support it we resumed its exhibition, adding porter to the diet. Again and again the same set of

symptoms occurred. It struck us that the stimulus to which the patient had been previously habituated might enable her to bear the debilitating action of the mercury. We directed her to take a glass or two of gin during the day; and immediately a change for the better was observed, and the mercury was taken without inconvenience.

Thus, then, we remark that salivation may result from a small amount of mercury and peculiar idiosyncrasy—from mercury rapidly thrown in for acute inflammation—and from ordinary doses in debilitated patients.

Supposing iodine to possess the power of checking salivation, we apprehend that it would be applicable only to cases of the first description. In the second class, salivation is probably salutary, and, at all events, its speedy arrest would be dangerous, if not destructive. In the third class, salivation is the consequence of weakness and exhaustion, and we need not repeat our reasons for believing that iodine is not the remedy for those conditions. Fresh air, good diet, beer, and stimulants, are infinitely more adapted to them.

If iodine has really the virtue ascribed to it, that virtue may be safely and judiciously exerted in instances of salivation dependent upon idiosyncrasy. The cautious reader may hesitate in admitting even this as proved. We can only refer to the assertions of Professor Helmenstreich, and the corroboration they receive from Dr. Graves. We are irresistibly inclined to acknowledge a degree of doubt.—*Med.-Chir. Review.*

CHOLERA-PHOBIA AND CHOLERA ECCLESIASTES.

THE following amusing quotation from a work recently published, may not prove unentertaining even to medical readers, though its chief point is ecclesiastical.

“Tourists are the most fortunate people in the world. They seldom fail to find some remarkable incident or occurrence, just happening when they visit a place, as if for the very purpose of being put on record by their fertile pens. It was my good or evil star to be in Inverness when an event occurred there, unprecedented in the annals of that capital, or even of the Highlands themselves. On the very day that I took up my quarters in the Caledonian Hotel, another, and I have the vanity to think, a much less welcome visitor, arrived in the town—the INDIAN CHOLERA! Having been formerly on terms of intimacy rather than of friendship with this unhallowed stranger in his native country, I was apprehensive, at first, that I might be suspected of introducing him clandestinely, and in defiance of the quarantine laws; but my fears were soon dispelled, by learning that the blame was universally cast on the guard of a mail-coach, who had died of cholera, or rather of cold winds and hot whiskey, at some place between Aberdeen and Inverness. I was therefore at liberty to go about, and observe the effects of the panic on the inhabitants at large, without suspicion of being an infected personage myself. Had not the poor guard been dead, and consequently irresponsible, I think my conscience would have compelled me to take the blame on myself, as I was far more likely to have carried the dire contagion to the capital of the