question, unable to seize a morsel of food lying in the angle formed by the glass front and bottom of the tank, "raised himself into a slanting posture, the head inclined upwards and the under surface of the body towards the food," and, by waving his fins, caused a current in the water which lifted the food straight to his mouth. Mr. Romanes adds that this observation is practically worthless "from the observer having neglected to repeat the conditions in order to show that the movements of the fish were not, in their adaptation to these circumstances, purely accidental."

I quite agree with Mr. Romanes that such observations should be tested in every possible way, and I should have been only too glad to repeat the conditions of this and other observations had I been able to do so. The fact is, however, that as neither the directors nor the curator of the Manchester Aquarium were willing to call in the aid of those extra attractions which you London people seem to have successfully employed in the case of the Westminster institution, the Manchester Aquarium came to an untimely end, and thus my observations were cut short. There are, however, two comments which I should like to make. On p. 351 of "Animal Intelligence" (the coincidence in the numbering of the respective pages may help the reader's memory), Mr. Romanes quotes a story by Mr. J. S. Hutchinson concerning a Polar bear at the "Zoo." A bun was thrown into a pond, and fell "at the angle" beyond the reach of the bear. The animal thereupon "commenced stirring the water with its paw, so that it established a sort of rotatory current which eventually brought the bun within reach." This story was communicated to Mr. Romanes privately, and my skate story was published in your columns four years before Mr. Romanes published his book (see NATURE, vol. xix. p. 160). No repetition of the conditions is mentioned in the case of the bear, yet Mr. Romanes speaks of the story as "a most remarkable observation." In justice to Mr. Romanes I must add that he appears to accept the bear story as a proof of intelligence in that animal because it corroborates a similar story communicated to Mr. Darwin by another observer. I feel, however, that I have a right to back my skate against either of the bears named, for the following reasons. Had I repeated the conditions in the case of the skate with precisely the same result, it would have appeared as though the skate acted in obedience to inherited habit, or instinct, and even the similar conduct of the bears suggests this inference in their case. On the other hand, had a second trial with the skate failed, it would not have been proved that the first case was accidental, and therefore not the result of a "happy thought" on the part of the skate; for it might still have been contended that the skate, like a man, might display presence of mind on one occasion, and not on another, and the chief interest of the incident lies in the assumed spontaneity of the action. Finally, if Mr. Romanes will reflect upon the attitude of the fish as described in my narrative, I think he will see that the movements could not be "purely accidental." For, from the position of the skate's eyes, it follows that, when in the slanting posture described, he could no longer see the food. Yet he opened his mouth and adroitly caught it, the waving of the fins and the opening of the mouth being necessarily rapidly consecutive actions. This fact seems to me to show that he expected the food to rise in the way in which it did rise.

F. J. FARADAY

Manchester, January 21

YOUR correspondent seems to think that I had some particular spite against his skate, and quotes my indulgence to a bear as proof of inconsistency. But the two cases are very different. Even apart from the unconscious corroboration to which he alludes (and which as evidence of a fact I consider better than even verification by the same observer), we must remember that the stirring of water for a long time in the same direction with its paw is not quite so habitual an action on the part of a bear as is the ordinary swimming movement on the part of a skate. As for any difficulty which the skate may have had in seeing the food approach its mouth, surely the fact of its opening its mouth when the food was near enough to grasp is no better evidence of design than of accident. In either case, under the conditions, and more especially the "attitude," described, the seizure of the food at the proper moment can only be ascribed to the sense of smell, which in the skate is so highly developed. Lastly, why does your correspondent begin by saying that verification would have been desirable, and end by arguing that it would have been of no use? Even if the experiment had failed on repetition, he says, his inference would not thereby have been negated. If

this is so, assuredly there would have been no object in repeating the conditions. I once told a terrier to fetch me the ace of hearts from a pack of cards, and he did it. I happened previously to have known that the ace of hearts was the top card. Suppose I had repeated the experiment fifty times, and the dog had every time brought the wrong card, should I have been justified in attributing the first success to a "happy thought"?

GEORGE J. ROMANES

The Storm of January 26

I SEND you inclosed particulars of the great storm of January 26 and 27 as observed at Newport, opposite Dundee. Another observer six miles to the north-east of Newport took readings which corresponded almost exactly with those at Newport for the fall, but were thirty to forty minutes later for the rise. They were as follows, being reduced and corrected:—

Hour	Inches	Hour	Inches
2.15 p.m. Saturday,	28'429	11 p.m. Saturday,	27'385
3.15 " " "	28'218	11.15 " " "	27'385
4.15 " " "	28'036	3 a.m. Sunday,	27'665
6.15 " " "	27'834	6 " " "	27'922
8.15 " " "	27'598	9 " " "	28'143
10.15 " " "	27'406	11 " " "	28'230

Dundee, January 30

DAVID CUNNINGHAM

Ozone at Sea

DURING my voyage hither from London in the *Maranoa*, via the Canal, and calling at Malta, Aden, and Colombo, I was surprised at the low values for ozone as registered by Moffat's tests, which I pinned to the "uprights" in Stevenson's thermometer screen. I tried periods of exposure varying from half an hour to twenty-four hours, and the highest value noted was but 5.6 for eight hours (scale 0 to 10). The test papers, however, were always tinted, more or less, sometimes to 3.0 in half an hour, whereas tests exposed at the same time and examined when eight hours had elapsed, only gave 4.6. At Ben Nevis and Fort William, and in the moorlands of Staffordshire I have recorded far higher ozone values than at sea under the same force of wind and like periods of exposure. From my long experience of these tests I cannot consider them satisfactory; but in the absence of a more reliable method I would strongly suggest that they would give results more intercomparable if uniformly exposed for an agreed hourly period, especially at the various land stations.

CLEMENT L. WRAGGE

Adelaide, South Australia, December 22, 1883

Meteor

AT 9.55 p.m. on Sunday, January 27, I saw a meteor start from a point in Taurus, near to Saturn, and fall vertically a distance of 20°, and then burst with a brilliant flash, giving off several colours that almost instantaneously died away. The meteor was visible about three seconds, and increased greatly in brightness from the time first seen until it burst. It was the most brilliant meteor I ever saw, and its greatest brightness much exceeded that of Venus.

E. HOWARTH

Museum, Sheffield, February 5

Ravens in the United States

OUR Natural Histories say *Ravens* are common all over the United States, but I have never met any one who was aware of having ever seen one. Are they common in Westchester County, near the Hudson, and confounded with crows?

New York, January 11

MANHATTAN

Unconscious Bias in Walking

THE thought has occurred to me that "unconscious bias in walking" may be the result of inequality in the length of the lower limbs caused by the manner in which young children are carried. Each person appears to nurse solely on one arm; I think the right is more frequently employed. I have noticed when a child is held in the arm the side which is nearer the nurse appears to be in a somewhat cramped and unnatural position, the leg more or less bent, while the outer side is comparatively straight and free. Would not this, while preventing the full play of the muscles of the inner leg, tend to arrest to some extent its

proper development at a time when growth is very rapid, and thus cause that difference in the length and strength of the limbs remarked by your correspondents?

SARA S. OWEN

4, Soames Street, Grove Park, S.E., February 2

ON THE HEIGHT OF THE AURORA BOREALIS

IT is with pleasure that I respond to the invitation of NATURE to give an account of the work of the Danish Meteorological Station, which was maintained, under the international scheme, at Godthaab in Greenland, in 1882-83, and of which I had the honour of being the chief. I intend, in the present article, to confine myself to the aurora borealis.

The results, which have been obtained from calculations of the height of the aurora borealis in the temperate zone, which lies south of the so-called auroral belt, all agree in fixing the minimum height of the aurora very high, as the auroræ seem to be confined to the part of the atmosphere where its density is only a fraction of that at the surface of the sea. However different the value may have been of the heights of the auroræ observed outside their true zone, the average of the minimum heights is hardly under two hundred kilometres. On the other hand, the observations in the Arctic regions show that the auroræ may descend to much lower elevations above the earth's crust, and that they may even reach down into regions of the atmosphere where the density is about the same as on the surface of the sea.

Dr. S. Fritz has thus, at Ivigtut in South Greenland, in February and March, 1872, measured auroræ the lower edges of which were only from 50 to 200 metres above the level of the sea, while in nearly every monograph of the aurora borealis cases are cited in which the auroræ appear to have reached much further down in the atmosphere. I may further mention some instances, which have, by the bye, not been made public before, observed by the eminent zoologist, Prof. Steenstrup, and which he has permitted me to publish here.

During Prof. Steenstrup's sojourn in Iceland, 1839-40, he saw, on several occasions, auroræ which hid the top of the mountain Esia, some 600 metres in height, lying behind Reykjavik. He further states that he has seen auroral streamers between the masts of a ship, in such a manner that they disappeared where there were sails, and reappeared where the space was free. The Professor asserts even that on one occasion, on January 28, 1840, when walking between Reykjavik and Bessastad with the chief magistrate, Herr Tvede, and Judge Jonasson, he, as well as these two gentlemen, saw auroral streamers appearing *between themselves*. The phenomenon was not a solitary one, but occurred three or four times during this walk, and in spite of the pedestrians keeping about a yard from each other.

Although many estimates of the low descent of auroræ in the Arctic regions may have been due to optical illusions, specially through irradiation, one cannot, even from a casual observation of this magnificent phenomenon, but come to the conclusion that, while some auroræ lie in the same great, indefinable distance from the observer as those observed in the temperate zone, there are others whose whole appearance has the character of being a phenomenon of a purely local nature. During our stay at Godthaab this point had my special attention, as it appeared to me of importance to demonstrate by measurements as accurate as possible whether this subjective impression answered to the true facts.

To this end the Danish international station at Godthaab, 64° 10' 36" N. lat. and 51° 40' 0" E. long., has, during October and December, 1882, effected a series of measurements. The site of the station was particularly suited for the solution of the problem, as it lies just at the northern border of the great Arctic auroral belt, *i.e.* in a place where the auroræ appear with all the peculiarities which distinguish them in their true zone.

The distance between the two points of observation, separated by the Godthaab Fjord, was 5·8 kilometres, and the direction between them coincided with the magnetic meridian. The two instruments used for the measurements, exactly similar in construction, were arranged as universal instruments. Instead of a telescope, a tube was employed, which had in one end a small opening, and in the other a metal cross of very fine wires. In order that the errors in the observations should not affect them very much, measurements were only made in the vertical plane between the two points of observation. The placing and reading of the instruments were effected by means of pre-arranged fire-signals, and only those measurements of which the reading signals were instantaneously answered, and for which the time of reading exactly coincided, were recorded. Only the lower edges of the auroral bands were measured, as these are nearly always the most clearly defined.

We have, during our evenings of observation, measured the height of thirty-two auroral bands. The subjoined figures, showing the result of these, demonstrate that the lower edge of the band certainly descends very low. Thus of the thirty-two auroræ measured by this method ten only had a parallax under 1°, for six the parallax was between 1° and 2°, four had a parallax of between 3° and 4°, two one between 5° and 6°, four one between 7° and 8°, while we measured six of 10°, 14°, 15°, 17°, 86°, and 143° respectively.

Leaving the auroræ whose parallax was under 1° out of the calculation, I have found the following heights for the other twenty-two lower edges:—

1	...	67·81 kilometres.	...	12	...	7·43 kilometres.
2	...	59·60	...	13	...	6·16
3	...	54·73	...	14	...	5·28
4	...	46·94	...	15	...	3·72
5	...	45·04	...	16	...	3·69
6	...	38·09	...	17	...	3·22
7	...	29·81	...	18	...	2·87
8	...	19·14	...	19	...	1·99
9	...	9·76	...	20	...	1·96
10	...	9·40	...	21	...	1·35
11	...	7·67	...	22	...	0·61

The three heights of 1·99, 2·87, and 3·22 kilometres belong to the same aurora as that whose edge was measured at an interval of two minutes between each measurement. The two auroræ of 1·35 and 0·61 kilometres stood both above the fjord between the observatories. From the southern one they were seen at an altitude of 13°·6 and 30°·3 respectively above the northern horizon, while at the northern station they were 80°·5 and 7°·25 respectively above the southern horizon. These two and the above-mentioned third, the height of which was measured three times, had the appearance of curtains with large folds, or of bunches or wreaths of streamers lying close together, separated by darker, faintly-shining spaces, but connected below through a common band. The others were bands or arcs without radiating streamers. The edges measured were all nearly at right angles with the magnetic meridian. Only the height of those edges which were distinctly defined, and whose course did not deviate greatly from perpendicularity on the magnetic meridian, were measured.

I must further, as regards the small height of the auroræ observed at Godthaab, state that not only three observers besides myself, but even a student as conversant with auroral phenomena as Herr Kleinschmidt, all agree that auroræ were seen *below the clouds* on several occasions during the winter of 1882-83. On this point we do not entertain the least doubt.

In conclusion I will describe some observations made on several occasions during our stay at Godthaab of the peculiar type of the aurora known as phosphorescent auroral clouds.

On September 21, 1882, Herr C. Petersen, one of my