The condition of the parts involved in the abscess at the left carpus, was the same as had been found at the top of the sternum. The ulna was disarticulated; its coronary ligament was destroyed; the carpal end of the ulna bare, partially carious, and of a dark red colour. The capsular ligament of the carpus was also in part destroyed, and the articular cavity filled with sanious pus; the bones forming that joint were all tinged with the hue of inflammation, but the two bones corresponding most nearly to the carpal end of the ulna were quite bare of cartilage, rough, and of a very dark red colour. The forearm was separated from the arm, and its muscles carefully dissected away, in order to preserve the carpus as a morbid specimen; and in this state the ulna presented a striking appearance at its humeral extremity. The olecranon, and the sigmoid concavity, but especially the coronoid process, were of a high red colour, not superficial, but evidently pervading the solid structure of the bone, as the colour was not discharged or diminished by maceration. The shaft of the ulna was of the same deep red hue for an inch in advance of the coronoid process, but in and around the latter the red colour was particularly intense. On careful inspection of those points of the ulna, I was convinced that the colour was an effect of a morbid state of the bone, and not a casual tinge from stasis, post mortem infiltration, or other common causes; the bones of the same arm no where else presented similar characters, and I could not but draw the conclusion, that had the subject of this violent disease lived a few days longer, abscess, with gangrenous caries, would have been developed within the elbow joint.

Baltimore, Dec. 22d, 1828.

ART. III. *Remarks on the Use of Prussic Acid.* By GEORGE HAYWARD, M. D. of Boston.

THERE can be no doubt, I think, in the minds of those who have any practical experience on the subject, that Prussic acid is possessed of valuable, as well as powerful properties, and yet, if I am not mistaken, the use of it as a remediate agent has diminished in no small degree within the three or four last years. This must be attributed to the extravagant praise that has been bestowed on it, to its having been extolled for properties which it does not possess, and recommended as a specific for diseases, over which it has since been found