number of inunctions or injections of mercury or salvarsan injections necessary to convert a positive into a negative reaction varies with each patient and should be controlled by the blood examination. A positive reaction in tubes is little affected by treatment.

5. If mercury is used, systematic and consecutive treatment is necessary and this should be begun early, and during the first year no lapses of over one month permitted or much that has gained will be lost.

6. The use of alcohol has a bad effect on syphilis, predisposing to secondary and later manifestations, and, to a certain extent, nullifying the treatment.

7. In patients recently treated by mercury, and giving a positive reaction, the time required for a negative Wassermann with salvarsan may be much reduced. In very recent cases, or older cases not under the influence of mercury, there may be no shortening of the time.

8. In cases of a mercury-produced Wassermann negative at the time of injection, the reaction will generally revert to $+$ $-$, or in some cases to $+$, and then at varying intervals go back to $-$, from which subsequent injections may not change it. This is suggested as a possible diagnostic procedure in cases in which the question of a cure is open to doubt and it is not desired to suspend treatment if the infection still persists.

9. The intravenous method is preferred to others as being easier, more efficacious, less painful and less time being required by the patient.

10. At least two injections should be given and more if indicated and these injections should be given about a month apart.

11. An otherwise healthy adult will take from 0.5 to 0.6 gm. intravenously without untoward effect.

12. Unless one is prepared to maintain a very thorough watch over the patient, the salvarsan treatment should be reinforced with mercurial medication, and in any event better results will be obtained thereby.

13. In syphilis the tonic effect of salvarsan is far beyond that of any other drug at our disposal. Its effect on the weight, strength, appetite, mental and general condition approach the marvelous.

14. Whether or not salvarsan will cure syphilis remains to be seen, but it must be admitted that it will do some things that mercury will not do, and therefore must have a place in our therapeutics, most especially in those cases in which mercury is not well tolerated.

15. Up to the time of this report there had been no recurrences of manifestations, but in those under observation for seven months the reaction has returned to $+$ or $+$ $-$ in 22 per cent. In those under observation for four months 10 per cent. of the reactions have lapsed as before stated.

The Open Field of Internal Medicine.—In its rational development internal medicine is the laggard among the medical sciences, having always been centuries behind such branches as surgery and anatomy. This is natural, since the data of the latter are superficial and obvious, while those of internal medicine are deep-seated, obscure and inaccessible, and allow much play for theory or imagination. Moreover, internal medicine lacks the spectacular appeal which is expected, for example, by surgery. Yet though its data are inaccessible, its problems difficult, its therapeutic possibilities limited, the greater the difficulties the greater is the glory of surmounting them, and it is evident that internal medicine has at last entered on a career in honor and efficiency second to none.—Nichols in *Popular Science Monthly*.

EIGHT MONTHS' EXPERIENCE WITH SALVARSAN AT THE NEW YORK SKIN AND CANCER HOSPITAL *

HOWARD FOX, M.D.

AND

WILLIAM B. TRIMBLE, M.D.
NEW YORK

A few months ago we expressed some opinions on the value of salvarsan, based in part on our own experience and in part on that of others. Our present communication is a detailed account of work that has been done during the past eight months, and the conclusions drawn are based entirely on our own experience.

The greater part of our observations have been made at the New York Skin and Cancer Hospital in the service of Dr. George Henry Fox. At the time of writing, 128 injections have been given to 100 selected patients. The first ten patients were treated by the subcutaneous method (Wechselmann), after which intramuscular injections of an alkaline solution (Lesser method) were tried in twenty cases. Later, the intravenous method was employed in sixty-nine cases and intramuscular injections of an oily suspension in twenty-nine cases. The treatment was repeated in twenty-one of our patients, three of them receiving three, and two receiving four separate injections.

We have been fortunate in being able to follow a large proportion of the cases both clinically and serologically. Shortly before completing our report, we had personally observed or received letters from eighty-five of our patients. Of the remaining fifteen cases, ten had been under observation for at least a month.

Three patients with primary lesions were treated before the appearance of any general manifestations. They were unfortunately not seen at an early enough period to attempt to abort a general infection, as at the time of treatment the Wassermann reaction had already become positive.

Our early opinions regarding some of the cutaneous lesions of syphilis have not materially changed as a result of further experience. The macular syphilids have not disappeared with any surprising rapidity, and the papular syphilids have generally proved pretty resistant to treatment. Gummatous lesions have generally yielded to salvarsan in a most satisfactory manner, while mucous patches and condylomata have regularly disappeared in a most surprisingly short time.

In four cases (68, 74, 58 and 77) of late syphilis of the palms and soles, lesions which are notoriously resistant to mercury, the results of single injections of salvarsan have been extremely gratifying. One of these patients (68) had suffered from an extensive syphilid of the sole for twelve years, in spite of a good deal of treatment with mercury by several competent syphilologists. One month after a single intramuscular injection, the eruption had completely disappeared. In the other cases, the lesions had existed for three years, two years and two months respectively and disappeared in an equally satisfactory manner. Indeed, in one case a verrucous syphilid of the palms entirely cleared up in one week after an intravenous injection.

* Read in the Section on Pharmacology and Therapeutics of the American Medical Association, at the Sixty-Second Annual Session, held at Los Angeles, June, 1911.

* Owing to lack of space, several pages of tables detailing the cases are here omitted. They are included in the Transactions of the Section and in the authors' reprints. A copy of the latter will be sent by the authors or by THE JOURNAL on receipt of a stamped, directed envelope.

Four of the patients suffering from early syphilis presented an iritis which disappeared at the end of three weeks. In one case the iritis became more severe after treatment and on the fourth day had involved the other eye. A most excellent result was obtained in a case (57) of syphilitic neuroretinitis in a patient who had been infected a year and a half previously. Before injection her vision (examined by Dr. H. H. Tyson) was 20/100 in the right and 20/70 in the left eye. At the end of sixteen days there was a decided clearing of the fundus and she could read 20/30 minus with both eyes. One week later the fundus was practically normal in appearance and the vision was 20/20 minus in the right and 20/30 minus in the left eye.

Headaches occurring in both the early and late stages of syphilis were generally very favorably affected by salvarsan. Pains accompanying early manifestations,

istration that we have used. Three patients relapsed after one injection and four cases after two injections had been given. Three patients relapsed or remained stationary after the first, but responded favorably to a second injection. The patients in whom failures, or at least unfavorable results were obtained, included the case of tabes and a case (16) in which the diagnosis was in some doubt (gummatous vs. varicose ulcers). In addition, a case of gummatous ulceration of the leg showed no response to treatment at the end of three weeks, while the results in three others (38, 43 and 87) must be considered unsatisfactory. In all there were fifteen cases in which either relapse or unsatisfactory results were observed.

Many of the patients in whom excellent results were obtained by salvarsan had been previously treated by mercury without success. It is probable, however, that

Fig. 1.—Tuberculo-squamous syphilid of twelve years' duration.

Fig. 2.—Same as Fig. 1, one month after one intramuscular injection.

those associated with gummatous lesions and especially with periostitis, also yielded very satisfactorily.

One patient suffering from incipient tabes was treated without any benefit. In one case of hereditary syphilis the result was excellent, in another moderately good. In a case of leukoplakia of the mouth no improvement was obtained. In a case of extensive leukoplakia of the tongue, there was a temporary lessening in the thickness of the patches, without any diminution in their extent.

The all-important question of the permanence of the action of salvarsan can hardly be fairly discussed until a period of several years at least has elapsed. It must be said, however, that the number of relapses in our cases has been somewhat less than would have been expected as a result of our earlier experience. One of our patients relapsed repeatedly after four different injections, comprising all of the four methods of admin-

most of these cases at least would have yielded in a satisfactory manner to mercury if its administration had been sufficiently vigorous and thorough. An example of the very rare cases in which mercury cannot be tolerated in any form was shown by a case (83) of extensive gummatous ulceration of the palate. For eighteen months an experienced syphilologist had attempted to treat this patient by injections, inunctions and internal administration of mercury. Small doses had proved of no avail and large doses had been followed by necrosis of portions of the gums. We spoke of this case in our first communication¹ as one in which a "disappointing relapse" had occurred; we now report that the ulcerations healed entirely after a second injection and that no further relapse had occurred at the end of seven months.

1. Fox and Trimble: THE JOURNAL A. M. A., March 4, 1911, p. 650.

The Wassermann test was made in every case before treatment and subsequently as often as circumstances permitted. More than 400 examinations were made in the course of eight months. The original Wassermann method was followed, using as antigen a reliable alcoholic extract of syphilitic liver.

The results of the Wassermann reaction may be conveniently considered in two groups, those of patients who received one and those of patients who received two or more injections. Of the patients who were treated by a single injection, seventy-one showed a positive Wassermann reaction before treatment. Of these cases, twenty-eight became negative at an average period of about six weeks. In these the reaction later became positive in two, weakly positive in three cases and faintly positive in one case. The twenty-two cases which remained negative (30 per cent.) were followed, on an average, two and a half months; some of them for six and eight months. Of the remaining forty-three cases, ten showed a temporary and twenty a final weakening in the reaction.

Fig. 3.—Gummatous ulceration of six months' duration.

In eleven cases the reaction remained the same. In two cases it became stronger.

Of the cases in which two or more injections were given, twenty were positive before treatment. Of these four became negative in periods varying from five to nine weeks. One of these later again became positive. Of the remaining fifteen cases five showed a temporary and eight a final weakening in the reaction, while eight remained the same. The total number of cases which remained negative after either single or multiple injections was twenty-five (27 per cent.).

In all except five of the cases ophthalmoscopic examination of the fundus was made, although we have for some time considered that such an examination was an unnecessary precaution. Routine examinations of the urine were made before and after injection and showed a transitory albuminuria in a few cases.

At the beginning of our experiments every patient was required to remain in bed for three days after treatment. More recently the patients receiving intramuscular injections of oily suspension have been allowed to go home after treatment and advised to keep as quiet as possible for several days. Patients receiving intravenous injections have been required to remain in bed only for twelve, or in some cases four or five hours.

There have fortunately been no serious ill effects from salvarsan up to the time of writing. Deafness appeared in one case three months after treatment, but was found on aural examination to be due to acute catarrhal otitis media. The patient presented a chronic rhinitis and pharyngitis and had also suffered from a discharge from the opposite ear before the administration of salvarsan. Another patient (68) who had complained of smarting of the eyes and blurring of vision, showed no abnormal changes in the fundus on ophthalmoscopic examination.

Among the general effects of salvarsan should be mentioned a marked improvement in general health that was

Fig. 4.—The lesions shown in Figure 3, view ten days after one intramuscular injection.

noted in quite a number of cases. Seventeen of our patients showed a decided gain in weight, one of them gaining 23 pounds in the course of seven months. In three cases, however, there was a loss of from 15 to 18 pounds, which had not been regained in two of the cases at the end of three months.

After subcutaneous and intramuscular (Lesser) injections the temperature rose as a rule on the second day and lasted from one to three days. In five cases there was no reaction whatever, while in the others the height of the temperature ranged from 99 to 102.6. After intravenous injections the temperature rose as a rule within a few hours, its height varying from 99 to 103 degrees.

No temperatures were taken after most of the oily injections, the patients being allowed to go home immediately after the treatment.

Among the general symptoms noted after intravenous injections were chills or chilly sensations in twenty-seven cases, vomiting in twenty-five, headaches in fourteen cases. One patient complained of dizziness, another of diarrhea for a day, and two suffered from sleeplessness for several days. In twenty-eight cases there were no symptoms whatever. Two patients who had shown a marked reaction after the first injection showed no symptoms after the second had been given.

The local effects of intramuscular injections of alkaline suspensions included pain, induration and lameness, which were very severe in nine cases, moderate in five and slight in five cases. The discomfort in some of the cases was severe enough to warrant the use of morphin. After intramuscular injections of oily suspensions the local effects were severe in ten, moderate in eleven and slight in eight cases. In the worst cases the patients were obliged to remain in bed for a week or more, and even at the end of several months they complained of some stiffness about the buttocks and thighs. Necrosis occurred in two cases (3 and 4) after subcutaneous injections, in one case (10) after the intramuscular injection of an alkaline solution, and in two cases (70 and 88) after intramuscular injections of oily suspensions. At the time of writing nine of the ten patients that had been treated by the subcutaneous method still presented the characteristic indurations between the scapulae; the tenth case had been early lost to observation.

Among the interesting by-effects that were observed may be mentioned a case (85) in which a herpes zoster appeared on the trunk ten days after the injection. A general toxic erythema occurred in two cases (41 and 68) on the seventh and ninth days respectively after treatment, lasting three days and accompanied by febrile symptoms. In one case the temperature rose on the first day to 104.

Treatment with salvarsan had no disturbing effect on pregnancy in one of our cases (21) which was followed from the sixth month to full term, and in another case which could only be followed for a month.

Disturbances in menstruation were noted in twelve cases after treatment. In two of the cases the irregularities were possibly due to the menopause. In another case the cessation of the flow was probably due to pregnancy. In one case the menstrual periods became more regular, while in the others they became more irregular after injection, being delayed in one case for seven and a half months.

One patient with nodular leprosy was experimentally treated with an intramuscular injection. In our first report we stated that this patient had been prostrated by the treatment and that no benefit had resulted from the injection. A change occurred, however, a month later when numerous ulcerating lesions of the face healed entirely. Two weeks later there was a partial relapse. No change in the solid nodular lesions was noted. A small dose of salvarsan was later given intravenously without producing any improvement or indeed any reaction whatsoever.

Two patients (17 and 90) who were treated for syphilis also presented complicating lesions of psoriasis. In one case the eruption was practically well at the end of three and a half months. In the other case, in which (according to the patient's statement) the eruption had existed for forty years, there was a complete disappearance of the psoriatic patches at the end of six weeks and at the end of three months no relapse had occurred.

In giving intravenous injections we have employed the gravity method in all of our cases except the first, in

which the more complicated Schreiber apparatus was tried. We have had practically no difficulty and no bad results. For most of the injections we have used our own model of the gravity apparatus described elsewhere.² In this communication we recommended the Schreiber form of needle having a 17 or 18 bore and a bevel that was not too long. Our experience with some of the short beveled and dull needles has led us to modify our views. We are now convinced that the most suitable needle is of the Schreiber type, having an 18 bore (Stubbs) and a bevel of about 3 or 4 mm., that is, not too sharp and not too dull. For diluting the salvarsan we use only sterile distilled water and dispense entirely with the salt solution. This may be the cause of the appearance of small nodules in the vein in eight out of sixty-nine injections. In two cases they were present at the last time of observation, the fourth and tenth weeks respectively after treatment. In the others they had disappeared in from one to four weeks. The simplicity of using water instead of salt solution is an advantage that in our opinion more than compensates for the appearance of some of these harmless nodules.

CONCLUSIONS

1. Salvarsan is a powerful symptomatic remedy for the treatment of syphilis.
2. It acts with greatest rapidity on lesions of the mucous membranes.
3. It is of decided value in obstinate palmar and plantar syphilids.
4. The permanence of its action cannot be determined until a number of years have elapsed.
5. Salvarsan should be used in conjunction with mercury and cannot entirely replace this valuable remedy, except in a few selected cases.
6. The effect of salvarsan on the Wassermann reaction is less favorable than on the clinical manifestations of syphilis.
7. Its effect on the Wassermann reaction is in general analogous to that of mercury.
8. The intramuscular and intravenous methods of administration are probably of equal efficiency.
9. The intravenous method is the one of choice where rapidity of action and the comfort of the patient are factors to be considered.

For the greater part of the material we are indebted to Dr. George Henry Fox. We also wish to express our thanks to Dr. Robert G. Reese for the ophthalmoscopic examinations and to Dr. George T. Jackson and Dr. John A. Fordyce, from whose clinics a few of the cases were obtained.

616 Madison Avenue—39 East Twenty-Eighth Street.

ABSTRACT OF DISCUSSION

ON PAPERS OF DRS. HIRSCHFELDER, FOX AND TRIMBLE, AND MATSON AND REASONER

DR. D. W. MONTGOMERY, San Francisco: The interesting statement made by Dr. Hirschfelder in regard to the toxic effect of salvarsan being in many cases due to decomposition after intramuscular or subcutaneous injections, is very important, and constitutes a weighty reason for giving it intravenously. We do not know what these decompositions are, but we do know that there is an enormous amount of arsenic put under the skin or into the muscles, and we also know that any decomposition would likely work detrimentally to the patient.

The statement, however, in regard to cephalic chancre being particularly apt to be followed by syphilis of the brain, I am

2. Fox and Trimble: Med. Rec., March 18, 1911, lxxix, p. 400.

not inclined to accept. I do not believe that chancres in this region are any more likely to be followed by syphilis of the brain than infection in other regions. Another interesting statement was made in regard to the milk of syphilitic mothers suckling syphilitic infants being curative of the syphilis in the infant. That is true. Why it is so is hard to tell. It is doubtful if the small amount of arsenic found in the maternal milk is enough to account for it. In testing for arsenic one must always bear in mind that it can be found anywhere. I suppose there is not a suit of clothes in this room that does not contain it. So that the mere finding of arsenic in the milk, except in large quantity, would not account for the curative action of salvarsan in the infant. I saw an infant in Frankfort a short time ago under Theodore Sachs. It was getting well without receiving any arsenic or salvarsan, or any other drug directly, but simply from suckling a mother who had received a gluteal injection of salvarsan. I have no personal experience with salvarsan in anything else than syphilis, and I have never seen it given in anything else excepting in a few cases of leprosy. One of these lepers I understand did excellently. Dr. Fox, I think, is correct in saying that the macular and papular syphilids do not disappear with very astonishing rapidity. Such eruptions do disappear rapidly with mercury and I do not know that they fade away any more rapidly with salvarsan. But gummata, when the patient is treated with salvarsan, often disappear with remarkable rapidity. The late palmar and plantar syphilids are particularly amenable to treatment with salvarsan, and it is a matter for congratulation that they do, as under the old modes of treatment they were among the most intractable eruptions, and we often had to resort to intense treatment, such as calomel injections. That salvarsan does just what one might expect, as arsenic is strongly epitheliotropic. The same can be said of leukoplakia, and it is very interesting that Dr. Fox found that it reduced the thickening in this intractable disease.

Dr. H. R. OLLIVER, San Francisco: Some cases of optic atrophy which have shown deleterious effects from a first injection of salvarsan, improve and the patients often recover under a second injection. The first injection causes a round-cell infiltration, whereas the second injection causes reabsorption and the case is bettered. In regard to nephritis after the injection of salvarsan, I have seen a large number of cases in which there were casts and albumin occurring after the injection, but in those cases the casts invariably were waxy, and it would be impossible for the salvarsan to have caused. The lapse of one or two days after an injection is not sufficient to permit the formation of waxy casts in the urine. In the other cases in which there was transitory albuminuria there were no casts present. The crises in parasymphilitic cases are acted on remarkably. I have seen patients in agony, who in twenty-four hours were entirely relieved of their pain. It is well known that all infants that are born syphilitic should suckle a mother who has been injected with salvarsan or a syphilitic wet-nurse who has been treated. In addition to this it is well to use about 0.015 of salvarsan every eight days as an intravenous injection in these cases, because if we use larger doses it may liberate so many spirochetes from the internal organs of these infants (which literally teem with these organisms) as to cause an overwhelming of the economy and deleterious results. In leprosy the cases which have been improved by the salvarsan have invariably shown a positive Wassermann reaction. In primary syphilis I have never been so fortunate as to get a Wassermann reaction before the tenth to the sixteenth day after the initial lesion. I cannot really conceive how we would get it until we get a systemic infection; that is, we would not have enough antibodies formed in so short a time as to be able to get a Wassermann reaction. I have seen one case of malignant syphilis in which I have done three or four Wassermann reactions with negative results. The patient became very emaciated, was then given salvarsan and in six days his blood gave a triple plus Wassermann reaction and he has rapidly improved. The time of the disappearance of the reaction after the administration of salvarsan corresponds quite

directly to the strength of the Wassermann reaction before the salvarsan is given. That is, if the patient has a triple plus Wassermann, as we understand it, before the injection, it takes that patient from two to three months to become negative. The patient with double plus before the injection takes one or two months. When he has single plus or partial Wassermann, it takes about one month to become negative. The reaction is reverse to that of mercury. With the mercurial treatment we may have a negative reaction caused by mercury without the patient being well, and if we take him off of mercury he gradually increases his reaction until it becomes thoroughly positive again, whereas with salvarsan it may become positive, and gradually comes down until it becomes negative.

Dr. M. H. FUSSELL, Philadelphia: To the ordinary everyday clinician the administration of so large an amount of liquid into a vein is rather a terrible affair, but it must be, and in my own experience has been, practically without any danger. Yet there are dangers in these intravenous injections. I think Dr. Hirschfelder spoke of two to which I wish to call particular attention. It is true that these accidents ought not to occur. That they do occur I think proves that we should be extremely careful either in perfecting our own technic, if we ourselves administer the salvarsan intravenously, or that we should have an expert to do it for us. The latter plan I always follow. However, I have seen both of the accidents occur that Dr. Hirschfelder spoke of—one the induration about the point of injection into the vein. In this particular instance there was not only an induration about the vein but there was a distinct and marked phlebitis of the entire vein; the man's arm was swollen to twice or three times its normal size and for three or four weeks he was extremely ill. That was an accident which should not occur, but it did occur as a direct result of the injection. If it had been given by an expert I presume it would not have occurred. Another one, a frightful affair, is the hemiplegia or sudden death after injection. There is one case in Philadelphia, which did not occur in my own work, in which a man had a sudden collapse, a hemiplegia and finally a recovery. It seems to me some blood must have been in the tube before the injection of salvarsan was begun. This can easily be prevented by allowing the solution to flow through the tube at the time it is injected into the vein. I saw such an accident occur in the hands of a resident of mine after an ordinary phlebotomy. I had told him to give an intravenous saline solution. He bled the man with the tube through which he was to inject the saline solution. The man was bleeding at the time he attached his tube, and instantly the man was dead. I presume he had a cerebral thrombosis. That these things can occur is certain and we should remember that it is no light job to put numbers of centimeters of liquids into a vein. I want to call attention to the apparatus Dr. MacKinney of Philadelphia uses for the purpose. His tube is in a sort of vacuum bottle and it retains the heat very well. I would like to ask Dr. Fox if he has any explanation for the disappearance of the psoriasis and if he has used salvarsan in psoriasis alone.

Dr. L. DUNCAN BULKLEY, New York: I have watched many cases in my service in the New York Skin and Cancer Hospital with a great deal of interest, and photographs have confirmed the remarkable results in some cases which I have not observed personally. Really it is a wonderful remedy, but the point is whether for practicing physicians this is going to take the place of our ordinary treatment, what its true value is in relation to other proper therapeutics. I endorse what has been said concerning the Wassermann test. The Wassermann test should be used wherever possible; in doubtful cases it is certainly invaluable as an index to treatment. I think it might well be made a legal requirement before marriage. It would be of immense value in stamping out the disease if every patient who has had syphilis were required to have his cure demonstrated by the repeated application of the Wassermann test.

I have heard a great many other discussions on salvarsan in various scientific societies, with the exhibition of many dozens of cases in process of cure, and have seen bad results from subcutaneous and muscular injections. I have seen one

case of terrific phlebitis from an intravenous injection, which required a surgical operation for removal of the vein. The patient was not under my care at the time but came to me afterward. Almost all who write or speak say, evidently with good reason, that some mercurial treatment must follow the use of salvarsan. I have not accepted the salvarsan treatment as the best in private practice, but I am waiting to know further results. As yet, with proper other treatment, I do not see the need of salvarsan in private cases. I have used it in hospital practice and get surprising results, but I do not advise my young men to rush into it. It may be peculiarly serviceable in certain cases in future, but I doubt if it will ever largely supplant other accepted treatment. It should be used by an expert. There is also a grave danger in regard to the salvarsan treatment, lest patients think that they have been cured of syphilis by one or two injections and then neglect treatment and have relapses of a serious nature. The quacks have taken it up and are charging large amounts for injections of something called salvarsan. The victims have it administered and then go away and think they are cured. If they get the real article from a regular physician there is still danger of relapses or, rather, later manifestations of the disease.

Dr. W. P. MURRAY, Caliente, Nev.: It seems that there is quite a scare in the minds of many lest something foreign get into the veins during the administration of salvarsan. As to the blood regurgitating back into the tube, I believe the point is well taken by the speaker who preceded me. Many physicians have a fear of allowing a few bubbles of air to gain access to the system. I have seen as much as 1 to 3 c.c. of atmospheric air gain access to the venous system without causing ill effect. Never in my practice have I seen ill effect from a few bubbles of air in the administration of any solution whatever. Again, in reference to the normal salt solution, I find that the only use we can make of it is to ascertain whether we have the needle in the vein or not. I suppose some better physicians than I still resort to cutting the vein and inserting the needle and then allowing the salvarsan to flow into the vein. I have found it entirely unnecessary, as, by having the patient clench the fist so that the veins stand out, it is a very simple matter to pick up the vein. The only use of normal salt solution is to tell whether or not we are in the vein. When we have a sharp needle and insert it obliquely and release the flow from the normal salt solution, and it begins to balloon up the solution is not entering the vein, but if it flows along and there is no ballooning or swelling it is in the vein. To do that is sometimes a difficult matter. In the administration of salvarsan I have yet the first time to see the fluid escape extravascularly without producing a slough. I have known it to occur after the most careful aseptic precautions. In reference to the intramuscular administration, I have my first case yet to see in which it did not produce inflammation and sloughing. Never, when proper precaution is taken as to sepsis, have I seen a slough after the intravenous method.

Dr. J. O. HIRSCHFELDER, San Francisco: In reply to Dr. Montgomery's criticism of the presumed presence of arsenic in the milk and the claim that he makes that arsenic is found everywhere, I would say that the Homeopathic Association might consider such presence, but I doubt whether one will be able to find arsenic in milk in demonstrable quantities, provided one uses arsenic-free tests in the examination, unless one uses enormous quantities of milk—probably 2 or 3 liters. I cannot speak from knowledge that these large quantities would show even traces of arsenic. In quantities like 100 c.c. I do not think that arsenic has ever been found in normal milk. In the human milk of syphilitic individuals who have received salvarsan the arsenic has been found in very appreciable quantities, so that in making the test the presence of arsenic was very plain indeed; likewise with goats which had received a small quantity of salvarsan. Then we are not to forget in speaking of the milk of syphilitics who have received salvarsan, that it contains large quantities of endotoxins, as well as antibodies that have been set free. The fact that, in the experience of some, syphilitic infants that have been treated with this milk have become violently ill is probably

due to the immense quantity of endotoxins that are set free and cannot be formed into antibodies in the body of the infant. Consequently, I think that we should not forget the advice that was given to administer salvarsan to cows and goats and to feed the infants with the milk derived from these animals. I want to thank Dr. Fussell for the suggestion that he has made in regard to the coagulation of the blood in the tube inserted into the vein and the possible production of emboli by coaguli. I am sure that can occur. If the simple precaution is taken to allow the blood to flow freely through the tube before the salvarsan is introduced I think there will be no difficulty. I have had no trouble in making the salvarsan injections. In answer to Dr. Murray, as well, I would state that in very fat people it is sometimes impossible to find the vein at the elbow no matter how thoroughly you may apply your tourniquet, without having bare the vein by a little incision, but in those cases I have always taken the vein at the wrist where it always stands out prominently. There is rarely sufficient accumulation of fat at this point to interfere with the finding of the vein and the introduction of the needle. In answer to Dr. Fussell, if 300 c.c. are allowed to flow into the tube in fifteen minutes that would be 0.33 c.c. per second. This quantity is rapidly disposed of by the circulating blood. In order to be cautious, I have my mixture heated to 110 F., but I think I do it out of respect to the figure rather than as the result of experience of any ill effects from using the solution at a lower temperature.

Dr. HOWARD FOX, New York: The fluctuations in the Wassermann reaction occurring after injections of salvarsan cannot, I think, be ascribed to faulty technique. Such changes are shown in a marked degree in a table published by Zieler. They were also noted in some of our own cases, the reaction changing from positive to negative, or vice versa, within a week. They are certainly not seen with such frequency after mercurial treatment.

Dr. MATSON spoke of the intravenous injections as being more efficient than the intramuscular injections of salvarsan. While the intravenous method is undoubtedly quicker, we are not prepared to say that its action is more efficient than the intramuscular method. This is a question of greatest importance, as it is plainly impossible to administer salvarsan in every case by the intravenous method. The drug cannot fulfill its greatest usefulness, unless it can be given in office and dispensary practice, just as injections of salicylate of mercury are given.

In reply to Dr. Fussell, I would say there are three signs by which infiltration of the tissues can be avoided when giving intravenous injections. The most important of these is pain at the point of injection, the patient being instructed to tell the moment he feels a sharp, stinging pain at the point of the insertion of the needle. It is therefore apparent why salvarsan should never be administered intravenously under a general anesthetic. In the second place, the nurse or assistant should see that the fluid is running steadily from the glass container, and finally, the operator should watch for the presence of swelling at the insertion of the needle. This is not as important as the other signs, because the infiltration may take place behind the needle and not be visible at once to the operator. It may be said that a very small amount of infiltration in the cellular tissue will do no harm.

In answer to Dr. Fussell's question about psoriasis I would say that I have treated only two cases, the psoriatic lesions in these cases being incidentally present. The action on psoriasis is probably due to the arsenic, which has a stimulating effect on the epithelial layers of the skin.

I agree with Dr. Bulkley, who warns against our depending entirely on salvarsan. It should merely be considered as one of our means of combating a very serious disease and should be used in combination with the older well-known remedies. I also think that Dr. Murray is right regarding the bogey of getting air bubbles into the vein. I have seen small air bubbles enter the vein without producing the slightest ill effects.

Dr. RAY W. MATSON, Portland, Ore.: I believe for the initial cases that the intravenous method of administration

would accomplish more nearly than any other the so-called *therapia sterilisans magna* of Ehrlich, since the salvarsan is taken up by the vascular system and permeates the whole body in the course of a few seconds or minutes. There is a possibility of destroying all of the spirochetes contained in the body in a very short period of time. For subsequent injections I think the intramuscular is theoretically better because the drug is absorbed and eliminated very slowly. In this way it exerts a continuous action. First, we have what might be termed the fatal blow, followed by the continuous action of the chemical. As to our method of giving salvarsan, of course it must be remembered that our patients have been in excellent shape to begin with, men in the prime of life. I have had absolutely no difficulty in regard to reaction. One patient with myocarditis complained of excruciating pain while the injection was being given, and it was so severe that I had to discontinue the method after giving the third injection. Captain Pearson of the Army Medical Corps devised the scheme of applying the tourniquet and distending the veins after freezing the vein with carbon dioxide. The introduction of the needle is very much easier because the vein is fixed absolutely to the skin. Regarding the length of time required for converting a negative to a positive reaction following the administration of salvarsan, I cannot agree with Dr. Oliver in every respect. I have seen the strongest positive Wassermann converted to a negative one in the space of three weeks following the administration of salvarsan. In one case of latent syphilis of twenty years' duration in which the patient complained of severe headache of one and one-half years' duration, he was relieved in two weeks and his Wassermann became negative at the end of three weeks following the intramuscular administration of 0.6 gm. of salvarsan.

A NEW ESOPHAGOSCOPE

A PRELIMINARY COMMUNICATION

RICHARD LEWISOHN, M.D.

NEW YORK

The principle on which all esophagoscopes have been constructed up to the present time has been seriously defective, as no attempt has been made to adapt the instrument to the anatomic conditions in the upper part of the alimentary tract. In the normal position, the oral cavity joins the upper esophagus at a right angle, and the only method hitherto in vogue to overcome this rectangular junction has been to extend the head forcibly and thus to approximate a straight line from the teeth to the cardia. This feature applies both to the simple straight tube and to the jointed instrument, and is the explanation of the unpopularity of esophagoscopy.

I have constructed an instrument intended to overcome this objection. As the accompanying illustrations show, my instrument is built on a rectangular plan similar to the gastroscope models of Poirier and Eckstein, although it differs from these in one essential particular. Their esophagus tube was a jointed one, while mine is a telescope formed by six small tubes which fit into one another. Each tube is 5 cm. long, and the diameter varies from 9 to 14 mm. The image is transferred at the rectangular junction by a mirror, while another perforated mirror in the eye-piece is used for the purposes of light reflection (Fig. 1).

It is beyond the scope of this preliminary report to give a detailed description of the minute mechanism of

the instrument. I would like to emphasize, however, that the construction is such that the entire instrument can be taken apart for the purposes of thorough cleansing.

The instrument is introduced with the telescope closed (Fig. 2). Then, by means of a spring, under the guidance of the eye (Fig. 1), the telescopic tubes are separated and thus pushed down into the esophagus. As no pressure is made on the tubes until they lie in the axis of the esophagus, the danger of in-

Fig. 1.—Diagram of esophagoscope, showing tube extended.

juring the posterior wall of the organ, which is ever present in all the other models, is entirely obviated.

I have applied the same principle of a rectangular telescope to the construction of a gastrosopic instrument, by simply adding another tube and inserting a

Fig. 2.—General view of esophagoscope with telescoping tube closed.

cystoscopic optical and illuminating system. For the time being, however, I have confined my practical tests to the esophagoscope.

I have tried the esophagoscope in a considerable number of cases. After some experience with the instrument, the introduction should be easily learned. The instrument is inserted with the patient in an upright sitting posture. Cocainization of the pharynx is advisable, but I have already introduced the instrument without even local anesthesia. The introduction